

Correction

Open Access

Correction: The burden of diarrhoea, shigellosis, and cholera in North Jakarta, Indonesia: findings from 24 months surveillance

Magdarina D Agtini*¹, Rooswanti Soeharno¹, Murad Lesmana², Narain H Punjabi², Cyrus Simanjuntak², Ferry Wangsasaputra^{1,2}, Dazwir Nurdin², Sri Pandam Pulungsih³, Ainur Rofiq¹, Hari Santoso⁴, H Pujarwoto¹, Agus Sjahrurachman⁵, Pratiwi Sudarmono⁵, Lorenz von Seidlein⁶, Jacqueline L Deen⁶, Mohammad Ali⁶, Hyejon Lee⁶, Deok Ryun Kim⁶, Oakpil Han⁶, Jin Kyung Park⁶, Agus Suwandono¹, Ingerani¹, Buhari A Oyoyo², James R Campbell², H James Beecham², Andrew L Corwin² and John D Clemens^{6,7}

Address: ¹National Institute of Health Research and Development, Jakarta Indonesia, Ministry of Health, Jakarta, Indonesia, ²United States Navy Medical Research Unit 2, Jakarta, Indonesia, ³Infectious Disease Hospital Sulianti Saroso, Ministry of Health, Jakarta, Indonesia, ⁴Communicable Disease Control and Environmental Health, Ministry of Health, Jakarta, Indonesia, ⁵Department of Microbiology, University of Indonesia, Jakarta, Indonesia, ⁶International Vaccine Institute, Seoul, Korea and ⁷National Institute of Child Health and Human Development, Bethesda, Maryland, USA

Email: Magdarina D Agtini* - magdarina@yahoo.com; Rooswanti Soeharno - rooswanti@litbang.depkes.go.id; Murad Lesmana - murad@namru2.org; Narain H Punjabi - narain@namru2.org; Cyrus Simanjuntak - cyrushs@hotmail.com; Ferry Wangsasaputra - ferry@namru2.med.navy.mil; Dazwir Nurdin - ferry@namru2.med.navy.mil; Sri Pandam Pulungsih - rspiss@idola.net.id; Ainur Rofiq - rofiq@idola.net.id; Hari Santoso - santoso@hotmail.com; H Pujarwoto - ferry@hotmail.com; Agus Sjahrurachman - Sjahrurachman@idola.net.id; Pratiwi Sudarmono - ferry@namru2.med.navy.mil; Lorenz von Seidlein - lseidlein@ivi.int; Jacqueline L Deen - jdeen@ivi.int; Mohammad Ali - mali@ivi.int; Hyejon Lee - hjee@ivi.int; Deok Ryun Kim - drkim@ivi.int; Oakpil Han - qphan@ivi.int; Jin Kyung Park - jkpark@ivi.int; Agus Suwandono - selitbang@litbang.depkes.go.id; Ingerani - ingerani@hotmail.com; Buhari A Oyoyo - oyofoba@namru2.med.navy.mil; James R Campbell - campbelljr@namru2.med.navy.mil; H James Beecham - beechamjh@namru2.med.navy.mil; Andrew L Corwin - corwinal@namru2.med.navy.mil; John D Clemens - jclemens@ivi.int

* Corresponding author

Published: 26 January 2007

Received: 10 October 2006

BMC Infectious Diseases 2007, 7:1 doi:10.1186/1471-2334-7-1

Accepted: 26 January 2007

This article is available from: <http://www.biomedcentral.com/1471-2334/7/1>

© 2007 Agtini et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This is a correction of an earlier published article.

Text

After the publication of this work[1], we became aware that our description of the methods used to identify

Shigella spp. did not describe the identification of a subset of *Shigella* spp. which were untypeable with commercial sera. Additional typing was conducted by Dr. Kaisar A.

Table 1: *S. flexneri* serotyping criteria using MASF antibodies

Serotype	Type antigen specific					MASF B	Group antigen specific			Provisional antigen specific	
	I (I:-) ^a	II (II:-)	IV-2 (IV:-)	V (V:-)	VI (VI:-)	B (-:I)	Y-5 (-:3.4)	6 (-:6)	7.8 (-:7.8)	IV-1 (-:EI037)	Ic
1a	+					+	+				
1b	+					+		+			
1c						+					+
2a		+				+	+				
2b		+				+			+		
3a						+		+	+		
3b						+	+	+			
3c						+		+			
4:-			+			+					+
4a			+			+	+				+/-
4b			+			+		+			
5a				+		+	+				
5b				+		+			+		
6					+	+					+/-
X						+			+		+/-
Y						+	+				+/-
4X						+					+

^a Shown in parentheses is the antigen recognized.

+ Positive reaction

Talukder and coworkers (Laboratory Sciences Division, ICDDR, B: Centre for Health and Population Research, GPO Box-128, Dhaka-1000, Bangladesh. Email:kai-sar@icddr.org.)

Brief description of the methods used for serotyping the untypeable *S. flexneri* strains using MASF

All *Shigella* strains which agglutinated with commercial polyvalent anti-*S. flexneri* serum (Denka Seiken, Tokyo, Japan) but were untypeable using commercial serotype specific antisera were examined using a panel of monoclonal antibodies against *S. flexneri* (MASF B) according to the typing scheme shown in Table 1 using monoclonal antibodies from one supplier (Reagensia AB, Stockholm, Sweden) with the exception of MASF 1c which was provided by Dr. N. I. A. Carlin (SBL, Sweden) [2-5]. Untypeable *S. flexneri* strains which agglutinated strongly with polyclonal MASF B and type antigen factor IV-2 but not with group factor Y-5 or 6 thereby excluding their characterization into serotype 4a or 4b were designated as serotype 4 if they agglutinated with a new antigenic determinant E1037 (IV-1). Untypeable strains which agglutinated with antibodies against the provisional antigen, MASF IV-1 specific for the new antigenic determinant E1037, were designated as 4X. Untypeable strains which agglutinated only with MASF 1c were designated as 1c.

References

1. Agtini MD, Soeharno R, Lesmana M, Punjabi NH, Simanjuntak C, Wangsasaputra F, Nurdin D, Pulungsih SP, Rofiq A, Santoso H, Pujarwoto H, Sjahrurachman A, Sudarmono P, von Seidlein L, Deen JL, Ali

M, Lee H, Kim DR, Han O, Park JK, Suwandono A, Ingerani, Oyofa BA, Campbell JR, Beecham HJ, Corwin AL, Clemens JD: **The burden of diarrhoea, shigellosis, and cholera in North Jakarta, Indonesia: findings from 24 months surveillance.** *BMC Infect Dis* 2005, **5**:89.

2. Carlin NI, Rahman M, Sack DA, Zaman A, Kay B, Lindberg AA: **Use of monoclonal antibodies to type *Shigella flexneri* in Bangladesh.** *J Clin Microbiol* 1989, **27**(6):1163-1166.
3. Talukder KA, Dutta DK, Safa A, Ansaruzzaman M, Hassan F, Alam K, Islam KM, Carlin NI, Nair GB, Sack DA: **Altering trends in the dominance of *Shigella flexneri* serotypes and emergence of serologically atypical *S. flexneri* strains in Dhaka, Bangladesh.** *J Clin Microbiol* 2001, **39**(10):3757-3759.
4. Talukder KA, Islam MA, Dutta DK, Hassan F, Safa A, Nair GB, Sack DA: **Phenotypic and genotypic characterization of serologically atypical strains of *Shigella flexneri* type 4 isolated in Dhaka, Bangladesh.** *J Clin Microbiol* 2002, **40**(7):2490-2497.
5. Talukder KA, Islam Z, Islam MA, Dutta DK, Safa A, Ansaruzzaman M, Faruque AS, Shahed SN, Nair GB, Sack DA: **Phenotypic and genotypic characterization of provisional serotype *Shigella flexneri* 1c and clonal relationships with 1a and 1b strains isolated in Bangladesh.** *J Clin Microbiol* 2003, **41**(1):110-117.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2334/7/1/prepub>