

Is intimate partner violence a risk factor for HIV infection? A systematic review and meta-analysis

Prof. Charlotte Watts Ph.D.

Research Director, STRIVE Research Programme Consortium

Head, Social and Mathematical Epidemiology Group

Department of Global Health and Development

London School of Hygiene & Tropical Medicine



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Debate about whether VAW responses a core part of HIV programming

- **Several large GBV-HIV initiatives**
 - UNAIDS Pillar for HIV prevention on addressing GBV
 - Large PEPFAR funding in Sub-Saharan Africa
 - UN Stop Rape Campaign
 - Inclusion of violence prevention activities in some sex worker HIV prevention programmes, including AVAHAN
- **However:**
 - Questions about whether interventions to address violence should form a core part of IPV programmes
 - Analysis of DHS data found no association between intimate partner violence (IPV) and HIV (Harling 2010)



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Aims & methods

Aims

- Compile existing epidemiological evidence on the association between exposure to intimate partner violence (IPV) and HIV infection

Methods

- Systematic review
- Searches of Pubmed, Embase, Cinahl, other databases until Dec 1 2010
- > 3,000 abstracts screened
- Inclusion: any population, any definition of IPV, HIV/STI
- Analysis stratified by study quality:
 - Prospective studies
 - High quality cross-sectional studies (biological outcome data, unexposed reference group)



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Results

- 35 papers, describing 41 datasets with 121,479 participants, reporting 115 estimates included
 - 5 prospective datasets
 - 3 large studies with biological outcomes
 - 2 HIV, 1 STI
 - 3 case-control datasets
 - 35 cross-sectional datasets
 - With biological outcome data AND unexposed reference groups
 - HIV: 12 datasets, 25 estimates
 - STI: 6 datasets, 6 estimates



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Prospective studies find associations


Study	Sample	Intimate partner violence measure	HIV/STI measure	Estimate
Jewkes et al	1099 women, vocational schools in rural Eastern Cape, South Africa, 2002	More than one episode of physical and/or sexual violence, WHO	Incident HIV, biologically confirmed, adjusted for HSV-2	aIRR=1.51 (1.04-2.21)
Weiss et al	1991 non-pregnant women aged 18-45, population registers of primary care centre Goa, India, 2001-2003	Physical violence, not further defined Sexual violence, 'the husband or partner forcing sex against the woman's wishes.'	Incident CT/GC/TV, biologically confirmed	aOR=1.40 (0.70-3.00) aOR=3.00 (1.20-7.50)
Zablotska et al.	3422 women aged 15-24, population-based Rakai, Uganda, 2001-2003	Sexual violence, "Sexual partner physically forced you to have sex when you did not want to.'	Incident HIV, biologically confirmed	1.6/ 100py in IPSV-, Alcohol- 2.2/ 100py in Alcohol+ 2.3/ 100py in IPSV+



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE





Different analyses of same cross-sectional data have different findings



Globally cross-sectional findings more mixed (HIV outcome)

FORM OF IPV CONSIDERED

PHYSICAL

Physical

Study	Odds Ratio (95% CI)	Total number of people in analysis
DHS ZIMBABWE 2005	1.35 (1.05, 1.73)	2287
NATIONAL FAMILY HEALTH SURVEY INDIA 2005	1.53 (0.76, 3.06)	28139
DUNKLE SOUTH AFRICA 2004	1.56 (1.21, 2.03)	800
Subtotal	1.45 (1.22, 1.73)	

SEXUAL

Sexual

DUNKLE SOUTH AFRICA 2004	0.88 (0.51, 1.53)	676
DHS ZIMBABWE 2005	0.95 (0.66, 1.37)	1852
Subtotal	0.93 (0.68, 1.26)	

PHYSICAL AND/OR SEXUAL

Either

DHS HAITI 2005	0.45 (0.23, 0.90)	2628
DHS LIBERIA 2006	0.87 (0.56, 1.35)	3278
DHS KENYA 2003	0.88 (0.62, 1.25)	1756
DHS ZAMBIA 2007	0.91 (0.77, 1.08)	3368
DHS RWANDA 2005	0.99 (0.59, 1.67)	2476
DHS MALAWI 2004	1.07 (0.81, 1.42)	2086
DHS MALI 2006	1.07 (0.51, 2.23)	2804
DHS ZIMBABWE 2005	1.11 (0.91, 1.34)	3203
DHS DOMINICAN REPUBLIC 2007	1.12 (0.67, 1.88)	7870
NATIONAL FAMILY HEALTH SURVEY INDIA 2005	1.35 (0.95, 1.90)	29783
DUNKLE SOUTH AFRICA 2004	1.49 (1.18, 1.89)	1336
Subtotal	1.05 (0.90, 1.21)	

PHYSICAL AND SEXUAL

Both

DHS ZIMBABWE 2005	1.38 (1.03, 1.85)	2061
DUNKLE SOUTH AFRICA 2004	1.66 (1.18, 2.32)	800
NATIONAL FAMILY HEALTH SURVEY INDIA 2005	3.92 (1.41, 10.94)	28139
Subtotal (I-squared = 48.9%, p = 0.141)	1.66 (1.17, 2.34)	

NOTE: Weights are from random effects analysis




Figure: Cross-sectional studies. Pooled OR, biological data only, HIV outcome, where reference group is no physical or sexual violence

Growing evidence globally that have a clustering of risk behaviours

- Men who are abusive to their partners are also more likely to have:
 - Concurrent sexual partners
 - A sexually transmitted infection
 - Problematic use of alcohol
 - Refuse to use a condom
- Clustering of risk linked to common underlying risk factors




STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Potential pathways of association between IPV & women's risk of HIV



Conclusions

- Violence is both a cause and consequence of HIV infection
- Prospective studies show an association between physical and/or sexual IPV and incident HIV in South Africa
- Prospective data also find association between sexual IPV & HIV in Uganda and sexual violence & STI in India
- Cross-sectional data analysis find less consistent findings
 - Many methodological factors make interpretation of existing evidence difficult
 - Consistent association between more severe IPV and HIV risk
- Unclear how generalisable findings are across different epidemic settings
- Pathways between IPV & HIV complex – need to be better understood to inform effective programmes



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



3 priorities to improve evidence base...

1. Identify opportunities to collect additional evidence from longitudinal studies

- Take advantage of opportunities within ongoing intervention trials with HIV outcomes

2. Make best use of DHS & other population data collection

- Ensure that DHS collects data on violence exposure from all partners
- Re-analysis of DHS to address methodological issues

3. Integrate questions on violence in HIV intervention research

- Provide deeper understanding of how violence and the fear of violence may undermine effectiveness of proven HIV interventions



STRIVE
Tackling the structural drivers of HIV

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Acknowledgements



Co-authors:

Karen Devries, Lori Heise, Claudia Garcia-Moreno, Joelle Mak

Funders:

- Sigrid Rausing Trust
- UK Economic and Social Research Council
- UK Department for International Development

**Copies of presentation at
strive.lshtm.ac.uk**



LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE

