

Supplementary Material Table 1. Literature on characteristics associated with LOS

Variable	Direction of association (Reference)
Admission-level characteristics	
Physical co-morbidities	Positive (Douzenis et al., 2012)
Diagnosis of psychosis	Positive (Chung et al., 2010; Huntley, Cho, Christman, & Csernansky, 1998; Lay, Lauber, & Rossler, 2006; Oiesvold et al., 1999; Pertile et al., 2011; Tulloch, Fearon, & David, 2011)
Co-morbid diagnosis of Substance Misuse	Negative (Compton, Craw, & Rudisch, 2006; Harman, Cuffel, & Kelleher, 2004; Huntley et al., 1998)
Legal status	Positive (Kallert, Glockner, & Schutzwohl, 2008; Pertile et al., 2011) Negative (Compton et al., 2006; Tulloch et al., 2011)
Social support	Positive (Oiesvold et al., 1999) Negative (Tulloch et al., 2011)
Prior service use	Positive (Huntley et al., 1998) Negative (Dausey, Rosenheck, & Lehman, 2002; Rothbard & Schinnar, 1996)
Deprivation	Positive (Abas, Vanderpyl, & Robinson, 2008) Negative (Dekker, Peen, Goris, Heijnen, & Kwakman, 1997)
Age	Positive (Huntley et al., 1998; Oiesvold et al., 1999; Pertile et al., 2011), Negative (Chung et al., 2010) Non-linear (Harman et al., 2004; McCrone & Lorusso, 1999)
Patient-level characteristics	
Gender	Positive for males (Chung et al., 2010; Rothbard & Schinnar, 1996) Positive for females (Oiesvold et al., 1999; Pertile et al., 2011; Tulloch et al., 2011)
Ethnicity	Positive for black ethnicity (Evison, 2009) (Padgett, Patrick, Burns, & Schlesinger, 1994)
Provider-level characteristics	
Hospital capacity	Positive (Chung et al., 2010; Imai et al., 2005; Tulloch et al., 2011)
Human resources of healthcare professionals	Negative (Chung et al., 2010; Imai et al., 2005)
Readmission rates	Positive (Korkeila, Lehtinen, Tuori, & Helenius, 1998) Negative (Appleby, Desai, Luchins, Gibbons, & Hedeker, 1993) (Boden, Brandt, Kieler, Andersen, & Reutfors, 2011) (Figueroa, Harman, & Engberg, 2004; Information, 2008) (Lin et al., 2006; Sytema & Burgess, 1999)

References:

- Abas, M. A., Vanderpyl, J., & Robinson, E. (2008). Socioeconomic deprivation and extended hospitalization in severe mental disorder: a two-year follow-up study. *Psychiatr Serv*, 59(3), 322-325. doi: 10.1176/appi.ps.59.3.322
- Appleby, L., Desai, P. N., Luchins, D. J., Gibbons, R. D., & Hedeker, D. R. (1993). Length of stay and recidivism in schizophrenia: a study of public psychiatric hospital patients. *Am J Psychiatry*, 150(1), 72-76.
- Boden, R., Brandt, L., Kieler, H., Andersen, M., & Reutfors, J. (2011). Early non-adherence to medication and other risk factors for rehospitalization in schizophrenia and schizoaffective disorder. *Schizophr Res*, 133(1-3), 36-41. doi: 10.1016/j.schres.2011.08.024
- Chung, W., Oh, S. M., Suh, T., Lee, Y. M., Oh, B. H., & Yoon, C. W. (2010). Determinants of length of stay for psychiatric inpatients: analysis of a national database covering the entire Korean elderly population. *Health Policy*, 94(2), 120-128. doi: 10.1016/j.healthpol.2009.09.004
- Compton, M. T., Craw, J., & Rudisch, B. E. (2006). Determinants of inpatient psychiatric length of stay in an urban county hospital. *Psychiatr Q*, 77(2), 173-188. doi: 10.1007/s11126-006-9005-z
- Dausey, D. J., Rosenheck, R. A., & Lehman, A. F. (2002). Preadmission care as a new mental health performance indicator. *Psychiatr Serv*, 53(11), 1451-1455.
- Dekker, J., Peen, J., Goris, A., Heijnen, H., & Kwakman, H. (1997). Social deprivation and psychiatric admission rates in Amsterdam. *Soc Psychiatry Psychiatr Epidemiol*, 32(8), 485-492.
- Douzenis, A., Seretis, D., Nika, S., Nikolaidou, P., Papadopoulou, A., Rizos, E. N., . . . Lykouras, L. (2012). Factors affecting hospital stay in psychiatric patients: the role of active comorbidity. *BMC Health Serv Res*, 12, 166. doi: 10.1186/1472-6963-12-166
- Evison, G. G. a. F. (2009). Use of new mental health services by ethnic minorities in England. England: North East Public Health Observatory.
- Figueroa, R., Harman, J., & Engberg, J. (2004). Use of claims data to examine the impact of length of inpatient psychiatric stay on readmission rate. *Psychiatr Serv*, 55(5), 560-565.
- Harman, J. S., Cuffel, B. J., & Kelleher, K. J. (2004). Profiling hospitals for length of stay for treatment of psychiatric disorders. *J Behav Health Serv Res*, 31(1), 66-74.
- Huntley, D. A., Cho, D. W., Christman, J., & Csernansky, J. G. (1998). Predicting length of stay in an acute psychiatric hospital. *Psychiatr Serv*, 49(8), 1049-1053.
- Imai, H., Hosomi, J., Nakao, H., Tsukino, H., Katoh, T., Itoh, T., & Yoshida, T. (2005). Characteristics of psychiatric hospitals associated with length of stay in Japan. *Health Policy*, 74(2), 115-121. doi: 10.1016/j.healthpol.2004.12.018
- Information, C. I. f. H. (2008). Hospital Length of Stay and Readmission for Individuals Diagnosed With Schizophrenia: Are They Related?. Ottawa: Canadian Institute for Health Information
- Kallert, T. W., Glockner, M., & Schutzwohl, M. (2008). Involuntary vs. voluntary hospital admission. A systematic literature review on outcome diversity. *Eur Arch Psychiatry Clin Neurosci*, 258(4), 195-209. doi: 10.1007/s00406-007-0777-4
- Korkeila, J. A., Lehtinen, V., Tuori, T., & Helenius, H. (1998). Frequently hospitalised psychiatric patients: a study of predictive factors. *Soc Psychiatry Psychiatr Epidemiol*, 33(11), 528-534.
- Lay, B., Lauber, C., & Rossler, W. (2006). Prediction of in-patient use in first-admitted patients with psychosis. *Eur Psychiatry*, 21(6), 401-409. doi: 10.1016/j.eurpsy.2005.12.004
- Lin, H. C., Tian, W. H., Chen, C. S., Liu, T. C., Tsai, S. Y., & Lee, H. C. (2006). The association between readmission rates and length of stay for schizophrenia: a 3-year population-based study. *Schizophr Res*, 83(2-3), 211-214. doi: 10.1016/j.schres.2006.01.012
- McCrone, P., & Lorusso, P. (1999). A comparison of psychiatric casemix in the UK and Italy. *Soc Psychiatry Psychiatr Epidemiol*, 34(8), 432-436.
- Oiesvold, T., Saarento, O., Sytema, S., Christiansen, L., Gostas, G., Lonnerberg, O., . . . Hansson, L. (1999). The Nordic Comparative Study on Sectorized Psychiatry--length of in-patient stay. *Acta Psychiatr Scand*, 100(3), 220-228.

- Padgett, D. K., Patrick, C., Burns, B. J., & Schlesinger, H. J. (1994). Ethnic differences in use of inpatient mental health services by blacks, whites, and Hispanics in a national insured population. *Health Serv Res*, 29(2), 135-153.
- Pertile, R., Donisi, V., Grigoletti, L., Angelozzi, A., Zamengo, G., Zulian, G., & Amaddeo, F. (2011). DRGs and other patient-, service- and area-level factors influencing length of stay in acute psychiatric wards: the Veneto Region experience. *Soc Psychiatry Psychiatr Epidemiol*, 46(7), 651-660. doi: 10.1007/s00127-010-0231-1
- Rothbard, A. B., & Schinnar, A. P. (1996). Community determinants of psychiatric hospitalization and length of stay. *Socio-Economic Planning Sciences*, 30(1), 27-38. doi: [http://dx.doi.org/10.1016/0038-0121\(95\)00023-2](http://dx.doi.org/10.1016/0038-0121(95)00023-2)
- Sytema, S., & Burgess, P. (1999). Continuity of care and readmission in two service systems: a comparative Victorian and Groningen case-register study. *Acta Psychiatr Scand*, 100(3), 212-219.
- Tulloch, A. D., Fearon, P., & David, A. S. (2011). Length of stay of general psychiatric inpatients in the United States: systematic review. *Adm Policy Ment Health*, 38(3), 155-168. doi: 10.1007/s10488-010-0310-3