



ETIOLOGY

BRIEF REPORT

The 12-item WHO Disability Assessment Schedule II as an outcome measure for treatment of common mental disorders

C.C. Thomas^{1,2,*}, S.D. Rathod², M.J. De Silva², H.A. Weiss³ and V. Patel^{2,4}

¹Department of Psychology, Princeton University, Princeton, NJ, USA

²Department of Population Health, London School of Hygiene and Tropical Medicine, London, UK

³MRC Tropical Epidemiology Group, London School of Hygiene and Tropical Medicine, London, UK

⁴Sangath, Alto-Porvorim, Goa, India

Global Mental Health (2016), 3, e14, page 1 of 7. doi:10.1017/gmh.2016.7

Background. Common mental disorders (CMD) are among the most significant contributors to disability worldwide. Patient-reported disability outcomes should be included as a key metric in the comparative assessment of value across global mental health interventions. This study aims to evaluate the validity of a widely used, cross-cultural tool – the 12-item World Health Organization Disability Assessment Schedule II (WHODAS) – as a functional outcome measure for CMD treatment.

Methods. The study population includes 1024 participants with CMD enrolled in the MANAS trial in India. CMD was assessed using the Revised Clinical Interview Schedule (CIS-R). Disability was assessed using the 12-item WHODAS II plus a measure of disability days. This analysis presents the correlations between these disability items and CMD symptom severity at 2 months after enrollment (convergent validity) and the items' associations with CMD recovery 4 months later (external responsiveness).

Results. All items showed a positive correlation of disability with CMD symptom severity ($p < 0.001$). The WHODAS items of 'standing,' 'household responsibilities,' and 'emotional disturbance' explained the most variance in CMD symptom severity. Improvements in 'disability days,' 'emotional disturbance,' 'standing,' 'household responsibilities,' 'day-to-day work,' and 'concentrating' were significantly associated with CMD recovery over follow-up.

Conclusions. Further research is recommended on a CMD-specific WHODAS subscale comprised of the six WHODAS items found to be most strongly associated with CMD severity and recovery. This shorter, CMD-specific disability subscale would critically serve as a common metric to compare intervention impact on patient-centered outcomes and, in turn, to allocate global mental health resources efficiently.

Received 26 November 2015; Accepted 3 February 2016

Key words: 12-item WHODAS, anxiety, depression, disability, etiology, validity.

Background

Common mental disorders (CMD), comprising depressive and anxiety disorders, cause significant disability worldwide, with depressive disorders alone estimated to be the second leading global contributor (Ferrari *et al.* 2013). CMD are associated with such

* Address for correspondence C. Thomas, 426 Peretsman-Scully Hall, Department of Psychology, Princeton University, Princeton, NJ, 08540, USA.

(Email: ct13@princeton.edu)

© The Author(s) 2016. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



disabling consequences as diminished economic productivity, loss of employment, and impaired social functioning and, overall, contribute substantial socioeconomic burdens on patients and communities (Ormel *et al.* 1994; Judd *et al.* 2000; Patel *et al.* 2010; Silva *et al.* 2013). In part due to the large burden of CMD globally, particularly in low- and middle-income countries with wide treatment gaps, the regular assessment of disability outcomes of mental health interventions has been selected as one of the top 25 research priorities for the field of global mental health through a Delphi panel of hundreds of international stakeholders (Collins *et al.* 2011).

Disability is defined by the World Health Organization (WHO) as the disruption of an individual's interaction with his or her environment (WHO, 2010). A common metric of self-reported disability is a necessary tool for the evaluation and comparison of intervention impact on outcomes that matter to patients. While the diversity and reach of psychosocial treatments for common mental disorders have been rapidly growing over the past several decades, a recent systematic search found that fewer than 5% of clinical trials for the treatment of depression reported a measure of functional outcomes (McKnight & Kashdan, 2009). In the burgeoning movement that extends and tailors mental health treatments to new cultures and contexts, there is a need for a common metric for mental health program evaluation that includes patient-centered outcomes.

Due to its widespread international use, sound psychometric properties, and ease of administration, the WHO Disability Assessment Schedule II (WHODAS) is a prime candidate for the routine evaluation of intervention impact on patient-reported disability outcomes (WHO, 2010). Developed in 1998, the WHODAS is a cross-cultural tool that captures social, occupational, physical, and role impairments associated with a health condition (WHO, 2010). As a component in disease burden calculations (i.e. through Disability-Adjusted Life Years), the WHODAS enables the comparison of cost-effectiveness across diverse health interventions, which in turn guides planning of health policies and programs. While the WHODAS has heretofore been used primarily to assess disability associated with physical illnesses, the WHODAS is being increasingly applied to study disability among psychiatric populations in low- and middle-income countries. For example, it has been applied to assess functional impairments associated with mental disorders, understand relationships between mental illness and physical comorbidities, and measure mental health treatment outcomes (Akinsulore *et al.* 2015, De Silva *et al.* 2015, Faye *et al.* 2015).

This study aims to take the first step in evaluating the WHODAS as an outcome measure for CMD

treatment by assessing the convergent validity of WHODAS disability items with CMD severity and the external responsiveness of items to recovery from CMD. The individual WHODAS items found to be more correlated to CMD severity and responsive to CMD recovery are furthermore recommended as candidates for future development of a CMD-specific WHODAS subscale. This subscale would strengthen assessment of functional outcomes in studies of CMD recovery.

Methods

Study sample

The study sample for this secondary analysis includes patients enrolled in a cluster randomized controlled trial of CMD treatments in Goa, India (the MANAS trial) (Chatterjee *et al.*, 2008; Patel *et al.* 2010, 2011). This sample includes patients diagnosed with CMD, according to the International Statistical Classification of Diseases and Related Health Problems – 10th revision (ICD-10), being treated in the collaborative stepped-care ($n=423$) and enhanced usual care ($n=601$) arms. Conducted between 2007 and 2009, the MANAS trial included 2796 adult primary care attenders who screened positive for CMD according to the 12-item General Health Questionnaire (GHQ) in 12 public health centers and 12 private general practitioner clinics. Details of the study design, the baseline characteristics of participants, and the treatments effects on CMD and disability outcomes have been published elsewhere (Chatterjee *et al.* 2008; Patel *et al.* 2008, 2010, 2011). The MANAS trial is registered with ClinicalTrials.gov with identifier NCT00445407.

Study design

Disability measurements were taken at the 2-, 6-, and 12-month time points after enrollment in the MANAS trial. Of the 2491 participants attending the 2-month follow-up, 1024 (41.1%) maintained a diagnosis of CMD. This subsample of 1024 participants was included in the current analysis as a cohort followed from 2 to 6 months after enrollment.

Instruments

Clinical outcomes were measured with the Revised Clinical Interview Schedule (CIS-R), a standardized, structured CMD diagnostic tool that has been field-tested for use in Goa (Lewis *et al.* 1992; Patel *et al.* 1998, 2011). Lay health workers used the CIS-R to generate an ICD-10 diagnosis and a symptom severity score (0–57) at each follow-up point (Patel *et al.* 2010). Recovery from CMD at the 6-month follow-up visit



was determined with the CIS-R according to ICD-10 criteria (Patel *et al.* 2010).

Disability outcomes were assessed using the 12-item WHODAS II, and two items from the 36-item WHODAS II. The WHODAS II measures items in six domains of functioning as experienced over the past 30 days: mobility, self-care, life activities, understanding and communicating (U&C), interpersonal interactions, and participation in society (WHO, 2010). The 12-item tool assesses each domain with two items that are measured on a 3-point scale in which 1 indicates no disability, 2 indicates mild to moderate disability, and 3 indicates severe to extreme disability. These items were summed to generate a total score between 12 (no disability) and 36 (maximum disability). A total number of disability days were computed from two items of the 36-item WHODAS assessing days of no work or of reduced work due to illness in the past 30 days and was assessed by lay health workers if patients reported at least mild disability on any of the 12 WHODAS items (Patel *et al.* 2008).

Statistical analyses

The convergent validity of each WHODAS item with the CIS-R symptom severity score was evaluated using a non-parametric equivalent of the Pearson’s correlation coefficient, Spearman’s rank-order correlation coefficient r_s , which indicates the extent of correlation between the disability item score and the CIS-R score. Similar to squaring Pearson’s correlation coefficient, Spearman’s rank-order correlation coefficient may be squared (r_s^2) to estimate the variance in CIS-R scores explained by the disability item score as an effect size statistic (Rosenthal, 1994; Pett, 1997). Qualitative evaluations were used to assess the degree of correlation, with Pearson’s correlation values of approximately 0 (no correlation to weak correlation), ± 0.50 (moderate correlation), ± 0.71 (high correlation), and ± 1 (perfect correlation) corresponding to Spearman’s rank correlations of 0, ± 0.48 , ± 0.69 and ± 1 (Kirkwood & Sterne, 2003).

To identify which specific aspects of disability were most responsive to recovery, we used logistic regression to estimate magnitude of the relationship between change in individual WHODAS items and the CMD recovery outcome (i.e. external responsiveness). Recovery from CMD between the 2- and 6-month follow-up points was analyzed as the dependent variable. The change scores for each of the 12 WHODAS items, the global WHODAS score, and the total number of disability days were coded into binary measures of 0 = ‘not improved’ and 1 = ‘improved’. Each WHODAS measure was analyzed as an independent variable in separate logistic regression models, controlled for baseline CIS-R score, sex, age, clinic type,

Table 1. Distribution of study sample characteristics at 2 months after recruitment into the MANAS trial, India, 2007–2009

Sociodemographic characteristics		Mean (S.D.)/ number (%)
Age ($n = 1024$)		48.0 (13.6)
Sex ($n = 1024$)	Female	878 (85.8)
	Male	146 (14.2)
Education level ($n = 1024$)	<1 year	512 (50.0)
	1–5 years	230 (22.5)
	≥ 6 years	282 (27.5)
Employment ($n = 1023$)	Unemployed	731 (71.5)
	Full-time	126 (12.3)
	Part-time/seasonal	150 (14.7)
	Other	16 (1.6)
Marital status ($n = 1024$)	Married	632 (61.7)
	Widowed or separated/divorced	339 (33.1)
	Never married	53 (5.2)
Ethnicity ($n = 1024$)	Goan	974 (95.1)
	Other	50 (4.9)
	Religion ($n = 1024$)	Hindu
Financial status ($n = 1024$)	Christian	288 (28.1)
	Muslim	16 (1.6)
	Living comfortably	64 (6.3)
Clinical characteristics	Just getting by	361 (35.3)
	Difficult to make ends meet	599 (58.5)
Clinical characteristics		Median (IQR; range)
CIS-R score ($n = 1024$)		18 (14–24; 0–57)
Twelve-item WHODAS II global score ($n = 1024$)		19 (16–22; 12–36)
Disability days score ($n = 965$)		15 (0–30; 0–30)

and intervention arm as potential confounders. Cluster robust standard errors (S.E.) were applied to adjust the 95% confidence intervals (CI) for clinic-level clustering.

Results

Sample characteristics

The sociodemographic and clinical characteristics of the sample at the 2-month follow-up are presented in Table 1. They had a mean age of 48.0 years (S.D. = 13.6) and were predominantly female (85.8%), married (61.7%), and Hindu (70.2%). Half had no formal




Fig. 1. Convergent validity of CMD symptom severity scores with WHODAS disability measures at 2 months among participants of the MANAS trial, India, 2007–2009.

education (50.0%), and a majority was unemployed (71.5%). The median CIS-R score was 18 out of 57 (IQR: 14–24), the median WHODAS score 19 out of 36 (IQR: 16–22), and the median disability days score 15 out of 30 (IQR: 0–30). Of the initial 1024 participants, 943 (92.1%) were seen at 6 months, and of these, 368 participants (39.0%) had recovered from CMD. At both 2 and 6 months, respectively, only four patients (0.4%) had missing values for the disability days score.

Convergent Validity

Convergent validity statistics are presented in Fig. 1. All of the disability items had weak but significant correlations with CIS-R scores, with r_s values between 0.14 and 0.39. ‘Household responsibilities,’ ‘emotional disturbance,’ and ‘standing’ explained the most variance in CIS-R scores (r_s^2 values of 0.15, 0.10, and 0.09, respectively). The WHODAS global score showed the greatest extent of correlation with the CIS-R scores ($r_s=0.42$), yet remained only moderate, explaining 18% of the variance in CIS-R scores ($r_s^2 = 0.18$).

External Responsiveness

As presented in Table 2, improvement in the WHODAS global score, disability days score, and five of the 12 WHODAS item scores were associated with CMD

recovery. Participants whose global WHODAS scores improved between 2 and 6 months follow-up had on average 2.40 times higher odds of having recovered from CMD [adjusted Odds Ratio (aOR) 95% CI: 1.62–3.56]. Those who reported fewer disability days at 6 months had 2.37 higher odds of recovery (aOR 95% CI: 1.66–3.37). Of the 12 individual WHODAS items, five displayed evidence of significant responsiveness to CMD recovery: ‘emotional disturbance,’ ‘standing,’ ‘household responsibilities,’ ‘day-to-day work,’ and ‘concentrating.’

Conclusions

This study constitutes, to our knowledge, the first assessment of the validity of the 12-item WHODAS II among middle-aged patients with CMD in a low- or middle-income country. Previous studies have analyzed associations between measures of disability and CMD at a single time point or changes in disability over time as estimates of internal responsiveness without reference to CMD caseness (Judd *et al.* 2000; Garin *et al.* 2010). As a longitudinal treatment study, this study importantly adds quantifications of the associations between concurrent changes in disability and CMD caseness to the evidence base.

The present study found that five items in the domains of participation (‘emotional disturbance’), physical mobility (‘standing’), life activities



Table 2. External responsiveness of improvements in disability items with CMD recovery at 6 months among participants of the MANAS trial, India, 2007–2009

	Domain	Item	Adjusted OR (95% CI)
12-item WHODAS II (<i>n</i> = 943)	Mobility	Standing	1.67 (1.19–2.34)***
		Walking a kilometer	1.23 (0.87–1.72)
	Self-care	Washing whole body	1.10 (0.69–1.75)
		Getting dressed	0.83 (0.51–1.34)
		Concentrating	1.30 (1.01–1.67)**
	Understanding and communicating	Learning a new task	0.99 (0.70–1.40)
		Maintaining a friendship	0.76 (0.36–1.58)
	Inter-personal	Dealing with strangers	0.94 (0.52–1.69)
		Household responsibilities	1.67 (1.19–2.34)***
	Life activities	Day-to-day work	1.40 (1.01–1.94)**
		Community participation	1.03 (0.70–1.50)
	Participation	Emotional disturbance	1.81 (1.27–2.58)***
Global score		2.40 (1.62–3.56)***	
Disability days in past month (<i>n</i> = 836)			2.37 (1.66–3.37)***

Odds ratios adjusted for baseline CIS-R score, sex, age, clinic type, and intervention arm, with cluster robust standard errors (S.E.).

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$, p value from Wald test in fully adjusted model.

(‘household responsibilities’ and ‘day-to-day work’), and understanding and communicating (‘concentrating’) most strongly correlated with symptom severity in the cross-section and responded to CMD recovery. In contrast, items in the interpersonal and self-care domains showed little to no relationship with severity or recovery. Despite weak convergent validity, the disability days score showed strong responsiveness to recovery. Although correlation coefficients for all WHODAS items with CIS-R scores were weak, five WHODAS items and the disability days item showed the greatest extent of correlation with CMD severity and significant associations with CMD recovery; thus, these six WHODAS items are recommended for further research in the development of a CMD-specific subscale.

This pattern of results largely aligns with studies of patients with depression in Europe and the USA in which the WHODAS domains of participation and life activities have been found to be the most internally responsive (i.e. change over time) (Chwastiak & Von Korff, 2003; Garin *et al.* 2010). In contrast to Chwastiak & Von Korff (2003), this study found the self-care domain to be responsive and the interpersonal domain to be largely unresponsive. However, in this study few patients with CMD experienced disability in this domain at 2 months, and it is possible that gains in interpersonal functioning occurred largely within the first 2 months of the trial.

Unlike much of the research on patients with depression in high-income countries, physical mobility items showed strong responsiveness. A qualitative

study of the MANAS trial found weakness/tiredness to be one of the two most commonly reported reasons for seeking care, suggesting a somatic manifestation of CMD in this context (Andrew *et al.* 2012). Thus, the physical mobility domain may be heterogeneously associated with CMD cross-culturally, and the relationship may differ for those with anxiety versus those with depressive disorders.

The observed external responsiveness of the disability days measure to CMD recovery is consistent with findings from a review of depression treatment studies in high-income settings. Notably, Mintz *et al.* (1992) found that work role impairments, including disability days, were more consistently alleviated when patients’ symptoms were also alleviated and, moreover, that most of the variance in work outcomes over the course of treatment was attributable to symptom remission.

This study has several limitations. First, sample selection limits the generalizability. As disability was not measured at trial baseline, this study includes patients who presented with CMD at 2 months and thus were experiencing more chronic episodes of CMD. Moreover, at this follow-up point, 305 patients (10.9%) enrolled in the trial at baseline were missing at 2 months and these missing patients were more likely to be male, younger, and have less severe CMD. The WHODAS assessment did not take place during trial enrollment, during patient primary care visits, in order to minimize the burden on participants. The WHODAS assessments took place at 2 and 6 months, with all other follow-up assessments, during



planned visits to participants' homes. Second, the 'U-shaped' distribution of the disability days scores, being clustered largely around 0 and 30 days with a small peak at 15 days, suggests recall bias. Because this bias has unpredictable implications for the results, the disability days statistics should be interpreted with caution. Lastly, during piloting the research team observed that participants found the fine judgments required of the 5-point WHODAS scale difficult and thus collapsed the WHODAS scale to three points in the trial. This 3-point response scale did not allow for a fine-grained assessment of responsiveness. In addition, an implication of this collapse of the scale for individual WHODAS items is non-differential misclassification of the WHODAS scores, and thus bias toward the null when assessing correlation and regression coefficients for WHODAS items.

This study provides a foundational step in the development of a common disability metric for evaluating mental health treatment outcomes. The findings suggest that a WHODAS subscale of the six items found to be most strongly associated with CMD severity and recovery could provide a more valid and meaningful measure of CMD recovery in evaluations of CMD interventions. Future research is recommended to confirm the scale reduction of the WHODAS, as evaluated here, by measuring the WHODAS items on a 5-point response scale and using item-response theory methods, such as those used in Bokken (1971). Prior to its use across mental health evaluation studies, the resulting subscale would need to be validated among primary care-seeking populations with CMD in several low- and middle-income countries. The routine use of this CMD-specific subscale of the WHODAS would serve as a critical component of an evidence-based, cost-efficient, and patient-centered response to the challenge of global mental health.

Acknowledgements

This article is based on data from the MANAS study that occurred in Goa, India from 2007 to 2009. For the use of this data, we thank the project team comprised of individuals from the London School of Hygiene & Tropical Medicine, Sangath, the Directorate of Health Services (Government of Goa) and the Voluntary Health Association of Goa. The MANAS study was funded by a Wellcome Trust Senior Researcher Fellowship in Clinical Science to VP (grant number 091834/Z/10/Z). MDS was funded by a Wellcome Trust/LSHTM fellowship.

Declaration of Interest

None.

Ethical Standards

Ethics approval was granted by the London School of Hygiene & Tropical Medicine (UK) and the Institutional Review Board of Sangath (India), the implementing organization for both the original trial and this study. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All persons gave their informed consent prior to inclusion in the study.

References

- Akinsulore A, Mapayi BM, Aloba OO, Oloniniyi I, Fatoye FO, Makanjuola RO (2015). Disability assessment as an outcome measure: a comparative study of Nigerian outpatients with schizophrenia and healthy control. *Annals of General Psychiatry* 14, 40.
- Andrew G, Cohen A, Salgaonkar S, Patel V (2012). The explanatory models of depression and anxiety in primary care: a qualitative study from India. *BMC Research Notes* 5, 499.
- Bokken RJ (1971). *A Theory and Procedure of Scale Analysis: with Applications in Political Research*. Mouton & Co: the Netherlands.
- Chatterjee S, Chowdhary N, Pednekar S, Cohen A, Andrew G, Araya R, Simon G, King M, Telles S, Weiss H, Verdelli H, Clougherty K, Kirkwood B, Patel V (2008). Integrating evidence-based treatments for common mental disorders in routine primary care: feasibility and acceptability of the MANAS intervention in Goa, India. *World Psychiatry* 7, 39–46.
- Chwastiak LA, Von Korff M (2003). Disability in depression and back pain. *Journal of Clinical Epidemiology* 56, 507–514.
- Collins PY, Patel V, Joestl SS, March D, Insel TR, Daar AS, Bordin IA, Costello EJ, Durkin M, Fairburn C, Glass RI (2011). Grand challenges in global mental health. *Nature* 475, 27–30.
- De Silva MJ, Rathod SD, Hanlon C, Breuer E, Chisholm D, Fekadu A, Jordans M, Kigozi F, Petersen I, Shidhaye R, Medhin G, Ssebunnya J, Prince M, Thornicroft G, Tomlinson M, Lund C, Patel V (2015). Evaluation of district mental healthcare plans: the PRIME consortium methodology. *The British Journal of Psychiatry* (<http://doi:10.1192/bjp.bp.114.153858>).
- Faye AD, Gawande S, Tadke R, Kirpekar VC, Bhawe SH, Pakhare AP, Tayade B (2015). Do panic symptoms affect the quality of life and add to the disability in patients with bronchial asthma? *Psychiatry Journal* 15, Article ID 608351, 8 pages.
- Ferrari AJ, Charlson FJ, Norman RE, Patten SB, Freedman G, Murray CJ, Vos T, Whiteford HA (2013). Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Medicine* 10, e1001547.
- Garin O, Ayuso-Mateos J, Almansa J, Nieto M, Chatterji S, Vilagut G, Alonso J, Cieza A, Svetskova O, Burger H,



- Racca V, Francescutti C, Vieta E, Kostanjsek N, Raggi A, Leonardi M, Ferrer M** (2010). Validation of the 'World Health Organization disability assessment schedule, WHODAS-2' in patients with chronic diseases. *Health and Quality of Life Outcomes* **8**, 51.
- Judd LL, Akiskal HS, Zeller PJ, Paulus M, Leon AC, Maser JD, Endicott J, Coryell W, Kunovac JL, Mueller TI, Rice JP, Keller MB** (2000). Psychosocial disability during the long-term course of unipolar major depressive disorder. *Archives of General Psychiatry* **57**, 375–380.
- Kirkwood B, Sterne J** (2003). *Essential Medical Statistics*. Blackwell Science Ltd: Oxford.
- Lewis G, Pelosi AJ, Araya R, Dunn G** (1992). Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychological Medicine* **22**, 465–486.
- McKnight PE, Kashdan TB** (2009). The importance of functional impairment to mental health outcomes: A case for reassessing our goals in depression treatment research. *Clinical Psychology Review* **29**, 243–259.
- Mintz J, Mintz LI, Arruda MJ, Hwang SS** (1992). Treatments of depression and the functional capacity to work. *Archives of General Psychiatry* **49**, 761.
- Ormel J, VonKorff M, Ustun T, Pini S, Korten A, Oldehinkel T** (1994). Common mental disorders and disability across cultures: results from the who collaborative study on psychological problems in general health care. *JAMA* **272**, 1741–1748.
- Patel V, Pereira J, Coutinho L, Fernandes R, Fernandes J, Mann A** (1998). Poverty, psychological disorder and disability in primary care attenders in Goa, India. *The British Journal of Psychiatry* **172**, 533–536.
- Patel VH, Kirkwood BR, Pednekar S, Araya R, King M, Chisholm D, Simon G, Weiss H** (2008). Improving the outcomes of primary care attenders with common mental disorders in developing countries: a cluster randomized controlled trial of a collaborative stepped care intervention in Goa, India. *Trials* **9**, 4.
- Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, De Silva MJ, Bhat B, Araya R, King M, Simon G, Verdelli H, Kirkwood B** (2010). Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *The Lancet* **376**, 2086–2095.
- Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, Bhat B, Araya R, King M, Simon G, Verdelli H, Kirkwood B** (2011). Lay health worker led intervention for depressive and anxiety disorders in India: impact on clinical and disability outcomes over 12 months. *The British Journal of Psychiatry* **199**, 459–466.
- Pett MA** (1997). *Nonparametric statistics for health care research: statistics for small samples and unusual distributions*. Sage Publications: Thousand Oaks, California.
- Rosenthal R** (1994). Parametric measures of effect size. In *The Handbook of Research Synthesis* (ed. H Cooper, LV Hedges), pp. 231–244. Sage: New York.
- Silva MJD, Cooper S, Li HL, Lund C, Patel V** (2013). Effect of psychosocial interventions on social functioning in depression and schizophrenia: meta-analysis. *The British Journal of Psychiatry* **202**, 253–260.
- World Health Organization** (2010). *Measuring Health and Disability: Manual for WHO Disability Assessment Schedule WHODAS 2.0*. WHO Press: Geneva.