

ANALYSIS

Europe's men need their own health strategy

A recent European report on men's health shows that it lags behind that of women. **Alan White and colleagues** analyse the problems and call for more policy, practice, and research aimed specifically at men

Alan White *professor of men's health*¹, Martin McKee *professor of European public health*², Noel Richardson *director*³, Richard de Visser *lecturer*⁴, Svend Aage Madsen *head of department of psychology, play therapy, and social counselling*⁵, Bruno C de Sousa *research fellow*⁶, Richard Hogston *director*¹, Witold Zatoński *director*⁷, Péter Makara *professor*⁸

¹Centre for Men's Health, Institute for Health and Wellbeing, Faculty of Health and Social Sciences, Leeds Metropolitan University, Leeds LS2 8NU, UK; ²London School of Hygiene and Tropical Medicine, London, UK; ³Centre for Men's Health, Institute of Technology Carlow, Carlow, Ireland; ⁴School of Psychology, University of Sussex, Falmer, UK; ⁵Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark; ⁶Centro de Malária e Doenças Tropicais, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisbon, Portugal; ⁷Department of Cancer Epidemiology and Prevention, Cancer Centre and Institute of Oncology, Warsaw, Poland; ⁸Institute of Public Health, Semmelweis University, Budapest, Hungary

Ten years ago the *BMJ* published a special issue on men's health.¹ It noted how, although men fare better than women in most conventional measures such as top jobs and earnings, this advantage is not reflected in their health. A report we produced this summer, *The State of Men's Health in Europe*,^{2,3} shows that little has changed. At any given age, men are still more likely than women to die from most of the leading causes, and in the European Union men have more than twice as many deaths a year as women throughout the working ages (15-64 years). This high level of premature mortality in men has psychological, social, and economic consequences for relatives, households, communities, and the workplace. Yet, in both national and European health policy, men and "masculinity" are largely taken for granted. This has limited the development of evidence based programmes that meet their health needs.

Differences in mortality and morbidity are not simply the result of biological factors; nor are they intractable. In fact, the health gap between men and women varies considerably. It is much greater in eastern Europe than in western Europe,⁴ and within countries it is influenced by class, education, employment, and other social determinants.⁵ The clustering of material, cultural, and psychosocial factors seems to be particularly detrimental to the health of many men.⁶ These factors contribute to gendered lifestyles and behaviours that have traditionally been seen as predominantly "masculine"⁷ and that cause many of the premature deaths in men. Traditional masculine attitudes are associated with unhealthy behaviours such as poor diet,⁸ smoking, excessive alcohol consumption,⁹ non-use or delayed uptake of health services,¹⁰ and higher likelihood of injury. All of these factors are more common among men living in eastern

Europe than those in western Europe and in poorer material and social conditions everywhere.¹¹ Men also seem to have adapted less well than women to the changes that have accompanied the political and social upheavals in eastern Europe in recent decades, such as more transient and unstable working conditions, increasing unemployment, and changing family structures (reduction in marriage and increased divorce).¹²

Yet, paradoxically, men often view themselves as having better health than women. There is some justification for this view: those men who survive into old age report less disability than women of the same age²; but what is overlooked is that fewer live this long.¹³ Though the average difference in life expectancy between men and women in the European Union is 6.1 years, it ranges from 11.3 years in Latvia to 3.3 years in Iceland and Lichtenstein.² Thus, men in general, and younger men in particular, tend to minimise the potential consequences of practices damaging to health and to see themselves as invulnerable to danger and risk.

The situation is at last beginning to change. *The State of Men's Health in Europe* provides a springboard for a range of policy measures, innovations in practice, and research into men's health.² The report, together with emerging men's health policy and initiatives in EU member states and the work of key non-governmental organisations (including the European Men's Health Forum, the Men's Health Forum, and the Nordic Men's Health Network) have begun to enhance our understanding of what can be done to improve men's health. Here we review the report's key conclusions.

Policy

Given the many influences on men's health, policy responses must take a broad perspective, acknowledging the variety of social positions, lifestyles, and health needs of men and women, rather than focusing only on differences between them. Many measures designed to improve opportunities for men in employment, education, and other sectors will also help women. However, men may benefit disproportionately when they are at greatest relative risk—as is the case for whole population initiatives in areas such as preventing road traffic injuries (where death rates are three times higher for men) and workplace health and safety (where men account for 95% of fatal incidents).

Because policies may impinge differently on women and men, it is important to consider the effects on each separately. To date, few EU member states (Ireland, UK, and Denmark being the exceptions) have identified men as a specific population group for the strategic planning of health.¹⁴⁻¹⁶ Ireland's approach to developing and implementing policy on men's health can be a particular source of inspiration for other countries (box 1).¹⁷

The large variations in trends in male mortality across Europe² highlight the potential benefits of a broader implementation of measures that have reduced male deaths. An obvious area is road traffic deaths, which have already fallen substantially in countries that have joined the EU since 2004, reflecting factors such as more effective enforcement of traffic laws and investment in safer roads. Although the percentage fall in deaths was similar in men and women between 2000 and 2009 (20.3% and 17.3%, respectively), the absolute reduction has been much greater for men (2640 fewer male deaths compared with 602 fewer female deaths).¹⁸

Similarly, the opening of markets in eastern Europe, and the accompanying improvements in diet and subsequently cardiovascular deaths have benefited men disproportionately.¹⁹ Nevertheless, cardiovascular disease still accounts for more deaths of men in eastern Europe than in western Europe (for example, 62% of male deaths in Bulgaria compared with 25% in France²). This points to the need for the broader implementation of established primary and secondary prevention measures that account for the sex/gendered nature of the disease—men develop the disease earlier than women and the risk factors are more prevalent in men.

Action is needed on several levels. The first is schools. Behaviours and values developed early in life have a critical influence on men's later health practices. There is a need for a visible, integrated focus on boys' and men's health within primary and secondary school curriculums that can foster positive models of physical, psychological, and social development.²⁰ This should challenge gender stereotypes that are harmful to men's health and develop boys' ability to maintain their emotional wellbeing, problem solving skills, and responsible decision making. This will lay the foundations for empowering men to manage their health more effectively, to make more informed decisions about their relative risk, and to use health services responsibly. An increased focus on schools must also tackle the problem of boys' lower educational performance.

The second level is the workplace. Employers and unions can work collaboratively to support policies and programmes to promote men's health in the workplace—not only through traditional occupational health measures but also by helping them to deal with lack of job security, prolonged sickness absence, and involuntary early retirement, all of which negatively affect men's health.²¹⁻²³ This requires governments and employers to consider the health of the workforce as an

investment and to recognise the increasingly strong evidence that work benefits physical and mental health and wellbeing. Unemployment not only impairs individuals' health but also leads to a loss of worth and self confidence, poorer social integration, and poverty. Work-life balance must also be considered. Measures that support and enable men to be more involved fathers (for example, paid paternal leave) benefit not only fathers but also their partners and children, and society as a whole.^{24 25} Although workplace interventions can help improve employees' health, men may be less likely than women to engage with them.²⁶

The third area for action is in policies that target marginalised subgroups of men, such as ethnic minorities, sexual minorities, disabled men, and those who are isolated in rural areas, homeless, or in prison, all of whom experience considerably higher morbidity and premature mortality.

Practice

Barriers to accessing health services still exist in many European countries, and these may exacerbate the traditionally lower use of appropriate care by men. It is thus important to identify and develop alternative services that engage men more effectively. However, these should not entrench or reinforce stereotypes that all men are unhealthy or uninterested in their health. An effective response must move beyond increasing access to family practitioners: it should extend to weight loss groups, counselling services, and health promotion activities. Paradoxically, although male socialisation patterns tend to result in men being less engaged with health and wellbeing issues than women, men are seldom the focus of targeted health education or health promotion.

Mental health is a case in point. Many men present their mental and emotional distress differently from women, so that the diagnosis is often missed. Men are less likely to contact health services because of mental health problems,²⁷ potentially placing them at risk of deterioration and suicide. Men commit suicide three to five times more often than women. Men may also feel compelled to use other coping strategies such as acting aggressively, being uncooperative with health professionals, rejecting offers of help, and resorting to alcohol misuse.

Although health initiatives aimed specifically at men have increased recently, much of this work has been small scale pilots with few assessed in definitive trials.²⁸ Nevertheless, longitudinal studies show that young men with more traditional beliefs about masculinity are significantly less likely to use health services²⁹ and that when they do seek help, flexible opening times, longer consultation times, the provision of lifestyle and behaviour modification programmes, and a comprehensive referral system are important to them.³⁰

Weaknesses in existing health services have stimulated interest in initiatives that bring health services to men in settings such as pubs, sports clubs, schools, and youth centres. The workplace is also an important setting in which to target services, screening initiatives, information, and support to men. One of the larger scale examples is the £1.6m (€1.85m; \$2.5m) invested in the English Premier League health initiative aimed at 18-35 year old male football supporters. As part of this three year programme, health trainers based in 16 clubs are engaging men to act on multiple aspects of physical, psychological, and social wellbeing. Clubs use a range of activities suited to their local male populations, including educational activities for supporters on match days and outreach work into schools and communities, weekly lifestyle classes, and football or multisport sessions for men. Baseline analysis shows that the initiative is reaching men

Box 1 Key steps in the development of Ireland's national men's health policy^{16 17}

- Men were identified as a specific population group
- A three year research project into men's health was commissioned to inform national policy
- A consultation process included national men's health days, targeted focus groups, and targeted and public calls for submissions
- Policy was based on an extensive review of national and international literature
- Meetings were held between government departments to finalise the policy and action plan

Underpinning principles

- To develop men's health, not just within a health policy context but in synergy with other government departments and across different sectors
- To adopt a social determinants and gender relations approach
- To include preventive and health promotion and community development to reach subgroups of men most in need.

Process for implementation

- National Implementation Group (including annual audit) has been established
- Independent evaluation will be undertaken after five years

Examples of actions resulting from policy

- Increased focus on men's health in the workplace and in community settings targeting disadvantaged men
- Development of guidelines for training in men's health
- Establishment of a national centre for men's health research
- Providing funding to the Men's Health Forum in Ireland to coordinate "Men's Health Week."
- The policy also called on national governing sporting bodies to provide a range of activities to stave off men's retirement from competitive sport. The Football Association of Ireland now runs a veterans football programme and the Gaelic Athletic Association has developed a social inclusion initiative for older men who are not participating in community life (www.gaa.ie/clubzone/gaa-social-initiative/description-of-initiative/)

in need.³¹ A clinical trial of the effectiveness of using football clubs as a conduit for improving men's fitness and weight is currently under way with the Scottish Premier League, under the banner of Football Fans in Training. The concept of "men's sheds," developed in Australia as a place for male focused activities outside the home and work,³² is now spreading into Europe, especially Ireland and the UK. Men's sheds may provide a captive audience for health promotion activities, and they may also be an important setting for challenging men's stereotypical beliefs about masculinity and healthcare.

Research

Research also presents a paradox. Although for many decades research on common diseases, and especially cardiovascular disease, was largely done on men and applied uncritically to women, this was mainly because of a combination of gender blindness and ease of sampling from workplaces rather than having men's specific needs as part of the research question. Until the mid-1990s, there was a dearth of research that had men as its main focus.

One limitation of much epidemiological and behavioural research is that the biological category "male" is conflated with the psychological and social concept "masculine." Yet how men construct beliefs, attitudes, and behaviours can have a crucial bearing on their health. More psychological and sociological research is needed into how different ideologies of masculinity influence healthy and unhealthy behaviour.⁷ For example, efforts to tackle binge drinking among young men will need to take account of the role that alcohol plays as a rite of passage to manhood for many young men and as a mechanism for defining collective gender practice.³³ Likewise, measures designed to stem the growing tide of male obesity will need to overcome men's scepticism and cynicism towards healthy eating messages and the strong connotations between dieting and femininity.⁸

More research into masculinity will enable a more nuanced analysis of differences in perspectives and experiences of health and illness, not just between men and women but between

different subpopulations of men.³⁴ Health practices of individual men must be interpreted within the broader economic and social contexts of their lives. We need to get to grips with the "intersections"³⁵ of gender with factors such as social class, educational attainment, employment status, race, and ethnicity in order to build evidence based and gender appropriate approaches to men's health.

Conclusion

Public debate on men's health tends to be dominated by negative portrayals of men and masculinity, where men are blamed for not attending health services and taking risks with their health. This is not helpful. We should use strategies that work with men in a constructive way.

A more focused approach to the physical and mental health of younger men needs to be balanced with one that also faces the health challenges of an ageing male population. The diversity of factors contributing to men's poorer health requires measures that not only recognise any gender equality issues but also highlight a more fundamental concern with equity. This relates to the right of all men—irrespective of social, cultural, political, or ethnic differences—to live long and fulfilling lives. In policy, practice, and research there is a pressing need to examine the "problems" with men's health and to tackle the underlying causes as well as the symptoms. This demands appropriate intersectoral and intergovernmental responses at both EU and national levels.

Contributors and sources: All the authors except MMcK were authors of the *State of Men's Health* report. AW was the lead author and editor of the report, BCdeS was the statistician, and the other authors either completed sections of the report or engaged in its compilation. MMcK was the chief reviewer of the report. AW, MMcK, NR, RdeV drafted the manuscript. All authors critically revised and approved the manuscript. AW is the guarantor.

Competing interests: All authors have completed the unified competing interest form at www.icmje.org/doi_disclosure.pdf (available on request

Box 2. Examples of innovative initiatives that have benefited the health of men

*Scottish well man health service pilots*³⁵

In 2003 the Scottish government funded 18 outreach projects aimed at improving men's health. It set out to reach men in deprived areas by setting up clinics in non-traditional healthcare settings:

- Fixed and mobile nurse led clinics
- Outreach work with homeless men
- Community based initiatives targeting disadvantaged men
- Pubs
- Workplace initiatives

Activities included awareness raising activities (health fairs and marketing events), brief health assessments, comprehensive health assessments with specially trained (usually nursing) staff, and community development endeavours such as the inclusion of a health dimension in adult education and training courses for unemployed men.

Evaluation showed variable success. Sessions held in community venues or workplaces were more likely to contact men who had not seen their general practitioner in the past two years. Over 3000 men received comprehensive health checks, 72% of whom were given healthcare advice and referred to another health service.³⁵

British Telecom (BT) Workfit

This workplace joint initiative between BT and the Communications Workers' Union comprised a 16 week internet based weight loss programme

- Over 16 000 participants were registered, 75% of whom were men
- About 5000 people could be tracked through the programme, with an average weight loss of 2.3 kg
- Over 65% of the participants reported sustained changes to their lifestyle as a direct result of the programme at 4 months
- A follow-up programme to increase cancer awareness is under way³⁶

Royal Mail health initiative

Men comprise 80% of the Royal Mail's workforce. The initiative included:

- Rapid access physiotherapy
- Locating clinics at main sorting offices
- Health screening
- Health promotion, including activities aimed at smoking and back pain

An initial investment of £46m produced estimated savings of over £227m by reducing worker absences from 7% to 5% between 2004 and 2007.³⁷

from the corresponding author) and declare: *The State of Men's Health in Europe* was funded by the European Commission; they have no financial relationships with any organisations that might have an interest in the submitted work in the previous three years, AW is a member of the executive of the European Men's Forum and chair of the Men's Health Forum (England).

Provenance and peer review: Not commissioned; externally peer reviewed.

- 1 Meryn S, Jadad AR. The future of men and their health. Are men in danger of extinction? *BMJ* 2001;323:1013-4.
- 2 European Commission. The state of men's health in Europe report. 2011. http://ec.europa.eu/health/population_groups/docs/men_health_report_en.pdf.
- 3 European Commission. The state of men's health in Europe: extended report. EC, 2011. http://ec.europa.eu/health/population_groups/docs/men_health_extended_en.pdf.
- 4 McKee M, Shkolnikov V. Understanding the toll of premature death among men in eastern Europe. *BMJ* 2001;323:1051-5.
- 5 Bates LM, Hankivsky O, Springer KW. Gender and health inequities: a comment on the Final Report of the WHO Commission on the Social Determinants of Health. *Soc Sci Med* 2009;69:1002-4.
- 6 Lohan M. How might we understand men's health better? Integrating explanations from critical studies on men and inequalities in health. *Soc Sci Med* 2007;65:493-504.
- 7 Courtenay WH. Constructions of masculinity and their influence on men's well-being: a theory of gender and health. *Soc Sci Med* 2000;50:1385-401.
- 8 Gough B, Conner MT. Barriers to healthy eating amongst men: a qualitative analysis. *Soc Sci Med* 2006;62:387-95.
- 9 De Visser RO, Smith JA, McDonnell EJ. "That's not masculine": masculine capital and health-related behaviour. *J Health Psychol* 2009;14:1047-58.
- 10 Galdas P, Cheater MF, Marshall P. Men and health help-seeking behaviour: literature review. *J Adv Nurs* 2005;49:616-23.
- 11 White A, de Sousa B, de Visser R, Hogston R, Madsen SA, Makara P, et al. Men's health in Europe. *J Men's Health* 2011;8:192-201.
- 12 Hajdu P, McKee M, Bojan F. Changes in premature mortality differentials by marital status in Hungary and in England and Wales. *Eur J Publ Health* 1995;5:259-64.
- 13 Andreev E, McKee M, Shkolnikov V. Health expectancy in the Russian Federation: a new perspective on the health divide in Europe. *Bull World Health Organ* 2003;81:778-89.
- 14 Department of Health. Improving outcomes: a strategy for cancer. DH, 2011.
- 15 Danish National Board of Health. Mænds sundhed—en oversigt over mænds sundhedstilstand og en gennemgang af effektive forebyggelsesmetoder [Men's health: An overview of men's health and review of effective prevention methods]. Danish National Board of Health, 2010.
- 16 Department of Health and Children. National men's health policy: working with men in Ireland to achieve optimum health and wellbeing. DHC, 2008.

- 17 Richardson N, Smith J. National men's health policies in Ireland and Australia: what are the challenges associated with transitioning from development to implementation. *Public Health* 2011;125:424-32.
- 18 World Health Organization. Health for All database. 2011. <http://data.euro.who.int/hfad/>.
- 19 Zatonski WA, McMichael AJ, Powles JW. Ecological study of reasons for sharp decline in mortality from ischaemic heart disease in Poland since 1991. *BMJ* 1998;316:1047-51.
- 20 Lloyd T, Forrest S. Boys' and young men's health: literature and practice review. NHS Health Development Agency, 2001.
- 21 Virtanen P, Vahtera J, Kivimäki M, Pentti J, Ferrie J. Employment security and health. *J Epidemiol Community Health* 2002;56:569-74.
- 22 Hammarström A, Janlert U. Early unemployment can contribute to adult health problems: results from a longitudinal study of school leavers. *J Epidemiol Community Health* 2002;56:624-30.
- 23 Wilkinson R, Marmot M. Social determinants of health. The solid facts, 2nd ed. WHO, 2003.
- 24 World Health Organization. Fatherhood and health outcomes in Europe. WHO Europe, 2007.
- 25 Settersten R Jr, Cancel-Tirado D. Fatherhood as a hidden variable in men's development and life courses. *Res Hum Dev* 2010;7:83-102.
- 26 Robroek SJW, van Lenthe F, van Empelen P, Burdorf A. Determinants of participation in worksite health promotion programmes: a systematic review. *Int J Behav Nutr Phys Act* 2009;6:26.
- 27 Möller-Leimkühler AM. Barriers to help seeking by men: a review of sociocultural and clinical literature with particular reference to depression. *J Affect Disord* 2002;71:1-9.
- 28 Robertson LM, Douglas F, Ludbrook A, Reid G, van Teijlingen E. What works with men? A systematic review of health promoting interventions targeting men. *BMC Health Serv Res* 2008;8:141.
- 29 Marcell AV, Ford CA, Pleck JH, Sonenstein FL, et al. Masculine beliefs, parental communication, and male adolescents' health care use. *Pediatrics* 2007;119:e966-75.
- 30 Leishman J, Dalziel A. Taking action to improve the health of Scottish Men. *Men's Health J* 2003;2:90-3.
- 31 Pringle A, Zwolinsky S, Smith A, Robertson S, McKenna J, White A. The pre-adoption demographic and health profiles of men participating in a programme of men's health delivered in English Premier League football clubs. *Public Health* 2011;125:411-6.
- 32 Ormsby J, Stanley M, Jaworski K. Older men's participation in community-based men's sheds programmes. *Health Soc Care Community* 2010;18:607-13.
- 33 De Visser RO, Smith JA. Alcohol consumption and masculine identity among young men. *Psychol Health* 2007;22:595-614.
- 34 White A, Richardson N. Gendered epidemiology: making men's health visible in epidemiological research. *Public Health* 2011;125:407-10.
- 35 Douglas F, Amaya M, Greener J, Ludbrook A, Reid G, Robertson L, et al. Evaluation of the well men health service pilots. Scottish Government Social Research, 2008.
- 36 Communications Workers Union, BT Workfit. www.cwu.org/bt-workfit.html.
- 37 Marsden D, Moriconi S. The value of rude health: report for Royal Mail Group. London School of Economics, 2008.

Cite this as: *BMJ* 2011;343:d7397

© BMJ Publishing Group Ltd 2011