| Data captured | Study A ¹⁹ | Study B ⁹ | Study C ¹¹ |
|-----------------------------|------------------------|-----------------------|--------------------------|
| How was moderate-to- | SASSAD of ≥10 | ≥3 flare ups in | Nottingham Eczema |
| severe eczema | | previous six months | Severity Scale (NESS) of |
| defined? | | | ≥9 |
| Definition of WCW | WCW: | WCW: | WCW: |
| used§ | ≤ two days where | ≤ two days where | ≤ two days when |
| | stepping up of | stepping up of | topical corticosteroids |
| | treatment was | treatment was | used and global bother |
| | required and ≤ two | required and ≤ two | score ≤4 for the week |
| | days with a bother | days with a bother | |
| | score >4 | score >4 | |
| Bother score (0-10) | Daily in paper diaries | Daily in electronic | Bother over the last |
| "How much bother did | | diaries | week assessed at |
| your (your child's) | | | baseline, weeks 8, 16 |
| eczema cause today?" | | | and 24 |
| Scratch score (0-10) | Daily | Daily | - |
| "How much did you | | | |
| (your child) scratch | | | |
| today? | | | |
| Treatment "stepped | Daily | Daily | - |
| up" (Individually | | | |
| defined at the start of | | | |
| the study) (Yes/No) | | | |
| Topical corticosteroid | Daily (Yes/No) | - | Weekly |
| use (Yes/No) | | | (number of days in the |
| | | | last week) |
| Topical calcineurin | Daily | - | Weekly |
| inhibitor use (Yes/No) | | | (number of days in the |
| | | | last week) |
| POEM scores ¹³ | Baseline, weeks 4, | Baseline, weeks 4, 8, | Weekly |
| (range 0-28)* | 12 and 16 | 12, 16, 20 and 24 | |
| TIS scores ¹⁵ | weeks 12 and 16 | Baseline, weeks 4, 8, | Baseline, weeks 8, 16 |
| (range 0-9)* | | 12, 16, 20 and 24 | and 24 |
| SASSAD scores ¹⁴ | At weeks 12 and 16 | - | - |
| (range 0-108)* | | | |
| EASI scores ²⁰ | - | - | Baseline, weeks 8, 16 |
| (range 0-72)* | | | and 24 |

Table 1. Data available from included studies

* Higher scores denote more severe disease

§ Higher WCW scores represents better disease control