

# Candidate Polymorphisms and Severe Malaria in a Malian Population

Ousmane Toure<sup>1</sup>, Salimata Konate<sup>1</sup>, Sibiri Sissoko<sup>1</sup>, Amadou Niangaly<sup>1</sup>, Abdoulaye Barry<sup>1,2</sup>, Abdourahmane H. Sall<sup>1,2</sup>, Elisabeth Diarra<sup>1,2</sup>, Belco Poudiougou<sup>1</sup>, Nuno Sepulveda<sup>3</sup>, Susana Campino<sup>4</sup>, Kirk A. Rockett<sup>5</sup>, Taane G. Clark<sup>3,6\*</sup>, Mahamadou A. Thera<sup>1,6\*</sup>, Ogobara Doumbo<sup>1,6\*</sup>, in collaboration with The MalariaGEN Consortium<sup>4,5</sup>

**1** Malaria Research and Training Centre, Faculty of Medicine, Pharmacy and Dentistry, University of Bamako, Bamako, Mali, **2** Centre hospitalier Universitaire Gabriel Toure Hospital, Bamako, Mali, **3** Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, United Kingdom, **4** Wellcome Trust Sanger Institute, Hinxton, Cambridge, United Kingdom, **5** Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, United Kingdom, **6** Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom

## Abstract

Malaria is a major health burden in sub-Saharan African countries, including Mali. The disease is complex, with multiple genetic determinants influencing the observed variation in response to infection, progression, and severity. We assess the influence of sixty-four candidate loci, including the sickle cell polymorphism (HbS), on severe malaria in a case-control study consisting of over 900 individuals from Bamako, Mali. We confirm the known protective effects of the blood group O and the HbS AS genotype on life-threatening malaria. In addition, our analysis revealed a marginal susceptibility effect for the CD40 ligand (CD40L)+220C allele. The lack of statistical evidence for other candidates may demonstrate the need for large-scale genome-wide association studies in malaria to discover new polymorphisms. It also demonstrates the need for establishing the region-specific repertoire of functional variation in important genes, including the glucose-6-phosphatase deficiency gene, before embarking on focused genotyping.

**Citation:** Toure O, Konate S, Sissoko S, Niangaly A, Barry A, et al. (2012) Candidate Polymorphisms and Severe Malaria in a Malian Population. PLoS ONE 7(9): e43987. doi:10.1371/journal.pone.0043987

**Editor:** David Joseph Diemert, The George Washington University Medical Center, United States of America

**Received:** April 15, 2012; **Accepted:** July 26, 2012; **Published:** September 5, 2012

**Copyright:** © 2012 Toure et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The MalariaGEN Project is supported by the Wellcome Trust (WT077383/Z/05/Z) and by the Foundation for the National Institutes of Health (566) as part of the Bill & Melinda Gates' Grand Challenges in Global Health Initiative. The Resource Centre for Genomic Epidemiology of Malaria is supported by the Wellcome Trust (090770/Z/09/Z). This research was supported by the Medical Research Council (G0600718; G0600230). The Wellcome Trust also provided core awards to the Wellcome Trust Centre for Human Genetics (075491/Z/04; 090532/Z/09/Z) and the Wellcome Trust Sanger Institute (077012/Z/05/Z; G0600230), the Wellcome Trust Biomedical ethics Enhancement Award (087285) and Strategic Award (096527). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: taane.clark@shtm.ac.uk (TC); mthera@icermali.org (MT); okd@icermali.org (OD)

† These authors contributed equally to this work.

## Introduction

Malaria is a life-threatening parasitic disease transmitted by mosquitoes. Despite the concerted and renewed efforts to control the disease, it still persists as a major health burden, being responsible for 655,000 deaths in 2010, mainly children in Sub-Saharan Africa [1]. In Mali, there are over 800,000 recorded cases of malaria among its ~14 million people every year, and it accounts for 17 percent of child deaths [1]. Malaria is a complex disease with many genetic and environmental determinants influencing the observed variation in response to infection, progression and severity. Several factors are important for these different phenotypes observed, such as parasite genetic make-up, and host age, state of immunity and genetic background [2]. It has been estimated that 25% of the total variation in mild and severe malaria in a Kenya cohort was explained by host genes [2]. The different geographic distributions of sickle-cell disease,  $\alpha$  thalassaemia, glucose-6-phosphatase deficiency (G6PD), ovalocytosis, and the Duffy-negative blood group are examples of the general principle that different populations have evolved different genetic

variants to protect against malaria (see [3], for a review). The most striking example is the beta-globin HBB gene, in which three different coding SNPs confer protection against malaria: Glu6Val (HbS), Glu6Lys (HbC), and Glu26Lys (HbE). The HbS allele is common in Africa but rare in Southeast Asia, whereas the opposite is true for the HbE allele. However, a more complex picture emerges at the local level, exemplified by the Dogon people of Mali, who have a much lower frequency of the HbS allele than do most other West African groups and instead have a high frequency of the HbC allele [4]. Striking differences in response to malaria infection have also been observed among ethnic groups who live in the same geographical region. For example, it has been observed that the Fulani of Burkina Faso [5] and of Mali [6] have a significantly lower prevalence of malaria parasitaemia and fewer malaria clinical attacks, when compared to other ethnic groups living in neighbouring villages. In addition to the sickle polymorphism (HbS) [7], G6PD (reviewed in [8]), and ABO blood group [9,10], a number of candidate polymorphisms have been proposed for the reduced risk of severe malaria. For example, these include genes that are relevant to immunity and inflammation such as the

tumour necrosis factor (TNF, MHC class III region, reviewed in [11]), Toll-like receptors (TLR-4,9) [12], CD40 ligand (CD40L) [13], the interferon gamma (IFNG) (reviewed in [14], and the Nitric oxide synthase type 2 (NOS2A) genes (reviewed in [15]). Here we investigate whether a number malaria candidate SNPs, including the HbC, HbS and ABO, are associated with severe malaria. Our study is the first to survey malaria candidate SNPs in a Malian population, and we seek to confirm genetic associations found in other studies. We consider a cohort of over 900 individuals recruited in Bamako, predominantly from the Bambara ethnicity, which is under-represented in other genetic epidemiological studies in Western Africa.

## Methods

### Participants, Materials and Methods

**Ethics Statement.** This study was approved by the Faculty of Medicine, Pharmacy and Dentistry (University of Bamako) Ethics Review Committee. All clinical and biological samples were collected and DNA was genotyped following approval by this committee. Written informed consent was obtained from the next of kin, carers or guardians on the behalf of the minors/children participants involved in this study.

**Study participants.** Patient samples were collected as part of ongoing epidemiological studies of severe malaria at the Centre Hospitalier Universitaire Gabriel Toure, Bamako, Mali (malaria cases 541 (57.9%); healthy controls 393 (42.1%)). They had a median age of ~3 years, and were predominantly from the Bambara ethnic group (53%) (see Table 1).

**Phenotypic definition.** All cases were children admitted to hospital with evidence of *P. falciparum* on blood film and clinical features of uncomplicated and severe malaria [16,17]. Subjects were defined as having had cerebral malaria (CM) if their Blantyre coma score was less than or equal to 2 on presentation or early during admission. A second phenotypic subset of severe malarial

anaemia (SMA) was defined as those subjects having had a haemoglobin concentration of less than 5 g/dl or a haematocrit less than 15%. Participants with co-existing severe or chronic medical conditions (e.g. bacterial pneumonia, kwashiorkor) unrelated to a severe malarial infection were excluded. Most cases had severe malaria (n = 458, 84.7%), but a minority had uncomplicated conditions (n = 83, 15.3%) (see Table 1). Controls were healthy individuals matched for age, ethnicity and residence to severe malaria cases (Table 1). For the purpose of analysis, we either excluded uncomplicated malaria cases or included them as part of the control group. Because of restrictions in sample size, we do not present an analysis on different ethnic groups or specific sub-clinical phenotypes.

**Sample preparation and genotyping.** Genomic DNA samples underwent whole genome amplification through Primer Extension Pre-amplification (PEP) [18], before genotyping on a Sequenom MassArray genotyping platform [19,20]. Sixty-four malaria candidate SNPs were genotyped, including: Haemoglobin variants C (HbC, rs33930165) and S (HbS, rs334), plus two SNPs that allow an estimate of the ABO blood group [10]. The rs8176719 derived allele results in a non-functional enzyme, and group O individuals are DD, while non-O Individuals are either II or ID. In addition, rs8176746 is involved in the enzyme's substrate selection and therefore defines either the A or B blood groups. The two Sequenom iPLEX reactions designed also included gender-typing SNPs. The selection of SNPs for genotyping was undertaken by the MalariaGEN Consortium, and were selected by interrogation of the literature and ongoing consortial experiments for evidence of association with severe malaria. Full details of polymorphisms can be found at [www.malariagen.net](http://www.malariagen.net), and a list of SNPs typed can also be found in Tables 2 and S1.

**Statistical analysis.** Genotypic deviations from Hardy-Weinberg equilibrium (HWE) were assessed using a chi-square statistical test. SNPs were excluded from analysis if they had at least 10% of genotype calls missing or there was significant deviation from HWE ( $p < 0.001$ ). Case-control association analysis using SNP alleles/genotypes was undertaken by logistic regression and included the covariates: ethnic group and the HbS polymorphism. In this approach we modelled the SNP of interest assuming several related genotypic mechanisms (additive, dominant, recessive, heterozygous advantage and general models) and reported the minimum p-value from these correlated tests. All analyses were performed using the R statistical package (<http://www.r-project.org>). Performing multiple statistical tests leads to inflation in the occurrence of false positives and using a permutation approach that accounted for correlation between markers and tests, we estimated a p-value cut-off of 0.002 to be statistically significant. Allele frequency differences were estimated using the *Fst* metric, with values potentially varying from zero (no population differentiation) to one (complete differentiation) [22].

## Results

Six SNPs were removed from the analysis because they were monomorphic (rs33950507, rs5743611, rs2814778), deviated from HWE in controls (rs2227478, rs2535611) or had high rates of missing genotype calls (rs1801274). Allelic-based tests revealed potential associations of HbS polymorphism (rs334, HBB gene), the O blood group (and its components rs8176746 and rs8176719), and rs1126535 (CD40L+220) with severe malaria ( $P < 0.002$ ) (Table 2). Figure 1 shows the minimum p-values from the genotypic tests applied to the autosomal SNPs, and confirms that the sickle cell (HbS) and ABO polymorphisms (rs8176746, rs8176719) are significantly associated with severe malaria. This

**Table 1.** Baseline and clinical characteristics.

|                       | Controls (n = 393) |             | Cases (n = 541) |             |
|-----------------------|--------------------|-------------|-----------------|-------------|
| Age* (median, range)  | (38.0)             | (4.0–178.0) | (36.0)          | (2.0–173.0) |
| Gender - male         | 195                | 49.6%       | 300             | 55.5%       |
| Bamako residence      | 140                | 35.6%       | 200             | 37.0%       |
| Ethnicity             |                    |             |                 |             |
| Bambara               | 219                | 55.7%       | 272             | 50.3%       |
| Malinke               | 55                 | 14%         | 83              | 15.3%       |
| Other                 | 48                 | 12.2%       | 78              | 14.4%       |
| Peulh                 | 36                 | 9.2%        | 56              | 10.4%       |
| Sarakole              | 35                 | 8.9%        | 52              | 9.6%        |
| Clinical phenotype    |                    |             |                 |             |
| Uncomplicated malaria | -                  | -           | 83              | 15.3%       |
| Any severe malaria    | -                  | -           | 458             | 84.7%       |
| Any SMA               | -                  | -           | 304             | 66.4%       |
| Any CM                | -                  | -           | 350             | 75.1%       |
| Both SMA+CM           | -                  | -           | 247             | 53.0%       |
| Any RD                | -                  | -           | 200             | 42.9%       |
| Malaria death         | -                  | -           | 72              | 15.7%       |

\*in months, SMA = severe malarial anaemia, CM = cerebral malaria, RD = respiratory distress.

doi:10.1371/journal.pone.0043987.t001

**Table 2.** Single nucleotide polymorphisms.

| SNP          | MajA     | MinA     | MAF          | Controls     | Cases        | HWE P         | OR           | LCL          | UCL          | P             |
|--------------|----------|----------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|---------------|
| rs3024500    | G        | A        | 0.363        | 0.367        | 0.358        | 0.3800        | 0.963        | 0.793        | 1.170        | 0.7079        |
| rs1800896    | C        | T        | 0.332        | 0.338        | 0.327        | 0.4408        | 0.951        | 0.781        | 1.158        | 0.6151        |
| rs1800890    | T        | A        | 0.185        | 0.195        | 0.175        | 0.0183        | 0.875        | 0.689        | 1.112        | 0.2743        |
| rs17047660   | G        | A        | 0.337        | 0.331        | 0.342        | 0.6354        | 1.047        | 0.860        | 1.275        | 0.6490        |
| rs17047661   | A        | G        | 0.219        | 0.223        | 0.216        | 0.5149        | 0.959        | 0.766        | 1.201        | 0.7182        |
| rs1803632    | C        | G        | 0.456        | 0.447        | 0.465        | 0.4712        | 1.075        | 0.892        | 1.295        | 0.4492        |
| <b>rs334</b> | <b>A</b> | <b>S</b> | <b>0.024</b> | <b>0.038</b> | <b>0.010</b> | <b>0.6769</b> | <b>0.255</b> | <b>0.122</b> | <b>0.533</b> | <b>0.0003</b> |
| rs33930165   | G        | A        | 0.054        | 0.059        | 0.049        | 0.1788        | 0.824        | 0.540        | 1.256        | 0.3669        |
| rs7935564    | G        | A        | 0.474        | 0.479        | 0.468        | 0.6407        | 0.955        | 0.792        | 1.152        | 0.6319        |
| rs542998     | T        | C        | 0.411        | 0.436        | 0.385        | 0.2672        | 0.810        | 0.669        | 0.980        | 0.0304        |
| rs2227507    | T        | C        | 0.022        | 0.022        | 0.021        | 0.6289        | 0.960        | 0.509        | 1.811        | 0.8994        |
| rs1012356    | A        | T        | 0.487        | 0.498        | 0.475        | 0.1593        | 0.912        | 0.758        | 1.099        | 0.3331        |
| rs2227491    | T        | C        | 0.312        | 0.314        | 0.309        | 0.1876        | 0.974        | 0.797        | 1.190        | 0.7972        |
| rs2227485    | G        | A        | 0.486        | 0.497        | 0.476        | 0.4513        | 0.920        | 0.764        | 1.109        | 0.3814        |
| rs229587     | C        | T        | 0.286        | 0.266        | 0.305        | 0.3735        | 1.211        | 0.981        | 1.493        | 0.0742        |
| rs1805015    | C        | T        | 0.484        | 0.488        | 0.479        | 0.0791        | 0.966        | 0.801        | 1.164        | 0.7152        |
| rs2230739    | G        | A        | 0.168        | 0.167        | 0.169        | 0.2478        | 1.014        | 0.791        | 1.300        | 0.9145        |
| rs10775349   | G        | C        | 0.163        | 0.158        | 0.167        | 0.9521        | 1.065        | 0.828        | 1.371        | 0.6229        |
| rs2297518    | A        | G        | 0.085        | 0.081        | 0.090        | 0.2139        | 1.112        | 0.797        | 1.551        | 0.5323        |
| rs1800482    | C        | G        | 0.092        | 0.089        | 0.096        | 0.1582        | 1.077        | 0.782        | 1.484        | 0.6500        |
| rs9282799    | T        | C        | 0.067        | 0.063        | 0.071        | 0.0786        | 1.146        | 0.791        | 1.660        | 0.4722        |
| rs8078340    | T        | C        | 0.251        | 0.257        | 0.245        | 0.3906        | 0.939        | 0.758        | 1.164        | 0.5659        |
| rs1799969    | G        | A        | 0.001        | 0.001        | 0.001        | 0.9812        | 1.035        | 0.065        | 16.566       | 0.9809        |
| rs5498       | G        | A        | 0.125        | 0.124        | 0.126        | 0.3716        | 1.018        | 0.769        | 1.348        | 0.9005        |
| rs373533     | T        | G        | 0.375        | 0.354        | 0.397        | 0.0643        | 1.200        | 0.989        | 1.456        | 0.0647        |
| rs461645     | T        | C        | 0.380        | 0.360        | 0.399        | 0.1011        | 1.180        | 0.974        | 1.429        | 0.0912        |
| rs17561      | T        | G        | 0.177        | 0.178        | 0.176        | 0.6004        | 0.986        | 0.774        | 1.257        | 0.9110        |
| rs1143634    | T        | C        | 0.120        | 0.113        | 0.126        | 0.4473        | 1.133        | 0.848        | 1.513        | 0.3977        |
| rs8386       | T        | C        | 0.146        | 0.145        | 0.148        | 0.3342        | 1.024        | 0.788        | 1.331        | 0.8578        |
| rs1128127    | G        | A        | 0.436        | 0.424        | 0.448        | 0.1413        | 1.102        | 0.912        | 1.332        | 0.3133        |
| rs187084     | C        | T        | 0.238        | 0.243        | 0.232        | 0.0059        | 0.944        | 0.755        | 1.182        | 0.6171        |
| rs6780995    | G        | A        | 0.459        | 0.455        | 0.463        | 0.9004        | 1.032        | 0.856        | 1.244        | 0.7443        |
| rs708567     | G        | A        | 0.483        | 0.462        | 0.503        | 0.4465        | 1.183        | 0.977        | 1.431        | 0.0849        |
| rs4833095    | T        | C        | 0.085        | 0.078        | 0.093        | 0.3797        | 1.220        | 0.873        | 1.704        | 0.2450        |
| rs5743810    | C        | T        | 0.002        | 0.002        | 0.001        | 0.9622        | 0.503        | 0.046        | 5.555        | 0.5748        |
| rs5743809    | T        | C        | 0.047        | 0.041        | 0.054        | 0.7487        | 1.329        | 0.852        | 2.073        | 0.2096        |
| rs2706384    | C        | A        | 0.435        | 0.449        | 0.422        | 0.0082        | 0.895        | 0.739        | 1.084        | 0.2579        |
| rs20541      | C        | T        | 0.176        | 0.173        | 0.177        | 0.3304        | 1.028        | 0.789        | 1.341        | 0.8361        |
| rs2243250    | T        | C        | 0.242        | 0.244        | 0.240        | 0.9126        | 0.979        | 0.786        | 1.219        | 0.8496        |
| rs1801033    | A        | C        | 0.445        | 0.432        | 0.457        | 0.9663        | 1.105        | 0.916        | 1.333        | 0.2951        |
| rs1555498    | C        | T        | 0.436        | 0.426        | 0.446        | 0.3849        | 1.089        | 0.903        | 1.313        | 0.3740        |
| rs2239704    | G        | T        | 0.378        | 0.383        | 0.372        | 0.8615        | 0.951        | 0.785        | 1.153        | 0.6093        |
| rs909253     | T        | C        | 0.419        | 0.411        | 0.426        | 0.1203        | 1.063        | 0.878        | 1.287        | 0.5316        |
| rs1799964    | T        | C        | 0.140        | 0.144        | 0.136        | 0.7781        | 0.942        | 0.720        | 1.232        | 0.6633        |
| rs1800750    | G        | A        | 0.014        | 0.013        | 0.015        | 0.7744        | 1.115        | 0.506        | 2.457        | 0.7876        |
| rs1800629    | A        | G        | 0.134        | 0.137        | 0.131        | 0.5698        | 0.947        | 0.721        | 1.243        | 0.6940        |
| rs361525     | G        | A        | 0.037        | 0.033        | 0.040        | 0.4682        | 1.242        | 0.758        | 2.035        | 0.3892        |
| rs3093662    | G        | A        | 0.072        | 0.067        | 0.078        | 0.4503        | 1.191        | 0.832        | 1.706        | 0.3401        |
| rs2242665    | G        | A        | 0.264        | 0.249        | 0.280        | 0.8738        | 1.175        | 0.950        | 1.452        | 0.1364        |

**Table 2.** Cont.

| SNP                 | MajA         | MinA     | MAF          | Controls     | Cases        | HWE P         | OR           | LCL          | UCL          | P                 |
|---------------------|--------------|----------|--------------|--------------|--------------|---------------|--------------|--------------|--------------|-------------------|
| rs17140229          | C            | T        | 0.393        | 0.387        | 0.399        | 0.7000        | 1.053        | 0.869        | 1.275        | 0.5988            |
| rs2075820           | A            | G        | 0.424        | 0.430        | 0.418        | 0.9496        | 0.954        | 0.790        | 1.152        | 0.6250            |
| rs3211938           | G            | T        | 0.143        | 0.157        | 0.129        | 0.0345        | 0.793        | 0.607        | 1.037        | 0.0900            |
| hCD36_G1439C        | C            | G        | 0.055        | 0.060        | 0.050        | 0.7433        | 1.216        | 0.805        | 1.836        | 0.3520            |
| rs4986790           | G            | A        | 0.110        | 0.120        | 0.100        | 0.0436        | 0.819        | 0.608        | 1.104        | 0.1893            |
| <b>rs8176746</b>    | <b>A</b>     | <b>C</b> | <b>0.244</b> | <b>0.213</b> | <b>0.276</b> | <b>0.0093</b> | <b>1.412</b> | <b>1.135</b> | <b>1.757</b> | <b>0.0020</b>     |
| <b>rs8176719</b>    | <b>I</b>     | <b>D</b> | <b>0.403</b> | <b>0.364</b> | <b>0.442</b> | <b>0.8022</b> | <b>1.384</b> | <b>1.145</b> | <b>1.673</b> | <b>0.0008</b>     |
| <b>ABO</b>          | <b>Non-O</b> | <b>O</b> | <b>0.356</b> | <b>0.438</b> | <b>0.295</b> | <b>NA</b>     | <b>0.539</b> | <b>0.438</b> | <b>0.662</b> | <b>&lt;0.0001</b> |
| rs3092945 F         | T            | C        | 0.334        | 0.387        | 0.285        | 0.3588        | 0.631        | 0.465        | 0.856        | 0.0031            |
| rs3092945 M         | T            | C        | 0.346        | 0.320        | 0.365        | NA            | 1.219        | 0.913        | 1.627        | 0.1795            |
| rs3092945 Ov        | T            | C        | 0.338        | 0.352        | 0.327        | NA            | 0.893        | 0.724        | 1.101        | 0.2891            |
| rs1126535 F         | T            | C        | 0.128        | 0.095        | 0.158        | 0.5737        | 1.798        | 1.157        | 2.796        | 0.0092            |
| rs1126535 M         | T            | C        | 0.098        | 0.077        | 0.113        | NA            | 1.534        | 0.954        | 2.468        | 0.0776            |
| <b>rs1126535 Ov</b> | <b>T</b>     | <b>C</b> | <b>0.112</b> | <b>0.085</b> | <b>0.134</b> | <b>NA</b>     | <b>1.671</b> | <b>1.209</b> | <b>2.309</b> | <b>0.0019</b>     |
| rs1050829 F         | A            | G        | 0.423        | 0.401        | 0.443        | 0.9805        | 1.188        | 0.890        | 1.585        | 0.2421            |
| rs1050829 M         | A            | G        | 0.418        | 0.425        | 0.413        | NA            | 0.954        | 0.724        | 1.259        | 0.7412            |
| rs1050829 Ov        | A            | G        | 0.421        | 0.413        | 0.427        | NA            | 1.060        | 0.868        | 1.294        | 0.5673            |
| rs1050828 F         | G            | A        | 0.162        | 0.146        | 0.178        | 0.9615        | 1.267        | 0.862        | 1.861        | 0.2281            |
| rs1050828 M         | G            | A        | 0.183        | 0.162        | 0.198        | NA            | 1.278        | 0.896        | 1.824        | 0.1757            |
| rs1050828 Ov        | G            | A        | 0.173        | 0.154        | 0.189        | NA            | 1.273        | 0.981        | 1.653        | 0.0699            |

MinA = minor allele, MajA = major allele, MAF = overall minor allele frequency, HWE P is the Hardy-Weinberg p-value, OR = odds ratio, 95% Confidence interval (LCL, UCL), P = P-value; rs8176746 and rs8176719 are used to infer ABO blood groups; for X chromosome SNPs (rs3092945 (CD40), rs1126535 (CD40), rs1050829 (G6PD-376), rs1050828 (G6PD-202/A-), analyses are presented for separately for females (F) and males (M) and pooled to obtain overall results (Ov), NA not applicable; rs33950507, rs5743611, rs2814778, rs2227478, rs2535611 and rs1801274 did not pass quality control filters and are not presented.  
doi:10.1371/journal.pone.0043987.t002

genotypic analysis adjusting for the potentially confounding effects of ethnicity (and where appropriate HbS) supports the heterozygous advantage effect of the HbS-AS genotype (Odds ratio AS vs. AA: 0.03,  $P < 6 \times 10^{-10}$ ), and the reduced risk in the O blood group (Odds ratio O vs. A/B/AB: 0.58,  $P = 0.0003$ ) (Table 3). An increased risk from the CD40L+220-C allele was observed for males (Odds ratio C vs. A 2.12,  $P = 0.05$ ) and females (Odds ratio additive C model 1.67,  $P = 0.03$ ), and the pooled result was marginally non-significant (Odds ratio additive C model 1.79,  $P = 0.0045$ ) (Table 3). The G6PD-202 polymorphism has been used as a molecular surrogate for the A- deficiency (see [8,21] for a review). We found no strong evidence of a G6PD association with severe malaria risk in males, females or overall (Tables 2 and 3,  $P > 0.06$ ); the direction of the odds ratios suggests those with the 202-A (A-) allele were at increased (rather than a decreased) risk of the disease. All our results were insensitive to the inclusion or exclusion of uncomplicated malaria cases in the control group. There were no major allele frequency differences between the Bambara and other ethnic groups (median  $F_{st}$  0.001, only 3  $F_{st}$  values greater than 0.01, maximum 0.015).


## Discussion

In our study, we set out to investigate the role of candidate malaria polymorphisms on severe malaria risk in a predominantly Bambara population in and around Bamako. To minimise errors we standardised procedures using case report forms, pre-defined definitions of severe malaria and sub-clinical phenotypes, and ensured all samples underwent genotyping on the same Sequenom MassArray platform with resulting low rates of missing data

(<5%). In addition, all analyses were adjusted for ethnicity, minimising the confounding effects and potential false positives arising from population stratification.

Data analysis confirmed the known (~90%) reduced severe malaria risk from the sickle cell AS genotype [7]. The low frequency of the S allele in the controls (~3.8%) is in keeping with other populations (see <http://www.map.ox.ac.uk/>) in West (Burkina Faso 5.2%, Cameroon 6.5%, Gambia 7.6%, Ghana 6.5%) and East (Kenya 6.4%, Malawi 2.7%, Tanzania 7.8%) Africa. The frequency of the HbC allele was 5% in cases and 6% in controls, both greater than the HbS allele, but there was no strong evidence of association ( $P > 0.3$ ). The higher frequency of the HbC allele has been observed in other West African populations [4].

Our analysis also confirms the known protective effects of the blood group O on life threatening malaria, which is thought to act on malaria pathogenesis through the mechanism of reduced *P. falciparum* rosetting [9,10]. An insight into an underlying molecular mechanism could lead to the development of a new anti-malarial therapy. Our inability to detect associations found in other studies may be due to previously reported false-positives, and methodological issues such as variation in phenotype definition, choice of controls, village surveys versus hospital-based studies, possible heterogeneity in the parasite population, immune status of subjects, and study location. There may also be issues with sample size (statistical power). At the present sample size, we have 90% power to detect a dominant odds ratio effect of 1.5 with a minor allele frequency of 0.20 and type I error of 5%. Reducing the odds ratio to 1.4 or MAF to 0.13 would lead to 80% power. It is clear that much larger studies are required to detect more modest



**Figure 1. Minimum p-values from tests of association for the autosomal SNPs** \*. \* Genotypic tests of dominant, recessive, general, heterozygous advantage, and additive models, adjusted for HbS and ethnicity; in this analysis controls include uncomplicated malaria cases; the dashed line represents a p-value of 0.002. doi:10.1371/journal.pone.0043987.g001

effects, especially if relying on markers in LD with a causal untyped polymorphism. In this setting, dense SNP genome-wide association strategies are required to discover new candidate genes.

Our analysis revealed a marginal susceptibility effect for the CD40 ligand (CD40L)+220C allele. CD40L is a glycoprotein involved in B cell proliferation, antigen presenting cell activation, and Ig class switching, and therefore important in the immune response to infection [13]. Previous work in the Gambia found the

CD40L+220C allele had a non-significant susceptibility effect, and instead identified the genotyped CD40L-726 polymorphism with a significant reduction in risk for severe malaria in males-hemizygous [13]. Similarly, we expected to see a reduced malaria risk with G6PD-202A (A- deficiency), as shown in another Malian population [8] with similar allele frequencies in controls to our study. These discrepancies in CD40L and G6PD results may be due to allelic heterogeneity. For example, it has been shown that the G6PD-202 may not be a good marker of A- deficiency, and other polymorphisms are required to confirm the protective effect [21]. In addition, discrepancies may arise due to differences in linkage disequilibrium (LD) patterns between populations. For example, it is known there are at least five classical haplotypes surrounding the sickle polymorphism in the HBB gene, and the resulting differences in LD can make it difficult to localise HbS using indirect associations [7]. In addition, some functional polymorphisms may be distal from candidate genes and polymorphisms genotyped. For example, it has been suggested that the causal polymorphisms regulating TNF and LTA response may be some distance from the genes [11,23].

In conclusion, our work reinforces in a Malian (mainly Bambara) population the importance of the sickle cell polymorphism and ABO blood group on severe malaria susceptibility. It may also demonstrate the need for establishing the region-specific repertoire of functional variation in important genes such as G6PD, before embarking on focused genotyping. Proposed large-scale genomewide association studies and resequencing of important genes in a number of Malian (and other African)

**Table 3. Odds ratios\* for HbS, ABO, CD40 and G6PD-202.**

| Polymorphism                   | Contrast     | OR    | LCL   | UCL   | P      |
|--------------------------------|--------------|-------|-------|-------|--------|
| HbS (rs334)                    | AS vs. AA    | 0.028 | 0.004 | 0.209 | <6E-10 |
| Blood Group (rs8176719)        | O vs. A/B/AB | 0.575 | 0.425 | 0.777 | 0.0003 |
| CD40 (rs1126535) – Female      | Additive C   | 1.674 | 1.045 | 2.683 | 0.03   |
| CD40 (rs1126535) – Male        | Additive C   | 2.119 | 0.990 | 4.537 | 0.05   |
| CD40 (rs1126535) – Overall     | Additive C   | 1.787 | 1.197 | 2.669 | 0.0045 |
| G6PD (rs1050828, 202) – Female | AG vs. GG    | 1.187 | 0.747 | 1.887 | 0.467  |
| G6PD (rs1050828, 202) – Female | AA vs. GG    | 1.557 | 0.427 | 5.684 | 0.502  |
| G6PD (rs1050828, 202) – Male   | A vs. G      | 1.205 | 0.718 | 2.022 | 0.480  |

\*adjusted for HbS and ethnicity, OR = odds ratio, 95% Confidence interval (LCL, UCL), P = P-value, in this analysis controls include uncomplicated malaria cases. doi:10.1371/journal.pone.0043987.t003

populations is likely to generate insights into how *P. falciparum* has shaped our genome. These approaches could expose new candidate protective polymorphisms, resulting in functional work to understand underlying molecular mechanisms, leading to the development of therapies and vaccines for malaria control.

## Supporting Information

**Table S1** List of all candidate polymorphisms considered in our work.  
(DOC)

## References

- World Health Organisation (2011) World Malaria Report 2011. Available: [http://who.int/malaria/world\\_malaria\\_report\\_2011/en/index.html](http://who.int/malaria/world_malaria_report_2011/en/index.html). Accessed 2012 Aug 14.
- Mackinnon MJ, Mwangi TW, Snow RW, Marsh K, Williams TN (2005) Heritability of malaria in Africa. *PLoS Med* 2: e340.
- Campino S, Kwiatkowski D, Dessein A (2006) Mendelian and complex genetics of susceptibility and resistance to parasitic infections. *Semin Immunol* 18: 411–22.
- Agarwal A, Guindo A, Cissoko Y, Taylor JG, Coulibaly D, et al. (2000) Hemoglobin C associated with protection from severe malaria in the Dogon of Mali, a West African population with a low prevalence of hemoglobin S. *Blood* 96: 2358–63.
- Modiano D, Petrarca V, Sirima BS, Bosman A, Nebie I, et al. (1995) *Plasmodium falciparum* malaria in sympatric ethnic groups of Burkina Faso, west Africa. *Parassitologia* 37: 255–9.
- Dolo A, Modiano D, Maiga B, Daou M, Dolo G, et al. (2005) Difference in susceptibility to malaria between two sympatric ethnic groups in Mali. *Am J Trop Med Hyg* 72: 243–8.
- Jallow M, Teo YY, Small KS, Rockett KA, Deloukas P, et al. (2009) Genome-wide and fine-resolution association analysis of malaria in West Africa. *Nat Genet* 41(6): 657–665.
- Guindo A, Fairhurst RM, Doumbo OK, Wellem TE, Diallo DA (2007) X-linked G6PD deficiency protects hemizygous males but not heterozygous females against severe malaria. *PLoS Med* 4: e66.
- Rowe JA, Handel IG, Thera MA, Deans AM, Lyke KE, et al. (2007) Blood group O protects against severe *Plasmodium falciparum* malaria through the mechanism of reduced rosetting. *Proc Natl Acad Sci U S A* 104: 17471–17476.
- Fry AE, Griffiths MJ, Auburn S, Diakite M, Forton JT, et al. (2008) Common variation in the ABO glycosyltransferase is associated with susceptibility to severe *Plasmodium falciparum* malaria. *Hum Mol Genet* 17: 567–76.
- Clark TG, Diakite M, Auburn S, Campino S, Fry AE (2009) Tumor necrosis factor and lymphotoxin-alpha polymorphisms and severe malaria in African populations. *J Infect Dis* 199: 569–75.
- Mockenhaupt FP, Hamann L, von Gaertner C, Bedu-Addo G, von Kleinsorgen C (2006) Common polymorphisms of toll-like receptors 4 and 9 are associated with the clinical manifestation of malaria during pregnancy. *J Infect Dis* 194: 184–8.
- Sabeti P, Usen S, Farhadian S, Jallow M, Doherty T, et al. (2002) CD40L association with protection from severe malaria. *Genes Immun* 5: 286–291.
- Stevenson MM, Riley EM (2004) Innate immunity to malaria. *Nat Rev Immunol* 4: 169–80.
- Clark IA, Rockett KA (1996) Nitric oxide and parasitic disease. *Adv Parasitol* 37: 1–56.
- Marsh K, Forster D, Waruiru C, Mwangi I, Winstanley M, et al. (1995) Indicators of life-threatening malaria in African children. *N Engl J Med* 332: 1399–404.
- World Health Organisation (1990) Severe and complicated malaria. World Health Organization, Division of Control of Tropical Diseases. *Trans R Soc Trop Med Hyg* 84 Suppl 2: 1–65.
- Zhang L, Cui X, Schmitt K, Hubert R, Navidi W, et al. (1992) Whole genome amplification from a single cell: implications for genetic analysis. *Proc Natl Acad Sci U S A* 89: 5847–51.
- Wilson JN, Rockett K, Jallow M, Pinder M, Sisay-Joof F, et al. (2005) Analysis of IL10 haplotypic associations with severe malaria. *Genes Immun* 6: 462–6.
- Ross P, Hall L, Smirnov I, Haff L (1998) High level multiplex genotyping by MALDI-TOF mass spectrometry. *Nat Biotechnol* 16: 1347–51.
- Clark TG, Fry A, Auburn S, Campino S, Diakite M, et al. (2009) G6PD deficiency and severe malaria: unrecognized allelic heterogeneity is confounding association studies in Africa. *Eur J Human Genetics* 17: 1080–5.
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38(6): 1358–1370.
- Diakite M, Clark TG, Auburn S, Campino S, Fry AE, et al. (2009) A genetic association study in the Gambia using tagging polymorphisms in the major histocompatibility complex (MHC) class III region implicates a BAT2 polymorphism in severe malaria susceptibility. *Human Genetics* 125: 105–9.

## Acknowledgments

We thank the participants and Bamako communities who made this study possible; we thank Professor Mamadou Marouf Keita and his team of healthcare workers at the Centre Hospitalier Universitaire Gabriel Toure who assisted with this work.

## Author Contributions

Conceived and designed the experiments: OT KR TC MT OD. Performed the experiments: OT SK SS AN AB AS ED BP. Analyzed the data: TC NS SC OT. Contributed reagents/materials/analysis tools: OD MalariaGen TC. Wrote the paper: TC OT OD MT KR SC.