

Introduction

The discovery of norovirus in 1972¹, changed the understanding of the aetiology of gastroenteritis, making it the virus to be identified as an agent of gastroenteritis in humans.

Today, norovirus is recognised as one of the commonest human infections and estimated to be associated with 125 million cases and 35,000 deaths worldwide in 2010². Better epidemiological surveillance and outbreak investigations³, coupled with wider implementation of molecular-based laboratory diagnostics⁴ are leading to better estimates of the burden of norovirus infections as well as improved outbreak control.

Data from challenge studies of prototype norovirus vaccines^{5, 6} demonstrated that protection against infection and disease can be achieved, however there remain significant challenges to development of a norovirus vaccine. Recent advances in cell culture systems for norovirus^{7, 8} and current research investigating the distribution of norovirus-associated disease in the population, for whom the disease burden is greatest, understanding host susceptibility factors, how to deploy novel technologies detecting norovirus in food and environmental matrices, and methodologies for ascertaining cases, are important in increasing our understanding of norovirus. Answers to these will help design strategies for vaccine and antiviral development, and how these might be best deployed to control norovirus infection.

Norovirus virology

At the time of discovery, the virus was referred to as the Norwalk agent, but as other related viruses were described in association with gastroenteritis, they became known as Norwalk-like viruses (NLVs), or – based on their morphology by electron microscopy – small, round, structured viruses (SLSVs). Following the cloning and sequencing of the Norwalk agent genome in 1993⁹, and subsequently other NLVs¹⁰, defining the genetic relatedness of these viruses led to their reclassification in the *Caliciviridae* family of viruses under the genus *Norovirus* in 2002¹¹.

Classification of this genetically diverse group of viruses^{12, 13} has described six established¹⁴ genogroups (GI-GVI), and a proposed seventh^{15, 16}. Two genogroups (GI and GII) are important pathogens of humans (GII also contains pathogens of animals, but there is no evidence of zoonotic transmission¹⁷) and genogroups are further subdivided into genotypes: nine GI and twenty-two GII genotypes have been described¹²⁻¹⁴. Norovirus names are presented as genotypes, e.g. genogroup II-genotype 4 (GII.4) and strains are named for the place and year of their first description: e.g. GII.4/New Orleans 2009¹⁸ or GII.4/Sydney 2012¹⁹.

The norovirus genome is a single strand of positive-sense ribonucleic acid (+ssRNA) that is ~7700 nucleotides in length, organised as three open reading frames (ORF1-3)²⁰. The 5'-proximal ORF1 encodes a polyprotein that is post-translationally processed by the virus-encoded protease into six non-structural proteins, including a genome-linked protein (VPg/NS5), protease (Pro/NS6) and an RNA-dependent RNA polymerase (RdRp/NS7). Both ORF2 and ORF3

encode a single protein each, VP1 and VP2 respectively, that are structural proteins involved in formation and stabilisation of the virus particle²¹.

Norovirus particles are 27-35nm in diameter and comprised of 180 copies of the VP1 protein, which itself is organised into three main domains: N-terminal (N), shell (S) and protruding (P), which is further arranged as P1 (subdivided as P1.1 and P1.2) and P2²¹. In the mature infectious virus particle, the N domain is internal, and the P2 domain is the most external part of the virus, making it highly surface-exposed and placed to coordinate many of the interactions between norovirus and its host environment.

The primary host cell receptor for human norovirus is unknown, but can interact with histo-blood group antigens (HBGAs)²²; these are glycans expressed on epithelial cells and in mucosal secretions, which determine ABO blood type groups. Norovirus strains may use HBGAs as attachment factors or co-receptors²³, and sites in the VP1 P2 domain have been identified as HBGA binding sites²⁴⁻²⁹. Synthesis of HBGAs occurs by sequential modification of a precursor, and the process is controlled by glycosyl-transferase enzymes from several genetic loci that exhibit polymorphism throughout the human population. The ABH and Lewis antigens are relevant to norovirus binding, and as such the phenotype of an individual for secretory ABH and Lewis antigens is a host susceptibility factor for norovirus infection. Specifically, individuals with a non-functional *FUT2* gene, which encodes an α 1,2-fucosyltransferase, have a 'non-secretor' phenotype, and are more resistant to norovirus infection than 'secretor' individuals^{30, 31}. Polymorphism at this locus may modulate susceptibility to other causes of diarrhoeal disease³².

Public health laboratory surveillance worldwide has demonstrated dominance of GII.4 viruses³³⁻³⁶, however other norovirus genotypes circulate consistently, if at a lower level, in the population. The GII.4 cluster of norovirus strains have been the most commonly detected noroviruses circulating worldwide since the mid-1990s, over time, distinct variants of the GII.4 virus evolve, emerge, and then recede to be replaced by a new variant³³. Emergence of a new GII.4 variant is associated with higher levels of infection and illness in the population and increased numbers of outbreaks³⁷, although severity of disease does not necessarily increase. These emergence events may be geographically contained (e.g. a 2003 variant emerged in Asia, and a 2006 variant spread in Europe)³⁸, or may be global, with new variants emerging and spreading worldwide over the course of a single year (as seen in 2002³⁹, 2006⁴⁰, 2009¹⁸ and 2012¹⁹).

Noroviruses have been shown to have high evolutionary rates, up to 10^{-2} substitutions/site/year in the VP1 protein⁴¹, due to the error-prone nature of the virus-encoded RdRp⁴². The rate of evolution is fastest in the P2 domain, which interacts with the host immune system. Immune response to norovirus infection appears to target this region of the virus capsid, and epitopes in this domain have been identified as important in defining the antigenic profile of GII.4 norovirus strains⁴³⁻⁴⁷, leading to the emergence of antigenically distinct viruses in the population, associated with epidemic/pandemic waves of gastroenteritis^{33, 37, 48, 49}.

The emergence of variant GII.4 strains is associated with mutations occurring in the virus at epitope A (VP1 amino acid positions 296-298, 368 & 372) and D (VP1 amino acid positions 393-395)^{44,50}. Mutations, particularly in epitopes A and D, will be selected for in the virus population if the mutations are such that existing immunity in the host population is evaded by the mutated virus, but the virus is otherwise not disadvantaged. Because much of the human population is exposed to antigenically similar noroviruses at a similar time, virus-specific immunity is likely similar between many people. As a result, the variant norovirus is advantaged, being more likely to successfully evade existing immunity, and subsequently establish more infections and be transmitted. In this way, the virus can spread quickly through the population. Eventually infections generate new immunological responses, which ultimately limit the success of this variant in the population, but in turn creates an ecological niche favourable for a new variant, and the process cycles again. This process has been observed for GII.4 noroviruses throughout the 1990s, 2000s and 2010s^{19, 33, 37, 45, 47}.

Norovirus surveillance

Surveillance of norovirus is complicated because most people do not contact medical services when they are ill. In the UK, it is suggested that for each laboratory report of norovirus around 300 cases go unreported⁵¹. This is largely related to the nature of the illness itself. The virus is highly infectious with an estimated infectious dose of around 10-100 virus particles (virions) needed to cause infection⁵², with a high probability of infection from ingesting a single particle⁵³. It has a short incubation period, anywhere between 12 and 72 hours, and symptoms typically last for around 24- 48 hours⁵⁴. Despite these difficulties it is still recognised as the commonest cause of gastrointestinal disease, not just in the UK, but worldwide⁵⁵. In the UK it is estimated between 3 and 4 million cases occur annually^{51, 56}, at a cost of £106 million to patients and the health care services. In the USA this estimate is around 21 million domestically acquired cases⁵⁷. Infections with norovirus occur in all age groups, however, the highest incidence is in children aged less than five years^{56, 58}.

The illness is often described as generally mild and self-limiting. The description of a mild infection can trivialise the effect of the illness; in England it has been estimated that 3000 admissions occur annually as a result of norovirus infection in adults⁵⁹ or 0.3% of emergency admissions in those aged over 65, and 0.1% in adults aged 16-64 years. The consequences of infection are also greater in vulnerable populations. In a study in the county of Avon, UK, hospital patients were ill for longer than care home residents and staff working in the hospitals or care home, with around 10% of inpatients affected still showing symptoms 7 days after becoming ill⁶⁰. There is also evidence that norovirus can contribute to mortality in the elderly. Modelling of deaths suggests that norovirus is associated with 20% in those aged over 65 years who died of infectious intestinal disease, and that 13% of deaths caused by non-infectious intestinal disease⁶¹.

Public Health England have conducted surveillance of gastrointestinal disease outbreaks since 1992⁶². Analysis of the first nine years of data highlighted the importance of norovirus outbreaks in hospitals; over 80% of all reported

outbreaks in hospitals were suspected or confirmed as norovirus, and 25% of all general outbreaks occurred in hospitals⁶³.

Recent developments

Surveillance

Since the recognition of the importance of norovirus as a cause of GI disease a more detailed online surveillance system was set up in 2009⁶³. The online system increased ascertainment of outbreaks in hospitals, with more outbreaks reported in the first year than the whole of the preceding system⁶³. Both systems highlighted the increased activity during the winter months, and the considerable burden it places on NHS hospitals in England. The online system suggests around 13000 patients and 3000 staff are affected each year, moreover, almost 9000 bed days are lost because of restrictions to admissions during outbreaks⁶³.

The key to surveillance of norovirus is allying the epidemiology with surveillance of virology. It is often difficult to achieve this. Recording the number of outbreaks, and laboratory reports indicates levels of infection, but they cannot directly ally this knowledge of circulating strains of the virus. The activity recorded in both Public Health England's hospital outbreak reporting scheme and laboratory reports suggests that seasonal activity varies considerably. The reasons for changes in seasonal activity need unpicking and modelling of changes in the circulating strains of norovirus against laboratory reporting provided evidence that modifications within the virus itself leads to changes in the epidemiology. In the autumn/winter of 2012 PHE recorded increased levels of norovirus activity; later attributed to the emergence of the Sydney 2012 strain³⁷. However, other reasons have been proposed, such as changes in winter conditions such as falling temperature⁶⁴.

Given the difficulty in surveillance of norovirus infections from direct sources, other developments need to be explored. For example, social media could provide early indications of increasing activity. There are a number of publications looking at the use of internet search and social media postings to provide information on increased disease activity⁶⁵⁻⁶⁷. Other forms of syndromic surveillance have been used such as the use of telephone helpline data to map diarrhoea and vomiting⁶⁸, difficulties with this approach fall mainly on disentangling the causes of the illness from syndromes (diarrhoea and vomiting). Norovirus is not the only cause of D&V and has a seasonality similar to that of rotavirus, similarly sapovirus has similar illness characteristics to norovirus.

Virus culture systems

Understanding the interactions of norovirus with host cells has been limited by the lack of an *in vitro* laboratory cell culture system. Attempts to establish conventional cell culture approaches were unsuccessful⁶⁹, after which alternative approaches were developed^{70,71}, however, these were limited in their usefulness.

More recently, progress has been made towards development of laboratory culture systems for human norovirus. Two systems have been described: one describes human norovirus replication in B cells⁸, and a second which describes human norovirus replication supported by stem cell-derived human enteroids⁷. These systems present exciting new opportunities to understand how norovirus interacts both with the host cell and with the host environment.

The system using human enteroids⁷ provides a model for processes of norovirus replication such as attachment/entry, genome replication, and virus assembly/release can be interrogated in a biologically relevant cell type. Advances in these areas will be crucial for identifying targets for virus-specific interventions, and evaluating how effective different antiviral therapies can limit norovirus replication. Further insights into virus entry and egress will enhance understanding of the interactions between virus and host receptors and identify novel interactions between virus and host that serve as intervention targets, for example antibodies which interfere with attachment or release processes, thus neutralising free virus.

The second system, in which norovirus replication is supported in B cells, uses commensal bacteria that express HBGAs to facilitate virus replication in this model^{8, 72}. Analysis of norovirus replication in this system could enhance understanding of the interaction between norovirus and HBGAs – and identify how these interactions might be disrupted– but also what interactions might occur with, and what role might be played by, the microbiome during norovirus infection⁷³.

Norovirus vaccines

Modelling studies have shown that norovirus vaccination would offer healthcare and economic benefit⁷⁴. These could help control and prevent the large-scale and often protracted outbreaks often seen in healthcare settings⁷⁵ and other settings such as in the military^{76, 77}.

Until very recently, development of candidate vaccines focussed on recombinant protein systems; expression of the norovirus capsid protein VP1 *in vitro* leads to self-assembly of the protein into virus-like particles (VLPs) that are antigenically and morphologically identical to infectious virus, but lacking a genome, VLPs are entirely non-infectious⁷⁸.

Early clinical studies of responses in humans to immunisation with VLPs demonstrated they were immunogenic when delivered orally^{79, 80} or intranasally⁸¹. A randomised, double-blind placebo-controlled trial conducted in healthy, susceptible adult volunteers investigated the safety and efficacy of vaccination using norovirus VLPs, followed by challenge with a homologous norovirus strain⁶. This trial demonstrated 70% of vaccine recipients had a virus-specific IgA response, and vaccination reduced the frequency of both infection and disease between placebo control group and vaccine recipients⁶.

However, the prototype vaccine (and challenge strain) used in this trial was based on a single norovirus strain – the prototype GI.1 Norwalk virus/1968 – which is uncommon, detected in <1% of norovirus strains characterised per year in surveillance programmes in developed countries. As the most significant disease burden is associated with the GII.4 genocluster, any candidate vaccine would need to elicit immunity to GII.4 norovirus strains, and cross-react to antigenically distinct GII.4 variants. A chimeric VLP was developed incorporating epitopes from antigenically distinct GII.4 viruses⁸², and induced broadly-reactive antibody responses⁸³.

A subsequent trial incorporated the chimeric VLP into a GI.1/GII.4 bivalent vaccine formulation and demonstrated vaccine induced seroconversion in 90% of vaccine recipients, and reduced gastroenteritis following challenge⁵. However, the predefined primary endpoints were not achieved in this study, and further studies are necessary to assess how effective this candidate vaccine would be in the general population, and specifically in paediatric and elderly populations. Furthermore, studies must address both the duration of and the inter-/intra-genotype breadth of protection.

Perspectives

Clearly, significant progress has been made in understanding the virology and epidemiology of norovirus in humans: but there remain significant gaps in our knowledge, important for development of therapeutic and preventative interventions, and ascertaining norovirus disease burden to understand how these should be utilised, and to measure their effectiveness. This is true in all economic settings, but especially in low economic settings.

One key question is to understand the emergence of norovirus strains. With no animal reservoir¹⁷, the virus must be sustained – and continuously evolve – in the human population. Using genomics approaches to measure and monitor virus diversity among circulating strains, and to characterise and measure whether observed genetic changes induce phenotypic changes will be crucial in developing the systems needed to understand and monitor emergence events, particularly those that lead to rapid pandemic spread of norovirus strains. There is increasing evidence that children may act as important reservoirs of norovirus, and the virus may exploit the more naïve immunological background in children to explore antigenic diversity, ultimately leading to virus diversification and subsequent emergence of novel strains⁸⁴.

Second, more detailed understanding of the burden of the disease, transmission dynamics and pathogenesis in risk groups, both those at risk of more severe disease (immunocompromised⁸⁵, elderly⁶¹), and those more likely to come into contact with or are at higher risk of transmitting the virus (food-handlers⁸⁶, healthcare workers⁸⁷, military personnel⁷⁶) is needed. There are complex epidemiological and virological questions relating to the distribution of norovirus-associated disease in the population, for whom the disease burden is greatest, as well as understanding host susceptibility factors. Integrated laboratory and epidemiological studies are crucial to investigate how norovirus is transmitted, disease attribution via different transmission pathways, how infections can be tracked in the

population and during outbreaks, and what role susceptibility factors such as HBGA phenotype or the individual microbiome composition may play in norovirus infection, development of disease and outcomes.

Third, alongside data on the direct burden of disease, enhanced data are needed to understand where interventions may alleviate transmission and disease overall, as many settings are interlinked. For example, administering a norovirus vaccine to patients in long term care homes might help prevent outbreaks in this environment, but might have limited effects on the population as a whole. However, it may be a worthwhile strategy if vaccination in care homes subsequently prevents outbreaks in hospitals and reduces bed blocking.

With the recent advances in laboratory culture systems for norovirus^{7, 88}, next generation sequencing technologies⁸⁹, improved diagnostics⁴ and measuring phenotypic characteristics of noroviruses⁹⁰, there are new opportunities to advance understanding of this common and important human pathogen.

Acknowledgments

The research was funded by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Gastrointestinal Infections at University of Liverpool in partnership with Public Health England (PHE), in collaboration with University of East Anglia, University of Oxford and the Institute of Food Research. DJA is based at the London School of Hygiene & Tropical Medicine. JPH is based at University of Liverpool. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, the Department of Health or Public Health England.

References

1. Kapikian AZ, Wyatt RG, Dolin R, Thornhill TS, Kalica AR, Chanock RM. Visualization by immune electron microscopy of a 27-nm particle associated with acute infectious nonbacterial gastroenteritis. *J Virol* 1972; 10:1075-81.
2. Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleeschauwer B, et al. World Health Organization Estimates of the Global and Regional Disease Burden of 22 Foodborne Bacterial, Protozoal, and Viral Diseases, 2010: A Data Synthesis. *PLoS Med* 2015; 12:e1001921.
3. Harris JP. Norovirus Surveillance: An Epidemiological Perspective. *J Infect Dis* 2016; 213 Suppl 1:S8-S11.
4. Kirby A, Iturriza-Gomara M. Norovirus diagnostics: options, applications and interpretations. *Expert Rev Anti Infect Ther* 2012; 10:423-33.
5. Bernstein DI, Atmar RL, Lyon GM, Treanor JJ, Chen WH, Jiang X, et al. Norovirus vaccine against experimental human GII.4 virus illness: a challenge study in healthy adults. *J Infect Dis* 2015; 211:870-8.
6. Atmar RL, Bernstein DI, Harro CD, Al-Ibrahim MS, Chen WH, Ferreira J, et al. Norovirus vaccine against experimental human Norwalk Virus illness. *N Engl J Med* 2011; 365:2178-87.
7. Ettayebi K, Crawford SE, Murakami K, Broughman JR, Karandikar U, Tenge VR, et al. Replication of human noroviruses in stem cell-derived human enteroids. *Science* 2016; 353:1387-93.
8. Jones MK, Watanabe M, Zhu S, Graves CL, Keyes LR, Grau KR, et al. Enteric bacteria promote human and mouse norovirus infection of B cells. *Science* 2014; 346:755-9.
9. Jiang X, Wang M, Wang K, Estes MK. Sequence and genomic organization of Norwalk virus. *Virology* 1993; 195:51-61.
10. Lambden PR, Liu B, Clarke IN. A conserved sequence motif at the 5' terminus of the Southampton virus genome is characteristic of the Caliciviridae. *Virus Genes* 1995; 10:149-52.
11. Mayo MA. Virus taxonomy - Houston 2002. *Arch Virol* 2002; 147:1071-6.
12. Zheng DP, Ando T, Fankhauser RL, Beard RS, Glass RI, Monroe SS. Norovirus classification and proposed strain nomenclature. *Virology* 2006; 346:312-23.
13. Kroneman A, Vega E, Vennema H, Vinje J, White PA, Hansman G, et al. Proposal for a unified norovirus nomenclature and genotyping. *Arch Virol* 2013; 158:2059-68.
14. Green K. Caliciviridae: the noroviruses. In: Knipe DM HP, Cohen JI, Griffin DE, Lamb RA, Martin MA, Racaniello VR, Roizman B, ed. *Fields virology*, 6th ed: Lippincott, Williams & Wilkins, Philadelphia, PA, 2013:583-609.
15. Tse H, Lau SK, Chan WM, Choi GK, Woo PC, Yuen KY. Complete genome sequences of novel canine noroviruses in Hong Kong. *J Virol* 2012; 86:9531-2.
16. Vinje J. Advances in laboratory methods for detection and typing of norovirus. *J Clin Microbiol* 2015; 53:373-81.
17. Wilhelm B, Waddell L, Greig J, Rajic A, Houde A, McEwen SA. A scoping review of the evidence for public health risks of three emerging potentially zoonotic viruses: hepatitis E virus, norovirus, and rotavirus. *Prev Vet Med* 2015; 119:61-79.
18. Vega E, Barclay L, Gregoricus N, Williams K, Lee D, Vinje J. Novel surveillance network for norovirus gastroenteritis outbreaks, United States. *Emerg Infect Dis* 2011; 17:1389-95.
19. van Beek J, Ambert-Balay K, Botteldoorn N, Eden JS, Fonager J, Hewitt J, et al. Indications for worldwide increased norovirus activity associated with emergence of a new variant of genotype II.4, late 2012. *Euro Surveill* 2013; 18:8-9.
20. Clarke IN, Lambden PR. Organization and expression of calicivirus genes. *J Infect Dis* 2000; 181 Suppl 2:S309-16.
21. Prasad BV, Hardy ME, Dokland T, Bella J, Rossmann MG, Estes MK. X-ray crystallographic structure of the Norwalk virus capsid. *Science* 1999; 286:287-90.
22. Huang P, Farkas T, Marionneau S, Zhong W, Ruvoen-Clouet N, Morrow AL, et al. Noroviruses bind to human ABO, Lewis, and secretor histo-blood group antigens: identification of 4 distinct strain-specific patterns. *J Infect Dis* 2003; 188:19-31.
23. Tan M, Jiang X. Norovirus and its histo-blood group antigen receptors: an answer to a historical puzzle. *Trends Microbiol* 2005; 13:285-93.

24. Shanker S, Choi JM, Sankaran B, Atmar RL, Estes MK, Prasad BV. Structural analysis of histo-blood group antigen binding specificity in a norovirus GII.4 epidemic variant: implications for epochal evolution. *J Virol* 2011; 85:8635-45.
25. Bu W, Mamedova A, Tan M, Xia M, Jiang X, Hegde RS. Structural basis for the receptor binding specificity of Norwalk virus. *J Virol* 2008; 82:5340-7.
26. Chen Y, Tan M, Xia M, Hao N, Zhang XC, Huang P, et al. Crystallography of a Lewis-binding norovirus, elucidation of strain-specificity to the polymorphic human histo-blood group antigens. *PLoS Pathog* 2011; 7:e1002152.
27. Choi JM, Hutson AM, Estes MK, Prasad BV. Atomic resolution structural characterization of recognition of histo-blood group antigens by Norwalk virus. *Proc Natl Acad Sci U S A* 2008; 105:9175-80.
28. Hansman GS, Biertumpfel C, Georgiev I, McLellan JS, Chen L, Zhou T, et al. Crystal structures of GII.10 and GII.12 norovirus protruding domains in complex with histo-blood group antigens reveal details for a potential site of vulnerability. *J Virol* 2011; 85:6687-701.
29. Cao S, Lou Z, Tan M, Chen Y, Liu Y, Zhang Z, et al. Structural basis for the recognition of blood group trisaccharides by norovirus. *J Virol* 2007; 81:5949-57.
30. Currier RL, Payne DC, Staat MA, Selvarangan R, Shirley SH, Halasa N, et al. Innate Susceptibility to Norovirus Infections Influenced by FUT2 Genotype in a United States Pediatric Population. *Clin Infect Dis* 2015; 60:1631-8.
31. Lindesmith L, Moe C, Marionneau S, Ruvoen N, Jiang X, Lindblad L, et al. Human susceptibility and resistance to Norwalk virus infection. *Nat Med* 2003; 9:548-53.
32. Bustamante M, Standl M, Bassat Q, Vilor-Tejedor N, Medina-Gomez C, Bonilla C, et al. A genome-wide association meta-analysis of diarrhoeal disease in young children identifies FUT2 locus and provides plausible biological pathways. *Hum Mol Genet* 2016; 25:4127-42.
33. Zakikhany K, Allen DJ, Brown D, Iturriza-Gomara M. Molecular evolution of GII-4 Norovirus strains. *PLoS One* 2012; 7:e41625.
34. Kroneman A, Harris J, Vennema H, Duizer E, van Duynhoven Y, Gray J, et al. Data quality of 5 years of central norovirus outbreak reporting in the European Network for food-borne viruses. *J Public Health (Oxf)* 2008; 30:82-90.
35. Vega E, Barclay L, Gregoricus N, Shirley SH, Lee D, Vinje J. Genotypic and epidemiologic trends of norovirus outbreaks in the United States, 2009 to 2013. *J Clin Microbiol* 2014; 52:147-55.
36. Mans J, Murray TY, Nadan S, Netshikweta R, Page NA, Taylor MB. Norovirus diversity in children with gastroenteritis in South Africa from 2009 to 2013: GII.4 variants and recombinant strains predominate. *Epidemiol Infect* 2016; 144:907-16.
37. Allen DJ, Adams NL, Aladin F, Harris JP, Brown DW. Emergence of the GII-4 Norovirus Sydney2012 strain in England, winter 2012-2013. *PLoS One* 2014; 9:e88978.
38. Siebenga JJ, Vennema H, Zheng DP, Vinje J, Lee BE, Pang XL, et al. Norovirus illness is a global problem: emergence and spread of norovirus GII.4 variants, 2001-2007. *J Infect Dis* 2009; 200:802-12.
39. Lopman B, Vennema H, Kohli E, Pothier P, Sanchez A, Negredo A, et al. Increase in viral gastroenteritis outbreaks in Europe and epidemic spread of new norovirus variant. *Lancet* 2004; 363:682-8.
40. Tu ET, Bull RA, Greening GE, Hewitt J, Lyon MJ, Marshall JA, et al. Epidemics of gastroenteritis during 2006 were associated with the spread of norovirus GII.4 variants 2006a and 2006b. *Clin Infect Dis* 2008; 46:413-20.
41. Victoria M, Miagostovich MP, Ferreira MS, Vieira CB, Fioretti JM, Leite JP, et al. Bayesian coalescent inference reveals high evolutionary rates and expansion of Norovirus populations. *Infect Genet Evol* 2009; 9:927-32.
42. Bull RA, Eden JS, Rawlinson WD, White PA. Rapid evolution of pandemic noroviruses of the GII.4 lineage. *PLoS Pathog* 2010; 6:e1000831.
43. Allen DJ, Gray JJ, Gallimore CI, Xerry J, Iturriza-Gomara M. Analysis of amino acid variation in the P2 domain of the GII-4 norovirus VP1 protein reveals putative variant-specific epitopes. *PLoS One* 2008; 3:e1485.
44. Allen DJ, Noad R, Samuel D, Gray JJ, Roy P, Iturriza-Gomara M. Characterisation of a GII-4 norovirus variant-specific surface-exposed site involved in antibody binding. *Virol J* 2009; 6:150.
45. Lindesmith LC, Donaldson EF, Lobue AD, Cannon JL, Zheng DP, Vinje J, et al. Mechanisms of GII.4 norovirus persistence in human populations. *PLoS Med* 2008; 5:e31.
46. Lindesmith LC, Donaldson EF, Baric RS. Norovirus GII.4 strain antigenic variation. *J Virol* 2011; 85:231-42.
47. Siebenga JJ, Vennema H, Renckens B, de Bruin E, van der Veer B, Siezen RJ, et al. Epochal evolution of GGII.4 norovirus capsid proteins from 1995 to 2006. *J Virol* 2007; 81:9932-41.

48. Debbink K, Lindesmith LC, Donaldson EF, Costantini V, Beltramello M, Corti D, et al. Emergence of new pandemic GII.4 Sydney norovirus strain correlates with escape from herd immunity. *J Infect Dis* 2013; 208:1877-87.
49. Debbink K, Lindesmith LC, Donaldson EF, Baric RS. Norovirus immunity and the great escape. *PLoS Pathog* 2012; 8:e1002921.
50. Lindesmith LC, Debbink K, Swanstrom J, Vinje J, Costantini V, Baric RS, et al. Monoclonal antibody-based antigenic mapping of norovirus GII.4-2002. *J Virol* 2012; 86:873-83.
51. Tam CC, Rodrigues LC, Viviani L, Dodds JP, Evans MR, Hunter PR, et al. Longitudinal study of infectious intestinal disease in the UK (IID2 study): incidence in the community and presenting to general practice. *Gut* 2012; 61:69-77.
52. Glass RI, Noel J, Ando T, Fankhauser R, Belliot G, Mounts A, et al. The epidemiology of enteric caliciviruses from humans: a reassessment using new diagnostics. *J Infect Dis* 2000; 181 Suppl 2:S254-61.
53. Teunis PF, Moe CL, Liu P, Miller SE, Lindesmith L, Baric RS, et al. Norwalk virus: how infectious is it? *J Med Virol* 2008; 80:1468-76.
54. Estes MK, Prasad BV, Atmar RL. Noroviruses everywhere: has something changed? *Curr Opin Infect Dis* 2006; 19:467-74.
55. Belliot G, Lopman BA, Ambert-Balay K, Pothier P. The burden of norovirus gastroenteritis: an important foodborne and healthcare-related infection. *Clin Microbiol Infect* 2014; 20:724-30.
56. Harris JP, Iturriza-Gomara M, O'Brien SJ. Re-assessing the total burden of norovirus circulating in the United Kingdom population. *Vaccine* 2017; 35:853-5.
57. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, et al. Foodborne illness acquired in the United States--major pathogens. *Emerg Infect Dis* 2011; 17:7-15.
58. O'Brien SJ, Donaldson AL, Iturriza-Gomara M, Tam CC. Age-Specific Incidence Rates for Norovirus in the Community and Presenting to Primary Healthcare Facilities in the United Kingdom. *J Infect Dis* 2016; 213 Suppl 1:S15-8.
59. Haustein T, Harris JP, Pebody R, Lopman BA. Hospital admissions due to norovirus in adult and elderly patients in England. *Clin Infect Dis* 2009; 49:1890-2.
60. Lopman BA, Andrews N, Sarangi J, Vipond IB, Brown DW, Reacher MH. Institutional risk factors for outbreaks of nosocomial gastroenteritis: survival analysis of a cohort of hospital units in South-west England, 2002-2003. *J Hosp Infect* 2005; 60:135-43.
61. Harris JP, Edmunds WJ, Pebody R, Brown DW, Lopman BA. Deaths from norovirus among the elderly, England and Wales. *Emerg Infect Dis* 2008; 14:1546-52.
62. Wall PG, de Louvois J, Gilbert RJ, Rowe B. Food poisoning: notifications, laboratory reports, and outbreaks--where do the statistics come from and what do they mean? *Commun Dis Rep CDR Rev* 1996; 6:R93-100.
63. Harris JP, Adams NL, Lopman BA, Allen DJ, Adak GK. The development of Web-based surveillance provides new insights into the burden of norovirus outbreaks in hospitals in England. *Epidemiol Infect* 2014; 142:1590-8.
64. Lopman B, Armstrong B, Atchison C, Gray JJ. Host, weather and virological factors drive norovirus epidemiology: time-series analysis of laboratory surveillance data in England and Wales. *PLoS One* 2009; 4:e6671.
65. Choi J, Cho Y, Shim E, Woo H. Web-based infectious disease surveillance systems and public health perspectives: a systematic review. *BMC Public Health* 2016; 16:1238.
66. Harris JK, Wong R, Matthew MG, Mansour R. Availability of Foodborne Illness Reporting Mechanisms for the Public on Local Health Department Web Sites. *J Public Health Manag Pract* 2017.
67. Harris JK, Hawkins JB, Nguyen L, Nsoesie EO, Tuli G, Mansour R, et al. Using Twitter to Identify and Respond to Food Poisoning: The Food Safety STL Project. *J Public Health Manag Pract* 2017.
68. Loveridge P, Cooper D, Elliot AJ, Harris J, Gray J, Large S, et al. Vomiting calls to NHS Direct provide an early warning of norovirus outbreaks in hospitals. *J Hosp Infect* 2010; 74:385-93.
69. Duizer E, Schwab KJ, Neill FH, Atmar RL, Koopmans MP, Estes MK. Laboratory efforts to cultivate noroviruses. *J Gen Virol* 2004; 85:79-87.
70. Katayama K, Murakami K, Sharp TM, Guix S, Oka T, Takai-Todaka R, et al. Plasmid-based human norovirus reverse genetics system produces reporter-tagged progeny virus containing infectious genomic RNA. *Proc Natl Acad Sci U S A* 2014; 111:E4043-52.

71. Chang KO, Sosnovtsev SV, Belliot G, King AD, Green KY. Stable expression of a Norwalk virus RNA replicon in a human hepatoma cell line. *Virology* 2006; 353:463-73.
72. Karst SM. Identification of a novel cellular target and a co-factor for norovirus infection - B cells & commensal bacteria. *Gut Microbes* 2015; 6:266-71.
73. Baldrige MT, Nice TJ, McCune BT, Yokoyama CC, Kambal A, Wheadon M, et al. Commensal microbes and interferon-lambda determine persistence of enteric murine norovirus infection. *Science* 2015; 347:266-9.
74. Bartsch SM, Lopman BA, Hall AJ, Parashar UD, Lee BY. The potential economic value of a human norovirus vaccine for the United States. *Vaccine* 2012; 30:7097-104.
75. Morter S, Bennet G, Fish J, Richards J, Allen DJ, Nawaz S, et al. Norovirus in the hospital setting: virus introduction and spread within the hospital environment. *J Hosp Infect* 2011; 77:106-12.
76. McCarthy M, Estes MK, Hyams KC. Norwalk-like virus infection in military forces: epidemic potential, sporadic disease, and the future direction of prevention and control efforts. *J Infect Dis* 2000; 181 Suppl 2:S387-91.
77. Centers for Disease C, Prevention. Outbreak of acute gastroenteritis associated with Norwalk-like viruses among British military personnel--Afghanistan, May 2002. *MMWR Morb Mortal Wkly Rep* 2002; 51:477-9.
78. Jiang X, Wang M, Graham DY, Estes MK. Expression, self-assembly, and antigenicity of the Norwalk virus capsid protein. *J Virol* 1992; 66:6527-32.
79. Ball JM, Graham DY, Opekun AR, Gilger MA, Guerrero RA, Estes MK. Recombinant Norwalk virus-like particles given orally to volunteers: phase I study. *Gastroenterology* 1999; 117:40-8.
80. Tacket CO, Sztein MB, Losonsky GA, Wasserman SS, Estes MK. Humoral, mucosal, and cellular immune responses to oral Norwalk virus-like particles in volunteers. *Clin Immunol* 2003; 108:241-7.
81. El-Kamary SS, Pasetti MF, Mendelman PM, Frey SE, Bernstein DI, Treanor JJ, et al. Adjuvanted intranasal Norwalk virus-like particle vaccine elicits antibodies and antibody-secreting cells that express homing receptors for mucosal and peripheral lymphoid tissues. *J Infect Dis* 2010; 202:1649-58.
82. Parra GI, Bok K, Taylor R, Haynes JR, Sosnovtsev SV, Richardson C, et al. Immunogenicity and specificity of norovirus Consensus GII.4 virus-like particles in monovalent and bivalent vaccine formulations. *Vaccine* 2012; 30:3580-6.
83. Debbink K, Lindesmith LC, Donaldson EF, Swanstrom J, Baric RS. Chimeric GII.4 norovirus virus-like-particle-based vaccines induce broadly blocking immune responses. *J Virol* 2014; 88:7256-66.
84. Allen DJ, Trainor E, Callaghan A, O'Brien SJ, Cunliffe NA, Iturriza-Gomara M. Early Detection of Epidemic GII-4 Norovirus Strains in UK and Malawi: Role of Surveillance of Sporadic Acute Gastroenteritis in Anticipating Global Epidemics. *PLoS One* 2016; 11:e0146972.
85. Bok K, Green KY. Norovirus gastroenteritis in immunocompromised patients. *N Engl J Med* 2012; 367:2126-32.
86. Rumble C, Addiman S, Balasegaram S, Chima K, Ready D, Heard J, et al. Role of Food Handlers in Norovirus Outbreaks in London and South East England, 2013 to 2015. *J Food Prot* 2017; 80:257-64.
87. Sabria A, Pinto RM, Bosch A, Bartolome R, Cornejo T, Torner N, et al. Norovirus shedding among food and healthcare workers exposed to the virus in outbreak settings. *J Clin Virol* 2016; 82:119-25.
88. Jones MK, Grau KR, Costantini V, Kolawole AO, de Graaf M, Freiden P, et al. Human norovirus culture in B cells. *Nat Protoc* 2015; 10:1939-47.
89. Brown JR, Roy S, Ruis C, Yara Romero E, Shah D, Williams R, et al. Norovirus Whole-Genome Sequencing by SureSelect Target Enrichment: a Robust and Sensitive Method. *J Clin Microbiol* 2016; 54:2530-7.
90. Debbink K, Donaldson EF, Lindesmith LC, Baric RS. Genetic mapping of a highly variable norovirus GII.4 blockade epitope: potential role in escape from human herd immunity. *J Virol* 2012; 86:1214-26.