

Supplemental Table. 2-Year Outcomes of patients discharged on dual anti-platelet therapy after percutaneous coronary intervention compared across baseline white blood cell count (WBC) quartiles

	WBC quartile 1 (2.10-6.60 x 10 ³ cells/uL) (n=1065)	WBC quartile 2 (6.61-8.00 x 10 ³ cells/uL) (n=1065)	WBC quartile 3 (8.01-9.83 x 10 ³ cells/uL) (n=1037)	WBC quartile 4 (9.84-29.02 x 10 ³ cells/uL) (n=1055)	p _{trend}
Major adverse cardiovascular event (composite of cardiac death, definite or probable stent thrombosis, spontaneous myocardial infarction, or target lesion revascularization) (%)	9.9 [8.2-11.9]	10.2 [8.5-12.2]	13.0 [11.1-15.3]	14.0 [12.0-16.3]	0.0004
Cardiac death (%)	1.9 [1.2-2.9]	3.3 [2.3-4.6]	3.9 [2.9-5.3]	4.0 [2.9-5.4]	0.005
Definite/probable stent thrombosis (%)	1.1 [0.6-2.0]	0.9 [0.5-1.7]	2.1 [1.4-3.2]	2.0 [1.3-3.0]	0.02
Spontaneous myocardial infarction (%)	3.3 [2.3-4.6]	3.4 [2.4-4.7]	3.9 [2.8-5.3]	4.5 [3.4-6.0]	0.11
Clinically indicated target lesion revascularization (%)	6.6 [5.2-8.3]	6.1 [4.8-7.8]	8.4 [6.8-10.3]	9.0 [7.3-10.9]	0.007

Data are shown as crude estimates [95% confidence intervals]

