

1 **Medical intensive care unit admission among patients with and without HIV,**
2 **hepatitis C virus, and alcohol-related diagnoses in the United States: a national,**
3 **retrospective cohort study, 1997-2014**

4 **Authors:** Christopher T. Rentsch, MPH^{1,2,3}; Janet P. Tate, ScD^{2,3}; Tessa Steel, MD⁴; Adeel A.
5 Butt, MD^{5,6}; Cynthia L. Gibert, MD^{7,8}; Laurence Huang, MD⁹; Margaret Pisani, MD³; Guy W. Soo
6 Hoo, MD^{10,11}; Stephen Crystal, PhD¹²; Maria C. Rodriguez-Barradas, MD^{13,14}; Sheldon T. Brown,
7 MD^{15,16}; Matthew S. Freiberg, MD^{17,18}; Christopher J. Graber, MD^{10,11}; Joon W. Kim, MD¹⁹; David
8 Rimland, MD²⁰; Amy C. Justice, MD^{2,3,21}; David A. Fiellin, MD^{3,21,22}; Kristina A. Crothers, MD⁴;
9 Kathleen M. Akgün, MD^{3,23}

10 ¹Epidemiology & Population Health, London School of Hygiene & Tropical Medicine, London,
11 UK

12 ²Veterans Aging Cohort Study Coordinating Center, VA Connecticut Healthcare System, West
13 Haven, CT, USA

14 ³Internal Medicine, Yale School of Medicine, New Haven, CT, USA

15 ⁴Internal Medicine, University of Washington School of Medicine, Seattle, WA, USA

16 ⁵Infectious Diseases, VA Pittsburgh Healthcare System, Pittsburgh, PA, USA

17 ⁶Weill Cornell Medical College, New York City, NY, USA and Doha, Qatar

18 ⁷Infectious Diseases, VA Medical Center, Washington, DC, USA

19 ⁸Medicine and Health Sciences, The George Washington University, Washington, DC, USA

20 ⁹Medicine, University of California, San Francisco, CA, USA

21 ¹⁰Medicine, VA Greater Los Angeles Healthcare System, Los Angeles, CA, USA

22 ¹¹David Geffen School of Medicine, University of California, Los Angeles, CA, USA

23 ¹²Center for Health Services Research, Rutgers University, New Brunswick, NJ, USA

24 ¹³Medicine, Michael E. DeBakey VA Medical Center, Houston, TX, USA

25 ¹⁴Medicine-Infectious Diseases, Baylor College of Medicine, Houston, TX, USA

26 ¹⁵Medicine, James J. Peters VA Medical Center, New York, NY, USA

27 ¹⁶Icahn School of Medicine, Mount Sinai, New York, NY, USA

28 ¹⁷Medicine, Vanderbilt University School of Medicine, Nashville, TN, USA

29 ¹⁸Geriatric Research Education and Clinical Centers, VA Tennessee Valley Healthcare System,
30 Nashville, TN, USA

31 ¹⁹Critical Care Medicine, James J. Peters VA Medical Center, Bronx, NY, USA

32 ²⁰Medicine, Emory University School of Medicine, Atlanta, GA, USA

33 ²¹Center for Interdisciplinary Research on AIDS, Yale School of Public Health, New Haven, CT,
34 USA

35 ²²Yale School of Public Health, New Haven, CT, USA

36 ²³Internal Medicine, VA Connecticut Healthcare System, West Haven, CT, USA

37
38 **Corresponding author:**

39 Christopher T. Rentsch, MPH

40 VA Connecticut Healthcare System

41 Yale University School of Medicine

42 London School of Hygiene & Tropical Medicine

43 Keppel Street

44 WC1E 7HT

45 London, UK

46 US Phone: +1(614) 388-8786

47 UK Phone: +44(0)79 8345 0440

48 Email: Christopher.Rentsch@va.gov

1 **Reprints:** Reprints will not be ordered.
2

3 **Preliminary stages of this research were presented as poster presentations at:**

- 4 1. International Workshop on HIV Observational Databases, Lisbon, Portugal, March 30-
5 April 1, 2017
- 6 2. American Thoracic Society International Conference, Washington, DC, May 19-24, 2017
7

8 **Conflicts of Interest and Source of Funding:** The authors declare no conflict of interest. This
9 work was supported by the US National Institutes of Health (NIH), including grants from the
10 National Institute on Alcohol Abuse and Alcoholism [U24-AA020794, U01-AA020790, U10-
11 AA013566-completed to ACJ] and National Heart, Lung, and Blood Institute [R01-HL090342 to
12 KC, K24-HL087713 to LH].
13

14 **Running title:** MICU rates by HIV, HCV, and alcohol use
15

16 **Article type:** Original Article

17 **Category:** Clinical Science

18 **Word count:** 3015

19 **Abstract:** 244

20 **Tables:** 3 + 3 supplemental

21 **Figures:** 1

22 **References:** 26
23
24

1 **ABSTRACT**

2 **Background:** HIV, hepatitis C virus (HCV), and alcohol-related diagnoses (ARD) independently
3 contribute increased risk for all-cause hospitalization. We sought to determine annual medical
4 intensive care unit (MICU) admission rates and relative risk of MICU admission between 1997
5 and 2014 among people with and without HIV, HCV, and ARD, using data from the largest HIV
6 and HCV care provider in the United States.

7 **Setting:** Veterans Health Administration.

8 **Methods:** Annual MICU admission rates were calculated among 155,550 patients in the
9 Veterans Aging Cohort Study by HIV, HCV, and ARD status. Adjusted rate ratios (RR) and 95%
10 confidence intervals (CI) were estimated with Poisson regression. Significance of trends in age-
11 adjusted admission rates were tested with generalized linear regression. Models were stratified
12 by calendar period to identify shifts in MICU admission risk over time.

13 **Results:** Compared to HIV-/HCV-/ARD- patients, relative risk for MICU admission decreased
14 among HIV mono-infected patients from 61% (95% CI 1.56-1.65) in 1997-2009 to 21% (95% CI
15 1.16-1.27) in 2010-2014, increased among HCV mono-infected patients from 22% (95% CI
16 1.16-1.29) in 1997-2009 to 54% (95% CI 1.43-1.67) in 2010-2014, and remained consistent
17 among patients with ARD only at 46% (95% CI 1.42-1.50). MICU admission rates decreased by
18 48% among HCV-uninfected patients (p -trend<0.0001) but did not change among HCV+
19 patients (p -trend=0.34).

20 **Conclusion:** HCV infection and ARD remain key contributors to MICU admission risk. The
21 impact of each of these conditions could be mitigated with combination of treatment of HIV,
22 HCV, and interventions targeting unhealthy alcohol use.

1 **Key Words:** Intensive care units; HIV; Hepatitis C; Alcoholism; Electronic health records;

2 Veterans

3

1 **Introduction**

2 People with HIV infection are living longer on combination antiretroviral therapy (ART),
3 but continue to experience excess risk for multiple chronic conditions compared to
4 demographically similar HIV-uninfected individuals, including pulmonary,¹ cardiovascular,² and
5 liver disease.³ Further, these chronic medical conditions associated with aging are major
6 contributors to risk of all-cause hospitalization⁴ and mortality⁵ and may progress more rapidly in
7 HIV-infected (HIV+) patients. Since ART became widely available, hospitalization rates among
8 HIV+ patients have decreased⁴ while rates of medical intensive care unit (MICU) admissions
9 have remained stable or even increased.⁶

10

11 A growing percentage of MICU admissions among HIV+ patients are due to
12 complications arising from hepatitis C virus (HCV) co-infection.⁷ HCV infection is more common
13 in patients with HIV than their HIV-uninfected counterparts.⁸ Compared with HIV mono-infected
14 patients, HCV co-infection is associated with approximately 50% increased risk of
15 hospitalization, most often attributed to infection-related diagnoses.⁹ Although safe and effective
16 treatments for HCV infection became widely available in the United States in 2014¹⁰, the risk of
17 MICU admission conferred by HCV infection before 2014 remains largely unknown.

18

19 While effective treatments for HIV and HCV infection have led to improved health
20 outcomes, common and potentially modifiable health behaviors, such as alcohol consumption,
21 represent the next challenge.¹¹ A previous study showed that one-third of MICU admissions
22 were associated with heavy alcohol consumption.¹² Alcohol-related diagnoses (ARD), including
23 alcohol use disorder and other conditions caused by alcohol (e.g., alcoholic liver disease,
24 alcoholic gastritis), are commonly associated with conditions that lead to MICU admission
25 among HCV+ patients.¹³ ARD are also common among HIV+ patients¹⁴, and sequelae such as

1 physiologic injury¹⁵ and mortality¹⁶ occur at lower levels of alcohol exposure than among
2 uninfected patients. Importantly, even low levels of alcohol use can aggravate deleterious
3 effects of HIV and HCV infection on the liver.¹⁷

4
5 As HIV has become a chronic medical condition, the growing relevance of HCV infection
6 and ARD on MICU admission requires quantification as a first step toward identifying new
7 targets for medical intervention. We used data from an observational cohort in the Veterans
8 Health Administration (VA), the largest HIV and HCV care provider in the United States, and
9 Medicare, the largest healthcare insurer in the United States, to calculate annual MICU
10 admission rates and relative risk for MICU admission between 1997 and 2014 among people
11 living with and without HIV, HCV, and ARD.

12
13 **Methods**

14 *Study design and cohort*

15 We conducted a retrospective cohort study among patients in the Veterans Aging Cohort
16 Study (VACS), which has been previously described.¹⁸ Briefly, VACS is a large observational
17 cohort based on data from the national VA electronic health record system that includes all
18 HIV+ patients in VA care (over 50,000 subjects) matched 1:2 with HIV-uninfected patients
19 based on age, sex, race/ethnicity, and site of care. For this analysis, we calculated annual MICU
20 admission rates among all VACS patients who were active in VA care (i.e., had at least one
21 outpatient or inpatient visit) in each calendar year between 1997 and 2014. Baseline date was
22 defined as the date of VACS enrollment, which is the first HIV diagnosis date within the VA for
23 HIV+ patients or date of matching for HIV-uninfected controls. Patients were followed until their
24 last VA visit date or death. HIV-uninfected controls (n=338) who seroconverted during follow-up
25 were excluded. The development of VACS has been approved by the Institutional Review

1 Boards of the VA Connecticut Healthcare System and Yale University School of Medicine, was
2 granted a waiver of informed consent, and deemed HIPAA compliant.

3

4 *MICU admission and principal diagnoses*

5 MICU admission data were extracted from national VA and Medicare databases. VA
6 MICU admission was identified by inpatient bed section codes 12 or 13. Bed section code 12
7 included cardiac ICU (CICU) admissions until 2008 when CICU was given a separate bed
8 section code, 13. For the purposes of this analysis, MICU refers to both MICU and CICU
9 admissions. Medicare MICU admissions were identified from MEDPAR data using MICU and
10 CICU indicator variables. Admissions that resulted from a transfer from another unit were
11 included in this analysis. For each MICU admission we obtained principal diagnosis for the
12 hospitalization, according to International Classification of Diseases, Ninth Revision (ICD-9)
13 code and categorized them as in previous studies^{4,7} (see Table, Supplemental Digital Content
14 1).

15

16 *Exposure groups*

17 Eight exposure groups were created using all combinations of HIV, HCV, and ARD
18 status. HIV status was determined as previously described in VACS.¹⁸ HCV infection was
19 defined by positive HCV RNA during the study period. Time-updated ARD included validated
20 ICD-9 codes for alcohol intoxication/ingestion, alcohol-induced mental disorders, alcohol
21 dependence syndrome, alcoholic polyneuropathy/cardiomyopathy/gastritis, and alcohol-related
22 liver diseases (see Table, Supplemental Digital Content 1).¹⁹ Patients were classified as ARD+ if
23 in the previous year they had at least two outpatient encounters with an ARD; they remained
24 ARD+ thereafter.

25

1 *Covariates*

2 We extracted data on demographics including age, sex, and race/ethnicity. Age was
3 categorized into 5-year age groups ensuring adequate numbers in each age and exposure
4 group category when calculating age-adjusted, annual MICU admission rates. Common
5 comorbid conditions were extracted using ICD-9 codes for psychiatric disorders, hypertension,
6 drug use-related diagnoses, anemia, cardiovascular disease (CVD), diabetes, chronic
7 obstructive pulmonary disorder (COPD), renal insufficiency, non-AIDS cancers, chronic hepatitis
8 B infection, and liver diseases. Comorbidities were defined by one inpatient or two outpatient
9 diagnostic codes (see Table, Supplemental Digital Content 1) and considered present at
10 baseline if the diagnosis date occurred within one year before or six months after baseline date.
11 We also extracted laboratory data, including CD4 cell count and FIB-4 (a composite measure of
12 liver injury incorporating values for age, platelet count, AST, and ALT), and among HIV+
13 patients, pharmacy data to identify ART initiation.

14

15 *Statistical analyses*

16 We compared baseline demographic and clinical characteristics across the eight
17 exposure groups using non-parametric Kruskal-Wallis tests for continuous measures and chi-
18 square (χ^2) tests for categorical measures. We calculated annual MICU admission rates by
19 dividing the total number of MICU admissions by the number of patients at risk in a given
20 calendar year (i.e., having at least one inpatient or outpatient visit). Patients with multiple MICU
21 admissions could only contribute one admission per calendar year of follow-up. We estimated
22 age-adjusted MICU admission rates, as well as rate ratios (RR) with 95% confidence intervals
23 (95% CI) for each HIV/ARD/HCV exposure group adjusted for baseline age, comorbidity (i.e.,
24 psychiatric disorders, hypertension, drug use-related diagnoses, diabetes, cardiovascular
25 disease, anemia, chronic obstructive pulmonary disorder, non-AIDS cancers, renal insufficiency,

1 chronic hepatitis B, and liver diseases), and calendar year using Poisson regression. Covariates
2 were selected based on *a priori* hypotheses informed by clinical relevance and because they
3 represent potential confounders in the relationship between HIV, HCV, or ARD and risk of MICU
4 admission, as well as availability of data. We tested for trends in annual MICU admission rates
5 using generalized linear regression models. We estimated the individual and combined
6 associations of HIV, HCV, and ARD with risk of MICU admission in three ways: 1) comparing
7 HIV/HCV/ARD groups to the HIV-/HCV-/ARD- group, 2) comparing HIV/ARD groups stratified
8 by HCV infection, and 3) comparing HCV/ARD groups stratified by HIV infection. We also
9 stratified the study period into two eras to assess differential impacts of each exposure between
10 1997-2009 and 2010-2014. We selected these specific time periods to highlight findings from
11 the most recent five years of follow-up that would be most relevant to current patients. Statistical
12 analyses were performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA). Statistical
13 significance was defined as $p < 0.05$.

14

15 **Results**

16 *Cohort characteristics*

17 Between 1997 and 2014, 155,550 patients met study inclusion criteria (contributing
18 1,525,267 person-years); 32% were HIV+, 13% were HCV+, and 30% had an ARD during
19 follow-up. Median baseline age was 47 years (interquartile range [IQR] 40-54 years) (Table 1).
20 Most patients were male (97%), 47% were black, and 40% were white. While 42% of the
21 sample had no baseline comorbidity, 29% had only one and the remaining 28% had two or
22 more. The most common comorbidities at baseline were psychiatric disorders (27%),
23 hypertension (27%), drug use-related diagnoses (14%), diabetes (12%), and CVD (6%).

24

1 Each of the eight exposure groups based on HIV, HCV, and ARD had $\geq 3,500$ patients,
2 the largest being the group with none of the exposures (n=69,511). ARDs were more common
3 in HCV+ patients (58%) and HIV+ patients (30%) than patients with neither HCV nor HIV (27%)
4 ($p < 0.0001$). Overall mortality was 2.9 deaths per 100 person-years of observation. Patients with
5 HIV infection (irrespective of HCV infection and ARD status) had higher mortality rates than
6 those without HIV infection (4.3 per 100 person-years vs. 1.9 per 100 person-years, $p < 0.0001$)
7 (see Table, Supplemental Digital Content 2).

8
9 Among HIV+ patients, 57% initiated ART within one year of VACS enrollment (i.e.,
10 diagnosis of HIV within the VA), and an additional 21% had initiated ART during the study
11 period. In 1997, 35% of HIV+ patients had CD4 < 200 cells/mm³ and 20% had CD4 ≥ 500
12 cells/mm³. At the end of the study period, only 8% of HIV+ patients had CD4 < 200 cells/mm³
13 and 58% had CD4 ≥ 500 cells/mm³. Among HCV+ patients, FIB-4 scores increased over the
14 study period. At study start, 36% had FIB-4 scores between 1.45-3.25 and 16% had scores
15 > 3.25 . By 2014, 51% had FIB-4 scores between 1.45-3.25 and 24% had scores > 3.25 . The
16 majority (69%) of HCV-uninfected patients had FIB-4 scores < 1.45 at study start and this
17 remained stable over the entire study period.

18
19 *MICU admissions*

20 During follow-up, 29,887 (19%) patients were admitted to a MICU for a total of 56,805
21 admissions; 61% had one admission, 33% had between two and four admissions, and 6% had
22 five or more admissions. HCV+ patients experienced proportionately more MICU admissions
23 than their HCV-uninfected counterparts in every HIV/ARD strata (42% of HCV+ had > 1
24 admission vs. 38% of HCV-uninfected, $p < 0.0001$) (see Table, Supplemental Digital Content 3).
25 The majority (63%) of MICU admissions occurred within the VA while the remaining 37% were

1 found in Medicare data (Table 2). The most common principal diagnoses for hospital stays with
2 a MICU admission were CVD (28%), respiratory (14%), GI and liver (8%), neurologic (8%),
3 severe sepsis or septic shock (6%), and other infections (6%).
4

5 More HIV+/HCV+/ARD+ patients experienced MICU admission than HIV-/HCV-/ARD-
6 patients (34% vs. 15%, $p<0.0001$) (see Table, Supplemental Digital Content 3). All HCV+
7 (irrespective of HIV infection and ARD status) were more likely than HCV-uninfected to have
8 principal diagnoses related to GI and liver diseases, severe sepsis/septic shock, other
9 infections, endocrine disorders, renal insufficiency, and alcohol- and drug use-related diagnoses
10 (all $p<0.0001$).
11

12 *Trends in MICU admission rates*

13 Overall, annual MICU admission rates decreased by 37% from 37.5 admissions per
14 1000 patients (37.5/1000) in 1997 to 23.7/1000 in 2014 ($t\text{-trend}=-4.8$, $p<0.0001$). The largest
15 decrease was seen among HCV-uninfected patients, irrespective of HIV infection or ARD
16 (Figure 1 with MICU admission rates decreasing by 48% over the study period from 34.0/1000
17 in 1997 to 17.6/1000 in 2014 ($t\text{-trend}=-17.2$, $p<0.0001$). Among HCV+ patients there was no
18 evidence of change in MICU admission rates from 1997 to 2014 ($t\text{-trend}=+1.0$, $p=0.34$).
19

20 *Relative risk for MICU admission*

21 There was minimal evidence of confounding by the adjusting covariates in each
22 exposure group, with relative differences between crude and adjusted RR ranging from 4% to
23 13% (Table 3). After adjusting for baseline age, baseline comorbidity, and calendar year, HIV
24 infection and ARD were each associated with 46% increased risk of MICU admission compared
25 to HIV-/HCV-/ARD- patients (RR 1.46, 95% CI 1.43-1.50 HIV infection; RR 1.46, 95% CI 1.42-

1 1.50 ARD) (Table 3). HCV infection alone was associated with 33% greater risk compared to
2 HIV-/HCV-/ARD- patients (RR 1.33, 95% CI 1.27-1.39). Models stratified by calendar period
3 showed that HIV infection was more strongly associated with risk of MICU admission between
4 1997-2009 (RR 1.61, 95% CI 1.56-1.65) than between 2010-2014 (RR 1.21, 95% CI 1.16-1.27).
5 Conversely, HCV infection was more strongly associated with MICU admission in later years
6 (RR 1.22, 95% CI 1.16-1.29 between 1997-2009; RR 1.54, 95% CI 1.43-1.67 between 2010-
7 2014). The association of ARD was similar in both time periods (RR 1.46, 95% CI 1.41-1.50
8 between 1997-2009; RR 1.44, 95% CI 1.38-1.50 between 2010-2014).

9

10 We found that the effect of ARD differed depending on the presence of other exposures.
11 For example, ARD was associated with 33% (95% CI 1.28-1.38) increased risk of MICU
12 admission among HIV+ patients and 47% (95% CI 1.42-1.52) increased risk among HIV-
13 uninfected patients (Table 3). Similar differential effects of ARD were observed between HCV+
14 and HCV-uninfected patients. Compared to patients with none of the exposures, those with all
15 three had the highest risk of MICU admission (RR 2.36, 95% CI 2.26-2.46).

16

17 **Discussion**

18 Using a national cohort of 155,550 patients and combined data from the largest
19 healthcare provider and insurer in the United States, we showed that HIV, HCV, and ARD were
20 each independently associated with increased risk of MICU admission between 1997 and 2014.
21 Compared to earlier years, risk of MICU admission in the last five years of follow-up decreased
22 among HIV mono-infected patients but increased among HCV mono-infected patients. These
23 findings illustrate that viral infections are associated with serious health conditions requiring
24 MICU admission, but this association may be attenuated by effective treatments as
25 demonstrated in HIV+ patients. Compared to HIV-/HCV-/ARD- patients, the contribution of ARD

1 to the risk of MICU admission remained consistent over the study period. Importantly, ARD
2 accounts for 3.3 million deaths (5.9% of all deaths) each year worldwide¹¹ and continues to be
3 associated with substantial MICU admission risk, signifying the urgency to address this
4 potentially modifiable health behavior.

5
6 MICU admission rates decreased by 37% between 1997 and 2014. This differs from
7 other national data that indicate increased MICU bed count and utilization over the past 15
8 years in the United States.²⁰ Decreasing MICU use in our study may be explained by several
9 factors, including the fixed number of acute care VA facilities, changes to criteria for MICU
10 admission, utilization of MICU resources, or expansion of step-down units. However, we
11 included admissions reimbursed by Medicare, which would be expected to partially offset this
12 effect. In addition, MICU admission rates in our study were substantially higher compared with
13 prior studies of non-surgical ICU admission among the general VA patient population.
14 Approximately 20% of participants in our cohort had at least one MICU admission during the
15 observation period compared with 8% in a cross-sectional study between 2009-2010.²¹ This
16 may be partially explained by differing study designs, including more HIV+ patients in the VACS
17 cohort, who have higher MICU admission rates.

18
19 MICU admission rates fell dramatically between 1997 and 2014 among patients without
20 HCV infection, but were unchanged among patients with HCV infection, suggesting that HCV
21 infection or associated risk factors played a role in MICU use during the study period. Without
22 treatment, the adverse impact of HCV infection on MICU admissions could increase given the
23 prevalence of HCV globally, estimated in 2015 at 1.0% of the general population (71.1 million
24 infections),²² and the lag time between infection and development of HCV-related
25 complications.²³ We anticipate that the influence of HCV on MICU admission rates may change

1 after 2014 when modern direct-acting antivirals (DAA) for treatment of HCV infection became
2 available to VA patients. These therapies are effective in achieving sustained virologic
3 response,¹⁰ but whether this translates to improved clinical outcomes of conditions that
4 commonly require MICU admission, such as sequelae of hepatic decompensation, remains to
5 be seen. The importance of optimally deploying HCV treatments is highlighted by our study's
6 findings of the substantial impact of HCV infection on MICU admission through 2014. Future
7 research should investigate the impact of DAA treatment on MICU admission rates among
8 HCV+ patients with and without HIV co-infection.

9
10 Our findings also highlight the importance of unhealthy alcohol use, a potentially
11 modifiable health behavior, on MICU admission risk in patient populations with or without
12 underlying HIV or HCV infection. In prior work, we showed that ARD are independently
13 associated with all-cause hospitalization.⁴ Our current findings further underscore that patients
14 with ARD also experience serious, potentially life-threatening complications that lead to MICU
15 admission. Although the mechanism through which ARD influences MICU admission risk
16 remains unclear, immune and neurocognitive impairments may play a role. In one study, alcohol
17 dependence was independently associated with sepsis, septic shock, and hospital mortality
18 among adult MICU patients.²⁴ However, other studies suggest that MICU patients with ARD had
19 similar²⁵ or even lower mortality compared to those without ARD.²⁶ Failure to detect harmful
20 effects of ARD in MICU patients must be viewed with caution as these studies were limited by
21 unbalanced comparator groups, with ARD groups being younger and with fewer comorbidities. It
22 is critical to better understand the impact of ARD on important health outcomes such as MICU
23 admission and determine how decreasing or eliminating alcohol use may offset these risks. We
24 postulate that development and subsequent implementation of effective interventions targeting

1 unhealthy alcohol use could reduce MICU admission risk for broad populations of patients and
2 decrease the morbidity and mortality of patients with ARD admitted to the MICU.

3

4 Our study had limitations. First, our findings may reflect bias from admitting clinicians
5 who have a lower threshold for admitting HCV+ patients to the MICU than other patient groups.
6 Some HCV+ patients are at greater risk for infection and may have more significant
7 derangements in biomarkers of infection such as lactate, a benchmark target for monitoring
8 sepsis. Second, our analysis relied on ICD-9 codes for ARD to define alcohol exposure, which
9 have low sensitivity for capturing true ARD status and therefore inflated MICU admission rates
10 among those we classified as ARD-. Finally, nearly all individuals in VACS are men, which
11 resembles the HIV population in many countries, but limits the ability to generalize our findings
12 to other HIV cohorts with higher proportions of women.

13

14 In summary, while MICU admission rates in this large, national cohort fell dramatically
15 between 1997 and 2014 among patients without HCV, there was essentially no change in MICU
16 admission rates among patients with HCV infection. While the additional risk of MICU admission
17 conferred by HIV mono-infection decreased over time, both HCV infection and ARD remained
18 major risk factors for MICU admission between 2010 and 2014. The impact of these conditions
19 could substantially change with the combination of safe and effective treatment for HIV and
20 HCV infections, and interventions targeting unhealthy alcohol use.

21

22 **Acknowledgements**

23 Support for VA/CMS data is provided by the Department of Veterans Affairs, Veterans Health
24 Administration, Office of Research and Development, Health Services Research and
25 Development, and VA Information Resource Center (SDR 02-237 and 98-004). This work was

1 supported by the US National Institutes of Health (NIH), including grants from the National
2 Institute on Alcohol Abuse and Alcoholism [U24-AA020794, U01-AA020790, U10 AA013566-
3 completed to ACJ] and National Heart, Lung, and Blood Institute [R01 HL090342 to KC, K24
4 HL087713 to LH]. The views expressed in this manuscript represent those of the authors and do
5 not necessarily represent those of the Department of Veterans Affairs.

6

7 **References**

- 8 1. Crothers K, Huang L, Goulet JL, et al. HIV infection and risk for incident pulmonary
9 diseases in the combination antiretroviral therapy era. *Am J Respir Crit Care Med.*
10 2011;183(3):388-395.
- 11 2. Freiberg MS, Chang CH, Skanderson M, et al. Association Between HIV Infection and
12 the Risk of Heart Failure With Reduced Ejection Fraction and Preserved Ejection
13 Fraction in the Antiretroviral Therapy Era: Results From the Veterans Aging Cohort
14 Study. *JAMA Cardiol.* 2017;2(5):536-546.
- 15 3. Crum-Cianflone NF, Grandits G, Echols S, Ganesan A, Landrum M, Weintrob A. Trends
16 and Causes of Hospitalizations Among HIV-Infected Persons During the Late HAART
17 Era: What Is the Impact of CD4 Counts and HAART Use? *J Acquir Immune Defic Syndr.*
18 2010;54:248-257.
- 19 4. Rentsch C, Tate JP, Akgun KM, et al. Alcohol-Related Diagnoses and All-Cause
20 Hospitalization Among HIV-Infected and Uninfected Patients: A Longitudinal Analysis of
21 United States Veterans from 1997 to 2011. *AIDS and Behav.* 2016;20(3):555-564.
- 22 5. Palella FJJ, Baker RK, Moorman AC, et al. Mortality in the Highly Active Antiretroviral
23 Therapy Era: Changing Causes of Death and Disease in the HIV Outpatient Study. *J*
24 *Acquir Immune Defic Syndr.* 2006;43:27-34.

- 1 6. Akgun KM, Pisani M, Crothers K. The changing epidemiology of HIV-infected patients in
2 the intensive care unit. *J Intensive Care Med.* 2011;26(3):151-164.
- 3 7. Akgun KM, Tate JP, Pisani M, et al. Medical ICU admission diagnoses and outcomes in
4 human immunodeficiency virus-infected and virus-uninfected veterans in the
5 combination antiretroviral era. *Crit Care Med.* 2013;41(6):1458-1467.
- 6 8. Platt L, Easterbrook P, Gower E, et al. Prevalence and burden of HCV co-infection in
7 people living with HIV: a global systematic review and meta-analysis. *Lancet Infect Dis.*
8 2016;16(7):797-808.
- 9 9. Crowell TA, Gebo KA, Balagopal A, et al. Impact of hepatitis coinfection on
10 hospitalization rates and causes in a multicenter cohort of persons living with HIV. *J*
11 *Acquir Immune Defic Syndr.* 2014;65(4):429-437.
- 12 10. Jakobsen JC, Nielsen EE, Feinberg J, et al. Direct-acting antivirals for chronic hepatitis
13 C. *Cochrane Database Syst Rev.* 2017;6:CD012143.
- 14 11. World Health Organization. Global status report on alcohol and health. 2014.
- 15 12. Gacouin A, Tadie JM, Uhel F, et al. At-risk drinking is independently associated with ICU
16 and one-year mortality in critically ill nontrauma patients. *Crit Care Med.* 2014;42(4):860-
17 867.
- 18 13. Mankal PK, Abed J, Aristy JD, et al. Relative effects of heavy alcohol use and hepatitis C
19 in decompensated chronic liver disease in a hospital inpatient population. *Am J Drug*
20 *Alcohol Abuse.* 2015;41(2):177-182.
- 21 14. Akgun KM, Gordon K, Pisani M, et al. Risk factors for hospitalization and medical
22 intensive care unit (MICU) admission among HIV infected Veterans. *J Acquir Immune*
23 *Defic Syndr.* 2013;62(1):52-59.

- 1 15. Justice AC, McGinnis KA, Tate JP, et al. Risk of mortality and physiologic injury evident
2 with lower alcohol exposure among HIV infected compared with uninfected men. *Drug*
3 *Alcohol Depend.* 2016;161:95-103.
- 4 16. Marshall BDL, Tate JP, McGinnis KA, et al. Long-term alcohol use patterns and HIV
5 disease severity. *AIDS.* 2017;31(9):1313-1321.
- 6 17. Lo Re V, III, Kallan MJ, Tate JP, et al. Hepatic Decompensation in Antiretroviral-Treated
7 Patients Co-Infected With HIV and Hepatitis C Virus Compared With Hepatitis C Virus-
8 Monoinfected Patients. *Ann Intern Med.* 2014;160:369-379.
- 9 18. Fultz SL, Skanderson M, Mole L, et al. Development and verification of a "virtual" cohort
10 using the national VA Health Information System. *Med Care.* 2006;44(8):S25-S30.
- 11 19. McGinnis KA, Justice AC, Kraemer KL, Saitz R, Bryant KJ, Fiellin DA. Comparing
12 alcohol screening measures among HIV-infected and -uninfected men. *Alcoholism,*
13 *clinical and experimental research.* 2013;37(3):435-442.
- 14 20. Wallace DJ, Angus DC, Seymour CW, Barnato AE, Kahn JM. Critical care bed growth in
15 the United States. A comparison of regional and national trends. *Am J Respir Crit Care*
16 *Med.* 2015;191(4):410-416.
- 17 21. Chen LM, Render M, Sales A, Kennedy EH, Wiitala W, Hofer TP. Intensive care unit
18 admitting patterns in the Veterans Affairs health care system. *Arch Intern Med.*
19 2012;172(16):1220-1226.
- 20 22. Polaris Observatory HCV Collaborators. Global prevalence and genotype distribution of
21 hepatitis C virus infection in 2015: a modelling study. *Lancet Gastroenterol Hepatol.*
22 2017;2(3):161-176.
- 23 23. Davis GL, Alter MJ, El-Serag H, Poynard T, Jennings LW. Aging of hepatitis C virus
24 (HCV)-infected persons in the United States: a multiple cohort model of HCV prevalence
25 and disease progression. *Gastroenterology.* 2010;138(2):513-521.

- 1 24. O'Brien JM, Jr., Lu B, Ali NA, et al. Alcohol dependence is independently associated with
2 sepsis, septic shock, and hospital mortality among adult intensive care unit patients. *Crit*
3 *Care Med.* 2007;35(2):345-350.
- 4 25. Lynch C, Pugh R, Battle C, Welsh Intensive Care Society Audit Research Group
5 (WICSARG). A multicentre prospective evaluation of alcohol-related admissions to
6 intensive care units in Wales. *J Intensive Care Soc.* 2017;18(3):193-197.
- 7 26. Hietanen S, Ala-Kokko T, Ohtonen P, Käkälä R, Niemelä S, Liisanantti JH. Treatment
8 Profile and 1-Year Mortality Among Nontraumatic Intensive Care Unit Patients With
9 Alcohol-Related Health Problems. *J of Intensive Care Med.* 2017. DOI:
10 10.1177/0885066617740071. Accessed Oct 5, 2018.

11

12 **Figure captions**

13 **Figure 1.**

14 **Title:** Age-adjusted medical intensive care unit admission rates per 1000 at-risk patients
15 enrolled in the Veterans Aging Cohort Study between 1997 and 2014, by HIV, hepatitis C virus
16 (HCV), and time-updated alcohol-related diagnosis (ARD)

17 **Note (below figure):** Patients were classified as ARD+ if in the previous year they had at least
18 two outpatient encounters with an ARD; they remained ARD+ thereafter.

19

20 **Supplemental Digital Content**

21 Table, Supplemental Digital Content 1.doc

1 Table, Supplemental Digital Content 2.doc

2 Table, Supplemental Digital Content 3.doc

Table 1. Demographic and clinical characteristics among patients receiving care in the Veterans Aging Cohort Study between 1997 and 2014, n=155,550

Characteristic	n (%)
Exposure group	
<i>HIV+ / ARD+ / HCV+</i>	5144 (3.3)
<i>HIV+ / ARD+ / HCV-</i>	9359 (6.0)
<i>HIV+ / ARD- / HCV+</i>	4743 (3.1)
<i>HIV+ / ARD- / HCV-</i>	30534 (19.6)
<i>HIV- / ARD+ / HCV+</i>	6682 (4.3)
<i>HIV- / ARD+ / HCV-</i>	25924 (16.7)
<i>HIV- / ARD- / HCV+</i>	3653 (2.4)
<i>HIV- / ARD- / HCV-</i>	69511 (44.7)
Age at baseline, median (IQR)	47 (40-54)
Male sex	151,379 (97.3)
Race/ethnicity	
<i>Black</i>	73,221 (47.1)
<i>White</i>	62,084 (39.9)
<i>Hispanic</i>	12,761 (8.2)
<i>Other or unknown</i>	7,484 (4.8)
Year of VACS enrollment	
<i>1997-2001</i>	82,144 (52.8)
<i>2002-2006</i>	33,134 (21.3)
<i>2007-2010</i>	20,659 (13.3)
<i>2011-2014</i>	19,613 (12.6)
Comorbidities at baseline	
<i>Psychiatric disorders</i>	41,614 (26.8)
<i>Hypertension</i>	41,253 (26.5)
<i>Drug use-related diagnoses</i>	21,415 (13.8)
<i>Diabetes</i>	18,106 (11.6)
<i>CVD</i>	9,570 (6.2)
<i>Anemia</i>	8,369 (5.4)
<i>COPD</i>	7,043 (4.5)
<i>Non-AIDS cancers</i>	5,340 (3.4)
<i>Renal insufficiency</i>	3,764 (2.4)
<i>Chronic hepatitis B</i>	1,868 (1.2)
<i>Liver diseases</i>	1,560 (1.0)
Mortality per 100 person-years	2.6
Ever MICU admission	29,887 (19.2)

Abbreviations: HIV - human immunodeficiency virus; VACS - Veterans Aging Cohort Study; HCV - hepatitis C virus; ARD - alcohol-related diagnosis; IQR - interquartile range; MICU - medical intensive care unit

Note: ARD status was used as an ever/never measure in this table

Table 2. Characteristics of medical intensive care unit (MICU) admissions, n=56,805

Characteristic	n (%)
Location	
VA	35,883 (63.2)
Medicare	20,922 (36.8)
LOS per admission in days, median (IQR)	6 (3-11)
Primary admission diagnosis	
CVD	16,151 (28.4)
Respiratory	7,682 (13.5)
GI and liver	4,770 (8.4)
Neurologic	4,374 (7.7)
Other infections	3,611 (6.4)
Severe sepsis/septic shock	3,469 (6.1)
Endocrine	3,248 (5.7)
HIV/AIDS	2,671 (4.7)
Renal insufficiency	2,224 (3.9)
Non-AIDS cancers	1,241 (2.2)
ARD	950 (1.7)
Psychiatric disorders	771 (1.4)
Drug use-related diagnoses	344 (0.6)
Other	5,299 (9.3)

Abbreviations: IQR - interquartile range; HIV - human immunodeficiency virus; HCV - hepatitis C virus; ARD - alcohol-related diagnosis; MICU - medical intensive care unit; VA - Veterans Affairs; LOS - length of stay; CVD - cardiovascular disease; AIDS - acquired immune deficiency syndrome


Table 3. Risk of medical intensive care unit (MICU) admission by HIV, hepatitis C virus (HCV), alcohol-related diagnosis (ARD), and calendar era, n=155,550

All groups						
Exposure group			Crude 1997-2014	Adjusted 1997-2014	Adjusted 1997-2009	Adjusted 2010-2014
HIV+	ARD+	HCV+	2.26 (1.17-2.34)	2.36 (2.26-2.46)	2.29 (2.17-2.42)	2.40 (2.23-2.58)
		HCV-	1.89 (1.83-1.95)	2.14 (2.06-2.22)	2.34 (2.24-2.44)	1.75 (1.64-1.86)
	ARD-	HCV+	1.64 (1.58-1.71)	1.74 (1.67-1.81)	1.66 (1.58-1.75)	1.94 (1.80-2.09)
		HCV-	1.38 (1.35-1.41)	1.46 (1.43-1.50)	1.61 (1.56-1.65)	1.21 (1.16-1.27)
HIV-	ARD+	HCV+	1.72 (1.66-1.79)	1.80 (1.73-1.88)	1.69 (1.60-1.78)	1.91 (1.78-2.04)
		HCV-	1.37 (1.33-1.40)	1.46 (1.42-1.50)	1.46 (1.41-1.50)	1.44 (1.38-1.50)
	ARD-	HCV+	1.25 (1.20-1.31)	1.33 (1.27-1.39)	1.22 (1.16-1.29)	1.54 (1.43-1.67)
		HCV-	ref	ref	ref	ref
By HIV						
HIV+	HCV+/ARD+		1.64 (1.57-1.70)	1.62 (1.55-1.70)	1.45 (1.37-1.53)	1.98 (1.83-2.14)
	HCV+/ARD-		1.19 (1.14-1.24)	1.19 (1.14-1.24)	1.04 (0.99-1.09)	1.60 (1.48-1.73)
	HCV-/ARD+		1.37 (1.32-1.42)	1.33 (1.28-1.38)	1.33 (1.28-1.39)	1.29 (1.21-1.38)
	HCV-/ARD-		ref	ref	ref	ref
HIV-	HCV+/ARD+		1.72 (1.66-1.79)	1.76 (1.69-1.84)	1.65 (1.56-1.74)	1.88 (1.75-2.02)
	HCV+/ARD-		1.25 (1.20-1.31)	1.29 (1.23-1.35)	1.19 (1.12-1.26)	1.52 (1.41-1.65)
	HCV-/ARD+		1.37 (1.33-1.41)	1.47 (1.42-1.52)	1.49 (1.43-1.55)	1.36 (1.29-1.44)
	HCV-/ARD-		ref	ref	ref	ref
By HCV						
HCV+	HIV+/ARD+		1.80 (1.69-1.92)	1.78 (1.66-1.90)	1.88 (1.73-2.04)	1.55 (1.39-1.74)
	HIV+/ARD-		1.31 (1.26-1.37)	1.31 (1.25-1.37)	1.36 (1.29-1.43)	1.26 (1.17-1.35)
	HIV-/ARD+		1.37 (1.32-1.43)	1.36 (1.30-1.42)	1.38 (1.31-1.46)	1.24 (1.14-1.34)
	HIV-/ARD-		ref	ref	ref	ref
HCV-	HIV+/ARD+		1.89 (1.83-1.95)	2.14 (2.06-2.22)	2.34 (2.24-2.44)	1.75 (1.64-1.86)
	HIV+/ARD-		1.38 (1.35-1.41)	1.46 (1.43-1.50)	1.60 (1.56-1.65)	1.21 (1.16-1.27)
	HIV-/ARD+		1.37 (1.33-1.40)	1.46 (1.42-1.50)	1.46 (1.41-1.50)	1.44 (1.38-1.50)
	HIV-/ARD-		ref	ref	ref	ref

Notes: reported as RR (95% confidence intervals); models adjusted for baseline age, comorbidity (psychiatric disorders, hypertension, drug use-related diagnoses, diabetes, cardiovascular disease, anemia, chronic obstructive pulmonary disorder, non-AIDS cancers, renal insufficiency, chronic hepatitis B, and liver diseases), and calendar year

Abbreviations: HIV - human immunodeficiency virus; RR - rate ratio; CI - confidence interval; ref - referent group

Figure 1. Age-adjusted medical intensive care unit admission rates per 1000 at-risk patients enrolled in the Veterans Aging Cohort Study between 1997 and 2014, by HIV, hepatitis C virus (HCV), and time-updated alcohol-related diagnosis (ARD)



Note: Patients were classified as ARD+ if in the previous year they had at least two outpatient encounters with an ARD; they remained ARD+ thereafter.

Supplemental Table 1. Categorization of ICD-9 codes for principal admission diagnoses and comorbidities

Category	ICD-9 codes
Alcohol-related diagnoses	291.X; 303.X; 305-305.03; 357.5; 425.5; 535.3; 571-571.3; 790.3; 980, 980.8-980.9; E860.0/.1/.8/.9; V11.3
Anemia	280-281.9; 282.1/.2/.3/.4/.5/.6X/.7/.8/.9; 283.X; 284/.8/.9; 285/.2X/.8/.9
Cardiovascular disease	401.X; 402.01-.91; 404-404.11; 404.13/.91/.93; 410-414.07; 414.8/.9; 423-428.9; 429.7; 433-437.1; 437.6-438.12; 438.2-.53; 438.81-.82; 438.89-441.9; 443/.1/.8X/.9; 447.1; 557.X; 780.2; 785.1-785.4; 786.5X; 996-996.1; 996.71/.72; V43.4; V45.81/.82
Chronic hepatitis B	070.2-070.33; 070.42/.52; V02.61
COPD	490/491.1/.2/.21/.8/.9; 492.X; 496
Diabetes	250-250.93; 357.2
Drug use-related diagnoses	292.X; 304-305.93
Endocrine	240-279.99
GI and liver	280.X; 520-579.99; 787.X; 789-789.09; V55.3
HIV/AIDS	042-044; 176.X; 200-202.98; V08
Hypertension	401-405.99; 437.2
Liver diseases	070; 070.2X/.44/.6; 456-456.2; 570; 571.2/.5/.6; 572.2/.3/.4/.8; 789.5
Psychiatric disorders	293-301.9; 309.X; 311; V11
Neurologic	290.4; 320-389.99; 430-459.99; 721.3; 722.X; 723.0/.4; 724.0X/.2/.4; 780.0X/.3X/.4/.97; 781.2/.3; 784; 852.2
Non-AIDS cancers	140-175.9; 179-199.2; 201.X; 202.3-.68; 203-208.92 (excluding remission codes); V58.0; V58.1; V58.11
Other infections	001-139.99 (excluding 042-044); 421.0; 424.90; 599.0; 604.X; 681-682.9; 711.0X; 730.X; 780.6; 790.7; 995.91-995.94; 996.6X; 997.62; 998.5X; 999.3X
Renal insufficiency	285.21; 403.01/.11/.90/.91; 404.02/.03/.12/.13/.92/.93; 580-594.9; 596.X; 599.7; 600.0X; 788.2; 792.5; 996.3X; 996.73; 996.81; 997.5; V42; V45.1; V56.X
Respiratory	415-416.9; 460-519.99; 780.57; 786.0X/.2/.52
Severe sepsis/septic shock	020.0; 038; 096.7; 112.5/.81; 286.6/.9; 287.4/.5; 293.0; 343.3; 348.1; 458.0; 570.0; 573.4; 584.0; 785.50/.52; 995.92

Abbreviations: ICD-9 - The International Classification of Diseases, Ninth Revision; COPD - chronic obstructive pulmonary disorder; GI - gastrointestinal; HIV - human immunodeficiency virus; AIDS - acquired immune deficiency syndrome

Supplemental Table 2. Demographic and clinical characteristics by HIV, hepatitis C virus (HCV) infection, and alcohol-related diagnosis (ARD) among patients receiving care in the Veterans Aging Cohort Study between 1997 and 2014, n=155,550

Characteristic	HIV+/ARD+		HIV+/ARD-		HIV-/ARD+		HIV-/ARD-	
	HCV+ (n=5,144)	HCV- (n=9,359)	HCV+ (n=4,743)	HCV- (n=30,534)	HCV+ (n=6,682)	HCV- (n=25,924)	HCV+ (n=3,653)	HCV- (n=69,511)
Age at baseline, median (IQR)	47 (42-51)	45 (39-51)	49 (44-54)	47 (39-55)	47 (43-52)	46 (40-52)	49 (45-54)	48 (40-56)
Male sex	5,050 (98.2)	9,086 (97.1)	4,671 (98.5)	29,713 (97.3)	6,626 (99.2)	25,545 (98.5)	3,597 (98.5)	67,091 (96.5)
Race/ethnicity								
<i>Black</i>	3,358 (65.3)	5,271 (56.3)	2,546 (53.7)	12,711 (41.6)	4,122 (61.7)	13,864 (53.5)	2,069 (56.6)	29,280 (42.1)
<i>White</i>	1,299 (25.3)	3,186 (34.0)	1,487 (31.4)	13,588 (44.5)	1,999 (29.9)	9,461 (36.5)	1,162 (31.8)	29,902 (43.0)
<i>Hispanic</i>	412 (8.0)	644 (6.9)	541 (11.4)	2,361 (7.7)	493 (7.4)	2,028 (7.8)	340 (9.3)	5,942 (8.6)
<i>Other or unknown</i>	75 (1.5)	258 (2.8)	169 (3.6)	1,874 (6.1)	68 (1.0)	571 (2.2)	82 (2.2)	4,387 (6.3)
Year of VACS enrollment								
1997-2001	3,407 (66.2)	5,355 (57.2)	2,709 (57.1)	15,078 (49.4)	4,251 (63.6)	15,271 (58.9)	2,054 (56.2)	34,019 (48.9)
2002-2006	1,040 (20.2)	1,916 (20.5)	1,040 (21.9)	6,125 (20.1)	1,388 (20.8)	5,011 (19.3)	801 (21.9)	15,813 (22.8)
2007-2010	464 (9.0)	1,272 (13.6)	511 (10.8)	4,395 (14.4)	576 (8.6)	2,994 (11.6)	467 (12.8)	9,980 (14.4)
2011-2014	233 (4.5)	816 (8.7)	483 (10.2)	4,936 (16.2)	467 (7.0)	2,648 (10.2)	331 (9.1)	9,699 (14.0)
Comorbidities at baseline, frequency								
<i>None</i>	1,358 (26.4)	2,841 (30.4)	2,221 (46.8)	15,438 (50.6)	1,856 (27.8)	8,099 (31.2)	1,383 (37.9)	32,733 (47.1)
<i>One</i>	1,386 (26.9)	2,696 (28.8)	1,297 (27.4)	8,286 (27.1)	1,929 (28.9)	8,394 (32.4)	1,094 (30.0)	20,529 (29.5)
<i>Two or more</i>	2,400 (46.7)	3,822 (40.8)	1,225 (25.8)	6,810 (22.3)	2,897 (43.4)	9,431 (36.4)	1,176 (32.2)	16,249 (23.4)
Comorbidities at baseline, by type								
<i>Psychiatric disorders</i>	2,069 (40.2)	3,762 (40.2)	1,005 (21.2)	6,344 (20.8)	2,696 (40.4)	10,350 (39.9)	995 (27.2)	14,393 (20.7)
<i>Hypertension</i>	1,123 (21.8)	1,936 (20.7)	1,029 (21.7)	5,908 (19.4)	1,829 (27.4)	7,127 (27.5)	1,205 (33.0)	21,096 (30.4)
<i>Drug use-related diagnoses</i>	2,754 (53.5)	3,605 (38.5)	698 (14.7)	1,733 (5.7)	3,053 (45.7)	7,463 (28.8)	443 (12.1)	1,666 (2.4)
<i>Diabetes</i>	434 (8.4)	558 (6.0)	453 (9.6)	2,491 (8.2)	755 (11.3)	2,430 (9.4)	625 (17.1)	10,360 (14.9)
<i>CVD</i>	205 (4.0)	482 (5.2)	229 (4.8)	1,692 (5.5)	267 (4.0)	1,348 (5.2)	225 (6.2)	5,122 (7.4)
<i>Anemia</i>	431 (8.4)	942 (10.1)	436 (9.2)	3,237 (10.6)	219 (3.3)	943 (3.6)	139 (3.8)	2,022 (2.9)
<i>COPD</i>	297 (5.8)	651 (7.0)	199 (4.2)	1,270 (4.2)	338 (5.1)	1,428 (5.5)	148 (4.1)	2,712 (3.9)

<i>Non-AIDS cancers</i>	117 (2.3)	262 (2.8)	145 (3.1)	1,332 (4.4)	158 (2.4)	615 (2.4)	111 (3.0)	2,600 (3.7)
<i>Renal insufficiency</i>	154 (3.0)	314 (3.4)	243 (5.1)	1,292 (4.2)	114 (1.7)	301 (1.2)	121 (3.3)	1,225 (1.8)
<i>Chronic hepatitis B</i>	221 (4.3)	336 (3.6)	153 (3.2)	746 (2.4)	134 (2.0)	139 (0.5)	34 (0.9)	105 (0.2)
<i>Liver diseases</i>	160 (3.1)	215 (2.3)	128 (2.7)	296 (1.0)	175 (2.6)	341 (1.3)	93 (2.6)	152 (0.2)
Mortality per 100 person-years	3.8	3.9	4.5	4.6	2.3	1.9	2.4	1.9
Ever MICU admission	1,736 (33.8)	2,445 (26.1)	1,369 (28.9)	6,117 (20.0)	1,811 (27.1)	5,097 (19.7)	797 (21.8)	10,515 (15.1)

Abbreviations: HIV - human immunodeficiency virus; VACS - Veterans Aging Cohort Study; HCV - hepatitis C virus; ARD - alcohol-related diagnosis; IQR - interquartile range; MICU - medical intensive care unit

Note: ARD status was used as an ever/never measure in this table; all comparisons across exposure groups $p < 0.0001$ using non-parametric Kruskal-Wallis test for continuous measures and chi-square tests for categorical measures

Supplemental Table 3. Characteristics of medical intensive care unit (MICU) patients by HIV, hepatitis C virus (HCV) infection, and alcohol-related diagnosis (ARD), n=29,887

	HIV+/ARD+		HIV+/ARD-		HIV-/ARD+		HIV-/ARD-	
	HCV+ (n=1,736)	HCV- (n=2,445)	HCV+ (n=1,369)	HCV- (n=6,117)	HCV+ (n=1,811)	HCV- (n=5,097)	HCV+ (n=797)	HCV- (n=10,515)
Admitted patients, median (IQR)								
Years to first MICU admission	6 (3-10)	4 (2-8)	6 (2-10)	4 (1-7)	6 (3-11)	6 (3-10)	7 (3-11)	6 (2-10)
Number of MICU admissions								
1 admission	981 (56.5)	1,525 (62.4)	844 (61.7)	4,146 (67.8)	1,021 (56.4)	3,038 (59.6)	457 (57.3)	6,192 (58.9)
2-4 admissions	632 (36.4)	784 (32.1)	455 (33.2)	1,715 (28.0)	644 (35.6)	1,690 (33.2)	273 (34.3)	3,612 (34.4)
≥5 admissions	123 (7.1)	136 (5.6)	70 (5.1)	256 (4.2)	146 (8.1)	369 (7.2)	67 (8.4)	711 (6.8)
	HIV+/ARD+		HIV+/ARD-		HIV-/ARD+		HIV-/ARD-	
	HCV+ (n=3,495)	HCV- (n=4,478)	HCV+ (n=2,470)	HCV- (n=10,195)	HCV+ (n=3,746)	HCV- (n=10,211)	HCV+ (n=1,612)	HCV- (n=20,598)
Admissions (n=56,805)								
Location								
VA	2,591 (74.1)	3,138 (70.1)	1,711 (69.3)	6,494 (63.7)	2,820 (75.3)	7,263 (71.1)	1,015 (63.0)	10,851 (52.7)
Medicare	904 (25.9)	1,340 (29.9)	759 (30.7)	3,701 (36.3)	926 (24.7)	2,948 (28.9)	597 (37.0)	9,747 (47.3)
LOS per admission, days	6 (3-13)	6 (3-13)	7 (3-13)	6 (3-13)	5 (2-11)	5 (2-10)	5 (3-11)	5 (3-10)
Primary admission diagnosis								
CVD	556 (15.9)	839 (18.7)	435 (17.6)	2,384 (23.4)	830 (22.2)	2,919 (28.6)	462 (28.7)	7,726 (37.5)
Respiratory	454 (13.0)	648 (14.5)	335 (13.6)	1,341 (13.2)	374 (10.0)	1,472 (14.4)	193 (12.0)	2,865 (13.9)
GI and liver	362 (10.4)	345 (7.7)	219 (8.9)	723 (7.1)	522 (13.9)	1,074 (10.5)	144 (8.9)	1,381 (6.7)
Neurologic	232 (6.6)	315 (7.0)	172 (7.0)	700 (6.9)	271 (7.2)	775 (7.6)	132 (8.2)	1,777 (8.6)
Other infections	243 (7.0)	279 (6.2)	215 (8.7)	845 (8.3)	222 (5.9)	396 (3.9)	137 (8.5)	1,274 (6.2)
Severe sepsis/septic shock	251 (7.2)	285 (6.4)	238 (9.6)	737 (7.2)	200 (5.3)	426 (3.2)	115 (7.1)	1,217 (5.9)
Endocrine	203 (5.8)	193 (4.3)	127 (5.1)	393 (3.9)	297 (7.9)	729 (7.1)	108 (6.7)	1,198 (5.8)
HIV/AIDS	284 (8.1)	519 (11.6)	274 (11.1)	1,594 (15.6)	-	-	-	-
Renal insufficiency	173 (5.0)	177 (4.0)	153 (6.2)	442 (4.3)	145 (3.9)	286 (2.8)	95 (5.9)	753 (3.7)
Non-AIDS cancers	54 (1.6)	69 (1.5)	49 (2.0)	208 (2.0)	82 (2.2)	184 (1.8)	34 (2.1)	561 (2.7)

<i>ARD</i>	57 (1.6)	51 (1.1)	11 (0.5)	20 (0.2)	64 (1.7)	105 (1.0)	10 (0.6)	26 (0.1)
<i>Psychiatric disorders</i>	84 (2.4)	110 (2.5)	12 (0.5)	54 (0.5)	92 (2.5)	233 (2.3)	13 (0.8)	173 (0.8)
<i>Drug use-related diagnoses</i>	57 (1.6)	51 (1.1)	11(0.5)	20 (0.2)	64 (1.7)	105 (1.0)	10 (0.6)	26 (0.1)
<i>Other</i>	431 (12.3)	519 (11.6)	228 (9.2)	745 (7.3)	457 (12.2)	1,121 (11.0)	167 (10.4)	1,631 (7.9)

Abbreviations: HIV - human immunodeficiency virus; HCV - hepatitis C virus; ARD - alcohol-related diagnosis; MICU - medical intensive care unit; IQR - interquartile range; VA - Veterans Affairs; LOS - length of stay; CVD - cardiovascular disease; AIDS - acquired immune deficiency syndrome

Notes: ARD status was used as an ever/never measure in this table; all comparisons across exposure groups $p < 0.0001$ using non-parametric Kruskal-Wallis test for continuous measures and chi-square tests for categorical measures