

Population-Based Estimate of Melioidosis, Kenya

**Esther M. Muthumbi, Nicola C. Gordon,
George Mochamah, Sammy Nyongesa,
Emily Odipo, Salim Mwarumba, Neema Mturi,
Anthony O. Etyang, David A.B. Dance,
J. Anthony G. Scott, Susan C. Morpeth**

Melioidosis is thought to be endemic, although underdiagnosed, in Africa. We identified 5 autochthonous cases of *Burkholderia pseudomallei* infection in a case series in Kenya. Incidence of *B. pseudomallei* bacteremia in Kenya's Kilifi County is low, at 1.5 cases per million person-years, but this result might be an underestimate.

Burkholderia pseudomallei, the causative agent of melioidosis, is a gram-negative bacillus endemic particularly in northern Australia and South and Southeast Asia. Worldwide, *B. pseudomallei* causes $\approx 165,000$ cases of disease and $\approx 89,000$ deaths annually (1). The presence of *B. pseudomallei* in Africa has been demonstrated by sporadic cases of melioidosis reported in travelers returning from countries including Kenya (2). Indigenous culture-confirmed cases have been reported in only 4 countries in Africa, mainly from research centers with diagnostic laboratory facilities (3).

The first case of melioidosis linked to Kenya was diagnosed in 1982 in a tourist from Denmark who had visited Nyali (an area of Mombasa City), ≈ 50 km south of the town of Kilifi (2). Follow-up clinical surveillance in Nairobi and environmental surveillance from other regions in Kenya yielded no *B. pseudomallei* isolates (4). However, growing concerns over possible underestimation of the disease in potentially endemic areas, including in tropical Africa, have led to calls for improved surveillance (5).

In 2010, at Kilifi County Hospital (KCH), we isolated *B. pseudomallei* from the blood culture of a 3-year-old child after a near-drowning accident in a seasonal river.

Author affiliations: KEMRI–Wellcome Trust Research Programme, Kilifi, Kenya (E.M. Muthumbi, N.C. Gordon, G. Mochamah, S. Nyongesa, E. Odipo, S. Mwarumba, N. Mturi, A.O. Etyang, J.A.G. Scott, S.C. Morpeth); London School of Hygiene & Tropical Medicine, London, UK (N.C. Gordon, D.A.B. Dance, J.A.G. Scott, S.C. Morpeth); Lao-Oxford-Mahosot Hospital–Wellcome Trust Research Unit, Vientiane, Laos (D.A.B. Dance); University of Oxford, Oxford, UK (D.A.B. Dance, J.A.G. Scott, S.C. Morpeth)

DOI: <https://doi.org/10.3201/eid2505.180545>

The identity of the isolate was confirmed by real-time PCR targeting the type III secretion system genes of *B. pseudomallei* (6), and the isolate was later sequenced for a study of geographic dissemination of *B. pseudomallei* (7). After this identification, we conducted a retrospective analysis of archived blood culture isolates collected during 1994–2012 to investigate possible missed cases of invasive *B. pseudomallei* infection.

The Study

During 1994–1998, blood culture was performed on all febrile patients admitted to the pediatric wards at KCH. Since 1998, all pediatric patients <15 years of age admitted, except those having trauma, burns, or elective surgery, have had blood samples drawn for culture. Surveillance for patients ≥ 15 years of age began in 2007; blood samples are drawn at admission for cultures on patients meeting clinical criteria for possible invasive bacterial disease. Since 2002, hospitalization events have been linked to the Kilifi Health and Demographic Surveillance System (KHDSS), which monitors the population of $\approx 280,000$ over an area of 891 km² (8). Informed consent is obtained from all patients participating in the surveillance, including for storage of isolates and future use of clinical data.

Blood samples for bacterial cultures were collected in BACTEC Peds Plus or BACTEC Plus Aerobic/F bottles (Becton Dickinson, <https://www.bd.com>) and incubated on a BACTEC FX 9050 Automated Blood Culture instrument (Becton Dickinson). Nonfastidious, oxidase-positive, gram-negative bacilli were identified by using API 20NE test kits (bioMérieux, <https://www.biomerieux.com>). We reviewed all gentamicin-resistant, glucose-nonfermenting, gram-negative rods, with the exception of *Pseudomonas aeruginosa*, even if the API 20NE identification was acceptable, to account for difficulties in speciating *Burkholderia* spp. with biochemical methods.

A total of 86,582 patients <15 years of age were admitted during 1994–2012 and 18,864 patients ≥ 15 years of age during 2007–2012. Surveillance identified 33 gentamicin-resistant, glucose-nonfermenting bacilli in 14,235 positive blood cultures from patients <15 years of age and 5 gentamicin-resistant, glucose-nonfermenting bacilli in 705 positive blood cultures from patients ≥ 15 years of age (Figure). We retrieved all 38 isolates from storage for PCR, which we performed using published primer and probe sequences (6).




Figure. Identification of gentamicin-resistant, glucose-nonfermenting bacilli and *Burkholderia pseudomallei* from isolates collected from patients at Kilifi County Hospital, Kilifi, Kenya, 1994–2012.

We identified 4 isolates as *B. pseudomallei* by PCR, including the index isolate from 2010 (Table 1; Appendix, <https://wwwnc.cdc.gov/EID/25/5/18-0545-App1.pdf>). One isolate was previously identified as *B. cepacia*, and 2 were previously labeled as *Pseudomonas* species. We identified a fifth *B. pseudomallei* case in July 2014 in a 68-year-old female patient with diabetes mellitus and bilateral cervical abscesses (Table 1; Appendix). Blood culture results were negative, but aspirated pus grew *B. pseudomallei*, identified by API 20NE and confirmed by PCR.

None of the case-patients had any history of travel outside Kilifi County. Three died during the course of their

admission. No further information is available for the 2 case-patients who survived because they were not residents of the area surveyed by KHDSS.

To estimate the incidence of melioidosis bloodstream infection, we divided the number of invasive *B. pseudomallei* cases among KHDSS residents by the sum of the annual midyear population counts during 2002–2012 for those <15 years of age and during 2007–2012 for those ≥15 years of age. We also adjusted for the sensitivity of the surveillance to account for the proportion of patients not consenting to the surveillance study and those who did not have a blood culture drawn. For the period before 2002, we extrapolated age-specific population estimates by using a log-linear model of age-specific population data based on subsequent enumerations. The estimated incidence was 1.3 cases/1 million person-years of observation for those <15 years of age and 2 cases/1 million person-years of observation for those ≥15 years of age (Table 2).

Conclusions

We identified 5 cases of melioidosis from a single surveillance site in Kenya. Despite reports suggesting that melioidosis is endemic but underdetected in the region (5), we demonstrated low incidence in this part of Kenya. Even so, *B. pseudomallei* has emerged as an underdiagnosed cause of sepsis in Kilifi County. The empirical treatment used for sepsis, ampicillin and gentamicin, does not cover *B. pseudomallei*. The lack of pathogenomic clinical features makes it difficult to detect melioidosis clinically, especially in areas to which the disease is not endemic. In the series we report, 2 case-patients died before receiving definitive treatment, and only 1 case-patient received antimicrobial drugs recommended to treat melioidosis.

The integrated, population-based bacterial surveillance system in Kilifi County provides a unique opportunity to estimate incidence. Routine blood culture sampling of all admitted patients <15 years of age and eligible patients ≥15 years of age eliminates reliance on clinical suspicion for bacteremic melioidosis. The use of molecular methods on isolates suspected to be *B. pseudomallei* will probably enhance case detection because *B. pseudomallei* is commonly misidentified or unidentified by culture (9). Only 2 isolates in our study were identified by using standard techniques, despite the reported good discriminatory performance of API 20NE in distinguishing *B. pseudomallei* and *B. cepacia* (10).

Our reported incidence rates might still be underestimated. Our data do not account for KHDSS residents who do not go to KCH. For example, ≈64% of deaths in children <5 years of age in the KHDSS area occur at home or in other healthcare facilities (8). Furthermore, as

Table 1. Clinical summary of patients with positive *Burkholderia pseudomallei* isolates, Kilifi, Kenya, 2002–2014*

Year	Age/sex	Clinical features	Risk factors	Diagnosis†	Culture source	Antimicrobial sensitivity					Days in hospital	Outcome
						AMC	STX	TET	CAZ	IMI		
2002	8 d/M	Fever, jaundice, respiratory distress	None identified	Neonatal sepsis	Blood	S	S	S	S	S	3	Died
2008	7 d/M	Respiratory distress	None identified	Severe pneumonia, neonatal sepsis	Blood	S	S	S	S	S	3	Survived
2010	3 y/F	Fever, respiratory distress	Near-drowning	Severe pneumonia, septic shock	Blood	S	S	S	S	S	6	Died
2011	52 y/M	Persistent fever and night sweats of unknown duration	None identified	Acute renal failure, meningitis	Blood	S	S	S	S	S	5	Died
2014	68 y/F	Fever, bilateral cervical neck swellings	Diabetes mellitus	Diabetes, cervical lymphadenitis	Pus swab	S	S	S	S	S	40	Survived

*AMC, amoxicillin/clavulanic acid; CAZ, ceftazidime; IMI, imipenem; S, susceptible; STX, sulfamethoxazole/trimethoprim; TET, tetracycline.
†Diagnosis at time of admission.

demonstrated by the fifth case, the incidence of nonbacteremic infection might be higher because non-blood culture samples are not systematically collected. Only 50%–75% of patients with melioidosis are bacteremic (11), and culture has an estimated sensitivity of 60.2% for melioidosis (12). In addition, our screening method excluded gentamicin-susceptible isolates. If gentamicin-susceptible *B. pseudomallei* is as common in Kenya as reported in other areas (13), additional surveillance that includes these organisms could increase the reported incidence rates. Finally, melioidosis often is unevenly distributed within endemic areas, as noted in Thailand (14). Despite these factors, our results suggest that, although *B. pseudomallei* is present in tropical Africa, the incidence of invasive melioidosis is surprisingly low.

The differences in disease incidence in Africa and Asia are striking. Host factors, such as diabetes mellitus, might contribute, but environmental factors and agricultural practices, such as rice farming, are probably more important in permitting exposure to and environmental persistence and proliferation of the organism. Nonetheless, Kenya has been identified as environmentally suitable for *B. pseudomallei* because of its soil type, agricultural practices, and rainfall (1). Our study demonstrates the presence of *B. pseudomallei* in Kenya. Changes in climate and agricultural practices might lead to future increases in melioidosis, and ongoing surveillance is necessary.

Acknowledgments

We thank the nursing, clinical, and clerical staff of Kilifi County Hospital and the patients and their families.

The surveillance work was funded by the Wellcome Trust (core support to KEMRI–Wellcome Trust Research Program grant no. 203077) and by Gavi, the Vaccine Alliance, through support for a study of pneumococcal vaccine impact. J.A.G.S. is supported by a Wellcome Trust clinical research fellowship (no. 098532) and D.A.B.D. is funded by Wellcome grant no. 106698/Z/14/Z.

E.M.M. is supported through the DELTAS Africa Initiative [DEL-15-003]. The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences' (AAS) Alliance for Accelerating Excellence in Science in Africa and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust, [107769/Z/10/Z] and the government of the United Kingdom. The views expressed in this publication are those of the authors and not necessarily those of AAS, NEPAD Agency, Wellcome Trust, or the UK government.

This paper is published with the approval of the director of the Kenya Medical Research Institute.

About the Author

Dr. Muthumbi is a medical epidemiologist at the KEMRI–Wellcome Trust Research Programme in Kenya. She is a doctoral student in infectious disease epidemiology.

Table 2. Incidence of melioidosis in Kilifi County Hospital, Kilifi, Kenya, 1994–2012*

Patient age group	No. case-patients		Study period	Person-years of observation	Crude incidence† (95% CI)	Adjusted incidence† (95% CI)
	No. cases	residing in KHDSS area				
<15 y	3	2	1994–2012	2,026,781	1.0 (0.12–3.56)	1.3 (0.17–5.17)
≥15 y	1	1	2007–2012	782,373	1.3 (0.03–7.1)	2.0 (0.08–15.6)
Overall	4	3	NA	2,809,154	1.1 (0.22–3.12)	1.5 (0.35–5.0)


*KHDSS, Kilifi Health and Demographic Surveillance System; NA, not applicable.

†Incidence per 10⁶ person-years of observation, adjusted for nonconsenters and missing blood cultures among eligible consenters.

References

1. Limmathurotsakul D, Golding N, Dance DAB, Messina JP, Pigott DM, Moyes CL, et al. Predicted global distribution of *Burkholderia pseudomallei* and burden of melioidosis. *Nat Microbiol*. 2016;1:15008 <http://dx.doi.org/10.1038/nmicrobiol.2015.8>
2. Bremmelgaard A, Bygbjerg I, Høiby N. Microbiological and immunological studies in a case of human melioidosis diagnosed in Denmark. *Scand J Infect Dis*. 1982;14:271–5. <http://dx.doi.org/10.3109/inf.1982.14.issue-4.05>
3. Birnie E, Wiersinga WJ, Limmathurotsakul D, Grobusch MP. Melioidosis in Africa: should we be looking more closely? *Future Microbiol*. 2015;10:273–81. <http://dx.doi.org/10.2217/fmb.14.113>
4. Batchelor BI, Paul J, Trakulsomboon S, Mgongo M, Dance DAB. Melioidosis survey in Kenya. *Trans R Soc Trop Med Hyg*. 1994;88:181. [http://dx.doi.org/10.1016/0035-9203\(94\)90286-0](http://dx.doi.org/10.1016/0035-9203(94)90286-0)
5. Currie BJ, Dance DAB, Cheng AC. The global distribution of *Burkholderia pseudomallei* and melioidosis: an update. *Trans R Soc Trop Med Hyg*. 2008;102(Suppl 1):S1–4. [http://dx.doi.org/10.1016/s0035-9203\(08\)70002-6](http://dx.doi.org/10.1016/s0035-9203(08)70002-6)
6. Novak RT, Glass MB, Gee JE, Gal D, Mayo MJ, Currie BJ, et al. Development and evaluation of a real-time PCR assay targeting the type III secretion system of *Burkholderia pseudomallei*. *J Clin Microbiol*. 2006;44:85–90. <http://dx.doi.org/10.1128/jcm.44.1.85-90.2006>
7. Chewapreecha C, Holden MT, Vehkala M, Valimaki N, Yang Z, Harris SR, et al. Global and regional dissemination and evolution of *Burkholderia pseudomallei*. *Nat Microbiol*. 2017;2:16263. <http://dx.doi.org/10.1038/nmicrobiol.2016.263>
8. Scott JAG, Bauni E, Moisi JC, Ojal J, Gatakaa H, Nyundo C, et al. Profile: the Kilifi Health and Demographic Surveillance System (KHDSS). *Int J Epidemiol*. 2012;41(3):650–7. <http://dx.doi.org/10.1093/ije/dys062>
9. Amornchai P, Chierakul W, Wuthiekanun V, Mahakhunkijcharoen Y, Phetsouvanh R, Currie BJ, et al. Accuracy of *Burkholderia pseudomallei* identification using the API 20NE system and a latex agglutination test. *J Clin Microbiol*. 2007;45:3774–6. <http://dx.doi.org/10.1128/jcm.00935-07>
10. Kiratisin P, Santanirand P, Chantratita N, Kaewdaeng S. Accuracy of commercial systems for identification of *Burkholderia pseudomallei* versus *Burkholderia cepacia*. *Diagn Microbiol Infect Dis*. 2007;59:277–81. <http://dx.doi.org/10.1016/j.diagmicrobio.2007.06.013>
11. Wiersinga WJ, Virk HS, Torres AG, Currie BJ, Peacock SJ, Dance DAB, et al. Melioidosis. *Nat Rev Dis Primers*. 2018;4:17107. <http://dx.doi.org/10.1038/nrdp.2017.107>
12. Limmathurotsakul D, Jansen K, Arayawichanon A, Simpson JA, White LJ, Lee SJ, et al. Defining the true sensitivity of culture for the diagnosis of melioidosis using Bayesian latent class models. *PLoS ONE*. 2010;5:e12485. <http://dx.doi.org/10.1371/journal.pone.0012485>
13. Podin Y, Sarovich DS, Price EP, Kaestli M, Mayo M, Hii K, et al. *Burkholderia pseudomallei* isolates from Sarawak, Malaysian Borneo, are predominantly susceptible to aminoglycosides and macrolides. *Antimicrob Agents Chemother*. 2014;58(1):162–6. <https://dx.doi.org/10.1128/aac.01842-13>
14. Vuddhakul V, Tharavichitkul P, Na-Ngam N, Jitsurong S, Kunthawa B, Noimay P, et al. Epidemiology of *Burkholderia pseudomallei* in Thailand. *Am J Trop Med Hyg*. 1999;60:458–61. <http://dx.doi.org/10.4269/ajtmh.1999.60.458>

Address for correspondence: Esther M. Muthumbi, KEMRI–Wellcome Trust Research Programme, PO Box 230, Kilifi, Kenya; email: emuthumbi@kemri-wellcome.org



Originally published
in November 2009

https://wwwnc.cdc.gov/eid/article/15/11/e1-1511_article

etymologia revisited

Burkholderia

[burk'hol-dēr'e-ə]

This genus of gram-negative, rod-shaped bacteria comprising animal and plant pathogens was named for American plant pathologist Walter H. Burkholder. Dr. Burkholder first described a particular species of this genus, later called *Burkholderia cepacia* (Latin for “like onion”), after an outbreak of infection in vegetable growers in New York State in 1949. Previously known to cause disease in onion bulbs, these organisms are now recognized as major bacterial lung pathogens in patients with cystic fibrosis. *B. mallei* causes glanders in horses, and *B. pseudomallei* is the etiologic agent of melioidosis in humans and animals. Dr. Burkholder is recognized for helping establish the role of bacteria as plant pathogens.

Source: Dorland's illustrated medical dictionary, 31st edition. Philadelphia: Saunders; 2007; De Soya A, Silipo A, Lanzetta R, Govan JR, Molinaro A. Chemical and biological features of *Burkholderia cepacia* complex lipopolysaccharides. *Innate Immunity*. 2008;14:127.

Population-Based Estimate of Melioidosis, Kenya

Appendix

Case Summaries of Patients with Melioidosis, Kilifi, Kenya

Patient 1. November 2002. An 8-day old male was admitted with a history of fever for 4 days and yellowness of eyes for 3 days. Mother reported normal breastfeeding. On examination he was febrile (temperature 39.0°C) and tachypneic (respiratory rate 84 cycles/min). He was pale, capillary refill time was prolonged, he had jaundice, chest wall indrawing and nasal flaring. The hemogram showed leukocytosis comprising mainly neutrophils, blood slide for malaria parasites was negative. Blood culture showed gram-negative rods that were later identified as *B. pseudomallei*. There was no growth from the cerebral spinal fluid culture. The patient died after 3 days of treatment in the ward. Records of antibiotic treatment were not available from the retrieved case record, but first line treatment for this condition in 2002 was penicillin and gentamicin (1).

Patient 2. June 2008. A 7-day-old male child was first seen with fever and a history of difficulty breathing for 3 days. The child had been born at home and his mother reported normal breastfeeding. On examination he was in respiratory distress and had chest wall indrawing, and a diagnosis of neonatal sepsis was made. There was no growth from the cerebrospinal fluid culture. He was treated empirically with ampicillin and gentamicin, and by day 3 had improved sufficiently to be discharged. Subsequently, a gentamicin-resistant nonglucose-fermenting gram-negative rod was isolated on day 5, which could not be identified further at that time. The child was not a resident of the KHDSS district, and no further follow-up information was available.

Patient 3. May 2010. A 3-year-old female child was admitted with a history of cough, difficulty breathing, and fever after a near drowning accident in a local seasonal river. She had been found unconscious with a distended abdomen. She later regained full consciousness but was not brought for treatment until 2 days later, when she developed respiratory symptoms. At

admission, she had a temperature of 38.9°C and a respiratory rate of 50 cycles/min; she had nasal flaring and chest wall indrawing. The white cell count showed leukocytosis (18.2×10^9 cells/L) and neutrophilia (73%) on the differential count. Her chest radiograph had patchy opacities and oxygen saturation measured by oximetry was 96% on room air. She had features of severe pneumonia, and was initially treated with ampicillin and gentamicin, which was changed to ceftriaxone after 48 hours because there was no improvement in the clinical condition. On day 5, cloxacillin and metronidazole were added to the treatment. On day 5, admission blood cultures grew gram-negative rods, which were identified as *Burkholderia pseudomallei* (API 20NE profile 1156576), susceptible to ceftazidime but resistant to gentamicin. The patient died on day 6 before any further changes to the treatment were effected.

Patient 4. October 2011. A 52-year-old man was seen with an acute respiratory illness after persistent fever and night sweats. The duration of symptoms was not recorded in the patient records we retrieved. He was febrile (temperature 38.4°C) at admission, but no other constituent clinical signs were elicited. Blood counts and white cell differential counts also were within normal ranges. Sputum was negative for acid-fast bacilli on direct microscopy. He was initially treated with penicillin and gentamicin. A blood culture taken at admission became positive on day 5 for gram-negative rods that were susceptible to ceftazidime and imipenem. The final identity of the organism was not fully determined at the time. The patient deteriorated and died on day 5. We could not retrieve any results showing renal function tests or cerebrospinal fluid studies from the patient's records although the discharge diagnosis on record was meningitis with acute renal failure.

Patient 5. July 2014. A 68-year-old female patient, with insulin-dependent diabetes mellitus, was seen with bilateral cervical abscesses. She had attended outpatient clinics 2 weeks previously for investigation of cervical lymphadenopathy suspected to be tuberculous lymphadenitis. A fine needle aspirate had been performed, the results of which were not available on request. At admission, she had hyperglycemia and bilateral neck abscesses that were incised and drained. The complete blood count, liver and renal function tests were normal. The pus aspirate grew *B. pseudomallei* (API 20NE Profile 1156573); blood culture results were negative. Abdominal ultrasound done was normal. She was managed on intravenous imipenem for 10 days and discharged on co-trimoxazole for the eradication phase regimen to continue for 3

months. She did not return for the follow up clinic appointments. She was not a resident of the KHDSS and no further follow-up information was available.

Reference

1. World Health Organization. Clinical guidelines for diagnosis and treatment of common conditions in Kenya. Nairobi: Ministry of Health; 2002 [cited 2018 Nov 01].
<http://apps.who.int/medicinedocs/documents/s16427e/s16427e.pdf>