

Epidemiology of *Chlamydia trachomatis* in the Middle East and north Africa: a systematic review, meta-analysis, and meta-regression

Alex Smolak*, Hiam Chemaitelly*, Joumana G Hermez, Nicola Low, Laith J Abu-Raddad



Summary

Background The epidemiology of *Chlamydia trachomatis* in the Middle East and north Africa is poorly understood. We aimed to provide a comprehensive epidemiological assessment of *C trachomatis* infection in the Middle East and north Africa.

Methods We did a systematic review of *C trachomatis* infection as well as a meta-analysis and meta-regression of *C trachomatis* prevalence. We searched PubMed and Embase, as well as regional and national databases up to March 13, 2019, using broad search terms with no language or year restrictions. Any document or report including biological measures for *C trachomatis* prevalence or incidence was eligible for inclusion. We extracted all measures of current (genital or rectal), recent, and ever infection with *C trachomatis*. We estimated pooled average prevalence in different populations using random-effects meta-analysis. Factors associated with prevalence and sources of between-study heterogeneity were determined using meta-regression.

Findings We identified a total of 1531 citations, of which 255 reports contributed to 552 *C trachomatis* prevalence measures from 20 countries. No incidence measures were identified. Pooled prevalence of current genital infection was 3.0% (95% CI 2.3–3.8) in general populations, 2.8% (1.0–5.2) in intermediate-risk populations, 13.2% (7.2–20.7) in female sex workers, 11.3% (9.0–13.7) in infertility clinic attendees, 12.4% (7.9–17.7) in women with miscarriage, 12.4% (9.4–15.7) in symptomatic women, and 17.4% (12.5–22.8) in symptomatic men. Pooled prevalence of current rectal infection was 7.7% (4.2–12.0) in men who have sex with men. Substantial between-study heterogeneity was found. Multivariable meta-regression explained 29.0% of variation. Population type was most strongly associated with prevalence. Additional associations were found with assay type, sample size, country, and sex, but not with sampling methodology or response rate (about 90% of studies used convenience sampling and >75% had unclear response rate). There was no evidence for temporal variation in prevalence between 1982 and 2018.

Interpretation *C trachomatis* prevalence in the Middle East and north Africa is similar to other regions, but higher than expected given its sexually conservative norms. High prevalence in infertility clinic attendees and in women with miscarriage suggests a potential role for *C trachomatis* in poor reproductive health outcomes in this region.

Funding National Priorities Research Program from the Qatar National Research Fund (a member of Qatar Foundation).

Copyright © 2019 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

With more than 100 million incident infections every year,¹ *Chlamydia trachomatis* is one of the most common sexually transmitted infections (STIs) worldwide.^{2,3} Although curable, control and early detection of *C trachomatis* infection are challenged by its largely asymptomatic nature.⁴ Untreated *C trachomatis* infection is associated with serious reproductive tract conditions including pelvic inflammatory disease, ectopic pregnancy, infertility among women, and epididymitis among men.^{5,6}

Despite burdensome sequelae, STI control has long languished on health policy agendas. The 2030 Agenda on Sustainable Development⁷ aims to remedy this situation and led to WHO's Global Health Sector Strategy on STIs.⁸ The strategy proposes an integrated approach for STI prevention and control that addresses core Sustainable Development Goals, mainly through securing

universal access to sexual and reproductive health-care services and rights.^{7,8} The first strategic direction of this STI Strategy is “the need to understand the sexually transmitted infection epidemic and response as a basis for advocacy, political commitment, national planning, resource mobilization and allocation, implementation, and programme improvement.”⁸

The epidemiology of STIs, including *C trachomatis*, remains poorly understood in the Middle East and north Africa—a region comprising 10% of the world's population.^{9–11} Here, political and sociocultural sensitivities have set STIs low on countries' public health agendas, resulting in limited capacity for surveillance and programmes targeting sexual health, despite the possibility of a hidden disease burden.⁹ For example, the prevalence of primary infertility in the Middle East and north Africa, based on demographic and reproductive health surveys, has been

Lancet Glob Health 2019;

7: e1197–225

See [Comment](#) page e1152

*Contributed equally

Infectious Disease Epidemiology Group, Weill Cornell Medicine–Qatar, Cornell University, Qatar Foundation–Education City, Doha, Qatar (A Smolak PhD, H Chemaitelly MSc, Prof L J Abu-Raddad PhD); Department of Communicable Diseases, HIV/Hepatitis/STIs Unit, World Health Organization Regional Office for the Eastern Mediterranean, Cairo, Egypt (J G Hermez MPH); Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland (Prof N Low PhD); Department of Healthcare Policy & Research, Weill Cornell Medicine, Cornell University, New York, NY, USA (Prof L J Abu-Raddad); and College of Health and Life Sciences, Hamad bin Khalifa University, Doha, Qatar (Prof L J Abu-Raddad)

Correspondence to: Prof Laith J Abu-Raddad, Infectious Disease Epidemiology Group, Weill Cornell Medicine–Qatar, Cornell University, Qatar Foundation–Education City, PO Box 24144, Doha, Qatar lja2002@qatar-med.cornell.edu

Research in context

Evidence before this study

In a context of continuing stigma and political and sociocultural sensitivities, the Middle East and north Africa region has a dearth of epidemiological data about sexually transmitted infections. The prevalence of *Chlamydia trachomatis* and its distribution among populations at differing levels of risk of exposure remain largely unknown. A PubMed search using the search criteria (“Chlamydia”[MeSH] AND “Review” [Publication Type]) identified no systematic review and meta-analysis of regional scope for all subpopulations for this infection in the Middle East and north Africa or elsewhere.

Added value of this study

Using rigorous state-of-the-art methodologies with current empirical evidence, this study provided the first comprehensive epidemiological assessment of *C trachomatis* infection in the Middle East and north Africa. The study searched diverse sources of data, beyond international electronic databases, and identified a large volume of published and unpublished data, some of which now appears in the literature for the first time. The scope of evidence allowed analyses that found revealing associations relevant for the Middle East and north Africa and elsewhere. Unexpectedly, given this region’s sexually

conservative norms, the study estimated a *C trachomatis* prevalence of 3% in the population at large, similar to estimates from other regions. The study also documented high *C trachomatis* prevalence levels in infertility clinic attendees and in women with miscarriage, with odds of infection three-times higher than in the general population.

Implications of all the available evidence

There is a substantial *C trachomatis* infection and disease burden in the Middle East and north Africa that is neglected and poorly recognised despite its social and economic toll in a region comprising 10% of the world’s population. *C trachomatis* infection appears to be consistently associated with infertility and poor reproductive health outcomes in this region, yet these conditions are not linked to the possibility of an underlying infectious cause. The Middle East and north Africa is far from achieving WHO’s Global Health Sector Strategy on Sexually Transmitted Infections, 2016–21. The findings of this study provide a scientific foundation to develop an evidence-informed public health response against *C trachomatis* and its burdensome sequelae. The challenge will be to implement effective targeted, culturally appropriate, and gender-specific programmes to tackle *C trachomatis* infection and improve sexual health in general.

estimated to be the highest worldwide (although that of secondary infertility seems to be the lowest).¹² Still, the contribution of *C trachomatis*, or other STIs, to this disease burden remains unknown. Against this background, our study aimed to characterise *C trachomatis* epidemiology in the Middle East and north Africa.

Methods

Search strategy and selection criteria

We did a systematic review as well as a meta-analysis and meta-regression. We followed systematic review methods proposed by the Cochrane Collaboration,¹³ and report findings following the Preferred Reporting Items for Systematic Reviews and Meta-analyses guidelines¹⁴ (appendix pp 5–6). We did exhaustive searches using PubMed and Embase, regional and national databases (WHO Index Medicus for the Eastern Mediterranean region, Iraqi Academic Scientific Journals database, and Iranian Scientific Information Database), abstract archives of International AIDS Society Conferences,¹⁵ as well as country-level and international organisations’ reports available through the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database.^{10,11}

Our searches were done up to March 13, 2019, using broad search terms (MeSH/Emtree terms exploded to cover all subheadings and free-text terms) with no language or year restrictions. The appendix (p 7) summarises the search criteria and search terms used. The Middle East and north Africa were defined as 23 countries extending from Morocco in the west to Pakistan in the east (appendix p 8).

This definition for the Middle East and north Africa follows earlier convention applied in HIV and hepatitis C research,^{10,16–22} and is based on definitions by WHO, UNAIDS, and the World Bank.

Search results were checked for duplicates using Endnote (version 8.2). We screened titles and abstracts of unique citations. Full texts of citations deemed relevant or potentially relevant were retrieved for further screening by AS and HC. Any document or report including biological measures for *C trachomatis* prevalence or incidence, or both, based on primary data was eligible for inclusion. Case reports, case series, editorials, commentaries, reviews, and reports about military personnel stationed in the Middle East and north Africa, but not from these countries, were excluded. Reference lists of literature reviews and all relevant articles were hand-searched for additional eligible reports.

In this Article, the term report refers to a document (article, conference abstract, or country-level report) containing outcome measures of interest (ie, prevalence or incidence) for one or more populations, and the term study refers to details of a specific outcome measure in a specific population. Consequently, one report could contribute multiple studies and one study could be published in different reports. Duplicate study results were included only once using the most detailed report.

Data analysis

Data from relevant reports were extracted by AS with input from LJA-R. Independent extraction was done by

See Online for appendix

HC, and discrepancies were settled by consensus, or by contacting authors. Data from non-English articles were extracted from the full text by native speakers.

We extracted all measures of current (genital or rectal), recent, and ever infection with *C trachomatis*. We stratified data according to the study population's risk of exposure to *C trachomatis* or clinical manifestations (panel). Populations were defined as per original study authors' specific population definition and inclusion criteria (such as for infertile populations or women with miscarriage). We classified women and men as symptomatic only if there was an indication for the presence of *C trachomatis*-related signs and symptoms. We subsequently synthesised data by type of assay used for *C trachomatis* detection and summarised these data using medians and ranges.

Studies applying the same assay to different biological specimens were included only once, based on a sequential order that prioritised, for women, *C trachomatis* detection in endocervical swabs, followed by vaginal and urine samples; and for men, detection in urethral swabs, followed by urine and semen samples. Studies applying nucleic acid amplification test (NAAT) and culture to the same biological specimen were included separately given our interest in studying their contribution to heterogeneity in *C trachomatis* prevalence, and in generating STI-estimation correction factors based on assay type.^{23–25} Studies applying other antigen detection assays to the same biological specimen were included only once based on assay sensitivity (direct fluorescence and enzyme-linked immunoassays on genital samples were prioritised over Giemsa staining).

We excluded studies using tissue specimens from the upper genital tract, or including less than ten participants. We stratified the analyses by sex where relevant. Studies reporting only an overall measure for men and women were classified according to the predominant sex in the sample.

We did risk of bias and precision assessments. Informed by the Cochrane approach,¹³ we classified studies as having low versus high risk of bias for each of three quality domains assessing rigour of sampling methodology (probability based *vs* non-probability based), type of *C trachomatis* ascertainment (biological assay *vs* other, such as self-report), and response rate ($\geq 80\%$ response rate or $\geq 80\%$ of target sample size reached [the latter for studies using respondent-driven sampling] *vs* $< 80\%$). Studies with unavailable information about any given domain were classified as having unclear risk of bias for that domain. Studies were considered of higher precision if 200 participants or more underwent testing for *C trachomatis*, which was judged as an acceptable level of precision assuming a mean prevalence of 3% in the general population.

We produced forest plots to visualise estimates of prevalence and 95% CIs for each at-risk population, stratified by type of assay. Pooled average prevalence and 95% CIs were then estimated using meta-analysis for each stratum. A Freeman-Tukey type arcsine square-root

Panel: Definitions for at-risk population classification

General populations (populations at low risk)

Populations at low risk of exposure to *Chlamydia trachomatis* infection such as antenatal clinic attendees, blood donors, and pregnant women.

Populations at intermediate risk

Populations who presumably might have some sexual contacts with populations engaging in high sexual risk behaviour, and have therefore a higher risk of exposure to *C trachomatis* than the general population. These populations comprise prisoners, people who inject drugs, truck drivers, migrant workers, and HIV-infected individuals in a setting where the HIV epidemic is driven by injecting drug use.

Populations at high risk

Populations at high risk of exposure to *C trachomatis* as a consequence of specific high sexual risk behaviours such as female sex workers, men who have sex with men, male sex workers, men-to-women transgenders, and HIV-infected individuals in a setting where the HIV epidemic is driven by sexual transmission.

Infertility clinic attendees

Infertile women or men and their partners were included in a separate category given the potential biological link between *C trachomatis* infection and infertility.

Women with miscarriage

These women were included in a separate category given the potential biological link between *C trachomatis* infection and miscarriage.

Women with ectopic pregnancy

These women were included in a separate category given the potential biological link between *C trachomatis* infection and ectopic pregnancy.

Symptomatic women

Women with clinical manifestations related to *C trachomatis* infection, or suspected of having a *C trachomatis* infection such as those with vaginal discharge.

Symptomatic men

Men with clinical manifestations related to *C trachomatis* infection, or suspected of having a *C trachomatis* infection such as those with urethral discharge.

transformation was first applied to stabilise variances of prevalence measures.^{26,27} Measures were then weighted using the inverse-variance method,^{27,28} before being pooled using a DerSimonian-Laird random-effects model.²⁹ This model assumes a normal distribution for true effect sizes (ie, prevalence) across studies, which factors in sampling variation and true between-study heterogeneity.³⁰

We did heterogeneity assessment using Cochran's *Q* statistic to confirm existence of heterogeneity across studies, *I*² to quantify magnitude of between-study variation that is due to true differences in effect size rather than chance, and prediction interval to estimate the 95% CI of the distribution of true effect sizes.^{30,31} We did subgroup meta-analyses whenever five studies or more were available, using the R software (version 3.4.2).³²

We did random-effects meta-regression analyses to identify sources of between-study heterogeneity and estimated the magnitude of their association with prevalence. We included risk of bias and precision domains in the meta-regression analyses. We considered the




Figure 1: Flow chart of the study selection process for the systematic review

The systematic review focused on *Chlamydia trachomatis* incidence or prevalence, or both, in the Middle East and north Africa. *Reasons for exclusion: 14 studies applied the same assay to different biological specimens, and prevalence was included only once based on a sequential order prioritising for women *C trachomatis* detection in endocervical swabs followed by vaginal and urine samples, and for men detection in urethral swabs followed by urine and semen samples; one study applied antigen detection assays (other than nucleic acid amplification assays and culture) to the same biological specimen, and prevalence was included only once based on assay sensitivity; two studies were conducted only on subsamples of an original sample, and the prevalence in the original sample was included in the systematic review; six studies and three reports used tissue specimens from the upper genital tract; and one study had a sample size of less than 10.

following predictors a priori: at-risk population (panel), assay type (NAAT, culture, other assays detecting current infection, serological assays detecting anti-*C trachomatis* immunoglobulins of class IgG, IgM, IgA, immunoglobulins not specified, and unclear), sampling methodology (non-probability-based sampling vs probability-based sampling), sample size (<200 vs ≥200 participants), response rate (≥80% vs <80% and unclear), year of publication, year of data collection, country (Egypt, Iran,

Pakistan, and remaining countries; Egypt, Iran, and Pakistan being the most populous in the Middle East and north Africa),³³ and sex (women vs men; men-to-women transgenders who were biologically males were considered as men).

Studies that assessed *C trachomatis* prevalence using different diagnostics or biomarkers were included independently. Missing values for year of data collection were imputed using data for year of publication adjusted

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
Kadi et al (1990) ³⁵	Algeria	Cross-sectional	Convenience	69	W	Gynaecology clinic	Gynaecology clinic attendees	Serum	MIF (IgG)	17.4%
Kadi et al (1990) ³⁵	Algeria	Cross-sectional	Simple random sampling	180	W	Hospital	Women seeking rubella tests	Serum	MIF (IgG)	26.6%
Abdel Monem et al (2005) ³⁶	Egypt	Case-control	Convenience	20	W	Antenatal clinic	Pregnant women	Endocervical	Culture	15%
Aboul Atta and Ibrahim (1995) ³⁷	Egypt	Case-control	Convenience	20	M	Hospital	Controls in STI study	Urethral	DFA	5%
Badary (1996) ³⁸	Egypt	Case-control	Convenience	32	W	Gynaecology clinic	Fertile women	Endocervical	DFA	12.5%
Berry and El Shabrawy (1996) ³⁹	Egypt	Case-control	Convenience	30	W	Family planning clinic	Family planning clinic attendees	Serum	EIA (IgG)	3.3%
Diab (1993) ⁴⁰	Egypt	Case-control	Convenience	30	W	Antenatal clinic	Women with full-term delivery	Serum	EIA (IgG)	0
Draz et al (2018) ⁴¹	Egypt	Case-control	Convenience	14	W	Gynaecology clinic	Healthy women	Endocervical	DFA	0
El-Sayed et al (2002) ⁴²	Egypt	Cross-sectional	Convenience	108	W	Family planning clinic	Family planning clinic attendees	Urine	NAAT	2.8%
El-Sayed et al (2002) ⁴²	Egypt	Cross-sectional	Convenience	604	W	Antenatal clinic	Antenatal clinic attendees	Urine	NAAT	1.3%
Mosbah and Nabel (2016) ⁴³	Egypt	Case-control	Convenience	90	W	Hospital	Pregnant women with pre-eclampsia	Endocervical	NAAT	4.4%
Mosbah and Nabel (2016) ⁴³	Egypt	Case-control	Convenience	90	W	Hospital	Normotensive pregnant women	Endocervical	NAAT	0
Mousa (1990) ⁴⁴	Egypt	Cross-sectional	Convenience	50	W	Gynaecology clinic	Gynaecology clinic attendees	Endocervical	DFA	2%
Nada et al (2015) ⁴⁵	Egypt	Case-control	Convenience	100	W	Gynaecology clinic	Gynaecology clinic attendees	Endocervical	NAAT	2%
Sullam et al (2001) ⁴⁶	Egypt	Cross-sectional	Multistage cluster sampling	1344	W	Community	Household survey of women	Endocervical	ELISA	4.2%
Zaki (1989) ⁴⁷	Egypt	Cross-sectional	Convenience	100	W	Antenatal clinic	Pregnant women	Endocervical	Culture	3%
Ahmadi et al (2016a) ⁴⁸	Iran	Case-control	Convenience	109	W	Family planning clinic	Family planning clinic attendees	Endocervical	NAAT	11.9%
Ahmadi et al (2018) ⁴⁹	Iran	Case-control	Convenience	165	M	Clinic	Fertile men	Semen	NAAT	0.6%
Ahmadnia et al (2016) ⁵⁰	Iran	Cross-sectional	Stratified cluster sampling	4274	W	Primary health-care centre	Primary health-care centre clinic attendees	Endocervical	Culture	1%
Badami and Salari (2001) ⁵¹	Iran	Cross-sectional	Convenience	250	W	Family planning clinic	Family planning clinic attendees	Serum	DFA	0.8%
Badami and Salari (2001) ⁵¹	Iran	Cross-sectional	Convenience	250	W	Family planning clinic	Family planning clinic attendees	Serum	Unclear	3.2%
Baghchesaraei et al (2011) ⁵²	Iran	Cross-sectional	Convenience	328	W	Gynaecology clinic	Gynaecology clinic attendees	Serum	EIA (IgM)	10.3%
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	60	W	Fertility centre	Pregnant women	Vaginal	NAAT	0
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	60	W	Fertility centre	Pregnant women	Serum	ELISA (IgA)	6.7%
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	60	W	Fertility centre	Pregnant women	Serum	ELISA (IgG)	1.7%
Behroozi (2001) ⁵⁴	Iran	Cross-sectional	Convenience	400	W	Antenatal clinic	Pregnant women	Unclear	DFA	2.8%
Chamani-Tabriz et al (2008) ⁵⁵	Iran	Cross-sectional	Convenience	991	W	Community	Married women	Urine	NAAT	12.8%
Cheraghi et al (2014) ⁵⁶	Iran	Cross-sectional	Convenience	1448	W	Health centres	Non-pregnant women	Endocervical	Unclear	0.2%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Antenatal clinic	Antenatal clinic attendees	Urine	NAAT	0
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Antenatal clinic	Antenatal clinic attendees	Serum	EIA (IgA)	0
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Antenatal clinic	Antenatal clinic attendees	Serum	EIA (IgM)	0
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Antenatal clinic	Antenatal clinic attendees	Serum	EIA (IgG)	12.8%
Goshayeshi et al (2015) ⁵⁸	Iran	Case-control	Convenience	30	W	Fertility centre	Fertile women	Endocervical	NAAT	3.3%
Haghighi Hasanabad et al (2013) ⁵⁹	Iran	Cross-sectional	Convenience	399	W	Antenatal clinic	Pregnant adolescents	Unclear	NAAT	12.3%
Jahromi et al (2010) ⁶⁰	Iran	Case-control	Convenience	200	W	Gynaecology clinic	Women with full-term delivery	Endocervical	DFA	5.2%
Javanmard et al (2018) ⁶¹	Iran	Cross-sectional	Convenience	210	W	Gynaecology clinic	Women undergoing routine pap smear	Endocervical	NAAT	11.4%

(Table 1 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†	
(Continued from previous page)											
	Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	125	W	Hospital	Pregnant women	Vaginal	NAAT	1.6%
	Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	125	W	Hospital	Pregnant women	Serum	ELISA (IgM)	1.6%
	Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	125	W	Hospital	Pregnant women	Serum	ELISA (IgG)	0
	Kajbaf and Gholamnezhad (1998) ⁶³	Iran	Case-control	Convenience	50	W	Antenatal clinic	Antenatal clinic attendees	Endocervical	DFA	4%
	Kajbaf and Gholamnezhad (1998) ⁶³	Iran	Case-control	Convenience	50	W	Antenatal clinic	Antenatal clinic attendees	Serum	ELISA (IgG)	6%
	Kamyabi (2009) ⁶⁴	Iran	Case-control	Convenience	35	W	Gynaecology clinic	Pregnant women	Serum	ELISA (IgG)	20%
	Khezerdoust et al (2009) ⁶⁵	Iran	Cross-sectional	Convenience	1114	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgG)	2.9%
	Marashi et al (2014) ⁶⁶	Iran	Case-control	Convenience	200	W	Antenatal clinic	Pregnant women	Endocervical	DFA	3.5%
	Marashi et al (2014) ⁶⁶	Iran	Case-control	Convenience	200	W	Antenatal clinic	Pregnant women	Endocervical	NAAT	8.7%
	Ministry of Health and Medical Education (2008) ⁶⁷	Iran	Case-control	Convenience	41	W	Clinic	Fertile women	Serum	ELISA (IgG)	2.4%
	Meidani (2009) ⁶⁸	Iran	Cross-sectional	Simple random sampling	140	M	Laboratory	Men—premarital or pre-employment screening	Urine	NAAT	0.7%
	Ministry of Health and Medical Education (2008) ⁶⁷	Iran	Cross-sectional	Convenience	70	W	Antenatal clinic	Pregnant women	Serum	ELISA (unclear)	4.3%
	Ministry of Health and Medical Education (2008) ⁶⁷	Iran	Case-control	Convenience	250	W	Family planning clinic	Healthy women	Endocervical	DFA	0.8%
	Mousavi et al (2014) ⁶⁹	Iran	Case-control	Convenience	104	W	Antenatal clinic	Antenatal clinic attendees	Endocervical	NAAT	5.8%
	Pourabbas et al (2018) ⁷⁰	Iran	Cross-sectional	Convenience	239	W	Hospital	Pregnant women	Endocervical	NAAT	15.5%
	Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	223	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgM)	1.8%
	Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	223	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgG)	5.0%
	Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	223	W	Antenatal clinic	Pregnant women	Urine	NAAT	8.5%
	Rohi et al (2011) ⁷²	Iran	Cross-sectional	Convenience	91	W	Hospital	Pregnant women	Serum	ELISA (IgG)	28.6%
	Rostami et al (2016) ⁷³	Iran	Cross-sectional	Simple random sampling	518	W	Gynaecology clinic	Gynaecology clinic attendees	Endocervical	NAAT	7.1%
	Safdari et al (2015) ⁷⁴	Iran	Cross-sectional	Convenience	70	W	Antenatal clinic	Antenatal clinic attendees	Endocervical	NAAT	10%
	Safdari et al (2015) ⁷⁴	Iran	Cross-sectional	Convenience	70	W	Antenatal clinic	Antenatal clinic attendees	Endocervical	Culture	8.6%
	Sattari et al (2017) ⁷⁵	Iran	Case-control	Convenience	100	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgM)	2%
	Sattari et al (2017) ⁷⁵	Iran	Case-control	Convenience	100	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgG)	18%
	Sisakht et al (2017) ⁷⁶	Iran	Case-control	Convenience	30	W	Gynaecology clinic	Women with full-term delivery	Urine	NAAT	4.7%
	Yeganeh et al (2013) ⁷⁷	Iran	Case-control	Convenience	100	M	Urology clinic	Asymptomatic men	Urine	NAAT	4%
	Zahirmia et al (2018) ⁷⁸	Iran	Cross-sectional	Convenience	76	W	Gynaecology clinic	Pregnant women	Vaginal	NAAT	10.5%
	Abdulkhudher et al (2014) ⁷⁹	Iraq	Case-control	Convenience	40	W	Antenatal clinic	Women with full-term delivery	Serum	ELISA (IgM)	0
	Abdulkhudher et al (2014) ⁷⁹	Iraq	Case-control	Convenience	40	W	Antenatal clinic	Women with full-term delivery	Serum	ELISA (IgG)	7.5%
	Abdul-Karim et al (2009) ⁸⁰	Iraq	Cross-sectional	Convenience	198	W	Hospital	Women with full-term delivery	Serum	ELISA (IgG)	13.7%
	Abdullah (2012) ⁸¹	Iraq	Case-control	Convenience	24	W	Hospital	Pregnant women	Serum	ELISA (IgM)	0
	Abdullah (2012) ⁸¹	Iraq	Case-control	Convenience	24	W	Hospital	Pregnant women	Serum	ELISA (IgG)	8.3%
	Ahmed (2008) ⁸²	Iraq	Case-control	Convenience	30	W	Hospital	Women with full-term delivery	Serum	ELISA (unclear)	0
	Al-Hamdani et al (2010) ⁸³	Iraq	Case-control	Convenience	17	W	Hospital	Pregnant women	Serum	ELISA (IgM)	14.0%

(Table 1 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†	
(Continued from previous page)											
	Al-Hamdani et al (2010) ⁸³	Iraq	Case-control	Convenience	17	W	Hospital	Pregnant women	Serum	ELISA (IgG)	14.0%
	Al-Hamdani et al (2010) ⁸³	Iraq	Case-control	Convenience	17	W	Hospital	Pregnant women	Serum	ELISA (IgA)	40.4%
	Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	100	W	Family planning clinic	Family planning clinic attendees	Serum	IFAT (unclear)	5%
	Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	100	W	Family planning clinic	Family planning clinic attendees	Endocervical	ELFA	4%
	Ali and Al-Kazaz (2018) ⁸⁵	Iraq	Case-control	Convenience	13	M	Clinic	Fertile men	Semen	NAAT	0
	Alkhafaf (2013) ⁸⁶	Iraq	Case-control	Convenience	122	W	Hospital	Married women	Serum	ELISA (IgG)	4.1%
	Alkhafaf (2013) ⁸⁶	Iraq	Case-control	Convenience	168	W	Hospital	Unmarried woman	Serum	ELISA (IgG)	2.4%
	Hwaid et al (2013) ⁸⁷	Iraq	Case-control	Simple random sampling	91	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgG)	5.5%
	Hwaid et al (2013) ⁸⁷	Iraq	Case-control	Simple random sampling	91	W	Antenatal clinic	Pregnant women	Serum	ELISA (IgM)	3.3%
	Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	50	W	Laboratories	General population women	Serum	ELISA (IgM)	10%
	Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	50	W	Laboratories	General population women	Serum	ELISA (IgG)	4%
	Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	50	W	Laboratories	General population women	Serum	ELISA (IgA)	2%
	Mohammed et al (2012) ⁸⁹	Iraq	Case-control	Convenience	23	W	Gynaecology clinic	Pregnant women	Serum	ELISA (IgM)	0
	Mohammed et al (2017) ⁹⁰	Iraq	Case-control	Convenience	20	W	Gynaecology clinic	Gynaecology clinic attendees	Endocervical	NAAT	0
	Mohammed et al (2017) ⁹⁰	Iraq	Case-control	Convenience	20	W	Gynaecology clinic	Gynaecology clinic attendees	Serum	ELISA (IgG)	0
	Yahya and Al-Siraj (2009) ⁹¹	Iraq	Cross-sectional	Convenience	296	M	Laboratory	Fertile men	Serum	Culture	0
	Abusarah et al (2013) ⁹²	Jordan	Case-control	Convenience	61	M	Urology clinics	Fertile men	Urine	NAAT	1.6%
	Al-Ramahi et al (2008) ⁹³	Jordan	Case-control	Convenience	146	W	Gynaecology clinic	Gynaecology clinic attendees	Endocervical	NAAT	0.7%
	As'ad (2004) ⁹⁴	Jordan	Cross-sectional	Convenience	144	W	Family planning clinic	Asymptomatic women	Vaginal	NAAT	0
	Awwad et al (2003) ⁹⁵	Jordan	Case-control	Convenience	61	M	Urology clinic	Non-urethritis patients	Urine	NAAT	0
	Awwad et al (2003) ⁹⁵	Jordan	Case-control	Convenience	39	W	Urology clinic	Non-urethritis patients	Urine	NAAT	0
	Mahafzah et al (2008) ⁹⁶	Jordan	Cross-sectional	Convenience	186	W	Gynaecology clinic	Family planning clinic attendees	Endocervical	NAAT	0.5%
	Jordan Ministry of Health (2004) ⁹⁷	Jordan	Cross-sectional	Convenience	213	W	Hospital	Asymptomatic women	Endocervical	NAAT	0.5%
	Al-Awadhi et al (2018) ⁹⁸	Kuwait	Cross-sectional	Convenience	65 338	W	Laboratory	Women undergoing pap smear 1997–2005	Endocervical	Unclear	0.1%
	Al-Awadhi et al (2018) ⁹⁸	Kuwait	Cross-sectional	Convenience	56 105	W	Laboratory	Women undergoing pap smear 2006–14	Endocervical	Unclear	0.04%
	Al-Sweih et al (2011) ⁹⁹	Kuwait	Cross-sectional	Convenience	5938	W	Primary health-care centre	Kuwaiti women	Vaginal	NAAT	1.9%
	Al-Sweih et al (2011) ⁹⁹	Kuwait	Cross-sectional	Convenience	2601	W	Primary health-care centre	Expatriate women	Vaginal	NAAT	2.3%
	Al-Sweih et al (2012) ¹⁰⁰	Kuwait	Case-control	Convenience	188	M	Gynaecology clinic	Fertile men	Semen	NAAT	3.7%
	Deeb et al (2003) ¹⁰¹	Lebanon	Cross-sectional	Multistage random sampling	506	W	Community	Ever-married women	Endocervical	ELISA	0
	Hancali et al (2015) ¹⁰²	Morocco	Cross-sectional	Convenience	760	W	Family planning clinic	Family planning clinic attendees in 1999	Unclear	NAAT	4.0%
	Hancali et al (2015) ¹⁰²	Morocco	Cross-sectional	Convenience	256	W	Family planning clinic	Family planning clinic attendees in 2011	Unclear	NAAT	4.4%

(Table 1 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†	
(Continued from previous page)											
	Hulstein et al (2018) ¹⁰³	Morocco	Cross-sectional	Simple random sampling	163	M	Community	General population men	Serum	IFAT (IgG)	31.3%
	Hulstein et al (2018) ¹⁰³	Morocco	Cross-sectional	Simple random sampling	174	W	Community	General population women	Serum	IFAT (IgG)	37.9%
	Morocco Ministry of Health (2001) ¹⁰⁴	Morocco	Cross-sectional	Convenience	323	W	Antenatal clinic	Pregnant women	Urine	NAAT	2.7%
	Morocco Ministry of Health (2001) ¹⁰⁴	Morocco	Cross-sectional	Convenience	518	W	Family planning clinic	Family planning clinic attendees	Urine	NAAT	5.2%
	The Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project (2017) ¹⁰⁵	Morocco	Cross-sectional	Convenience	252	W	Antenatal clinic	Pregnant women	Unclear	NAAT	3.6%
	The Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project (2017) ¹⁰⁵	Morocco	Cross-sectional	Convenience	537	W	Family planning clinic	Family planning clinic attendees	Unclear	NAAT	3%
	Radouani et al (1998) ¹⁰⁶	Morocco	Case-control	Convenience	81	W	Hospital	Pregnant women	Serum	MIF (unclear)	14.8%
	Radouani et al (1998) ¹⁰⁶	Morocco	Case-control	Convenience	200	M	Hospital	Blood donors	Serum	MIF (unclear)	4.5%
	Takourt et al (1995) ¹⁰⁷	Morocco	Case-control	Convenience	200	M	Hospital	Blood donors	Serum	MIF (unclear)	5.0%
	Takourt et al (1995) ¹⁰⁷	Morocco	Case-control	Convenience	200	W	Hospital	Blood donors	Serum	MIF (unclear)	10.0%
	Mir et al (2009) ¹⁰⁸	Pakistan	Cross-sectional	Multistage systematic random sampling	2383	M	Community	General population men	Urine	NAAT	0
	Wasti et al (1997) ¹⁰⁹	Pakistan	Cross-sectional	Convenience	300	W	Antenatal clinic and family planning clinic	Antenatal clinic and family planning clinic attendees	Endocervical	DFA	5.3%
	Al-Thani et al (2013) ¹¹⁰	Qatar	Cross-sectional	Convenience	133	W	Primary health-care centre	Qatari women	Endocervical	NAAT	5.3%
	Al-Thani et al (2013) ¹¹⁰	Qatar	Cross-sectional	Convenience	218	W	Primary health-care centre	Non-Qatari women	Endocervical	NAAT	5.5%
	Alzahrani et al (2010) ¹¹¹	Saudi Arabia	Cross-sectional	Simple random sampling	95	W	Antenatal clinic	Pregnant women	Endocervical	ELISA	10.5%
	Awad et al (2013) ¹¹²	Saudi Arabia	Cross-sectional	Convenience	144	W	Gynaecology clinic	Antenatal clinic attendees	Urine	NAAT	11.1%
	Bashi (1987) ¹¹³	Saudi Arabia	Cross-sectional	Convenience	100	W	Primary health-care centre	Primary health-care centre attendees	Serum	MIF (IgG)	0
	Bashi (1987) ¹¹³	Saudi Arabia	Cross-sectional	Convenience	100	M	Primary health-care centre	Primary health-care centre attendees	Serum	MIF (IgG)	2%
	Ghazi et al (2006) ¹¹⁴	Saudi Arabia	Cross-sectional	Simple random sampling	1600	W	Antenatal clinic	Saudi pregnant women	Serum	ELISA (IgG)	8.7%
	Ghazi et al (2006) ¹¹⁴	Saudi Arabia	Cross-sectional	Simple random sampling	1460	W	Antenatal clinic	Saudi pregnant women	Serum	ELISA (IgM)	1.5%
	Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	112	M	Hospital	Blood donors	Serum	MIF (IgM)	0
	Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	112	M	Hospital	Blood donors	Serum	MIF (IgG)	1.8%
	Kamel (2013) ¹¹⁶	Saudi Arabia	Randomised controlled trial‡	Convenience	100	W	Antenatal clinic	Antenatal clinic attendees	Serum	ELISA (IgG)	4.0%
	Massoud et al (1991) ¹¹⁷	Saudi Arabia	Case-control	Convenience	100	W	Hospital	Asymptomatic women	Serum	IFAT (unclear)	0
	Massoud et al (1991) ¹¹⁷	Saudi Arabia	Case-control	Convenience	100	M	Hospital	Asymptomatic men	Serum	IFAT (unclear)	2.0%
	Ismail et al (1990) ¹¹⁸	Somalia	Cross-sectional	Convenience	194	W	Community	Women	Endocervical	EIA	12.4%

(Table 1 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†	
(Continued from previous page)											
	Ismail et al (1990) ¹¹⁸	Somalia	Cross-sectional	Convenience	189	M	Community	Men	Urethral	EIA	6%
	Nur et al (2000) ¹¹⁹	Somalia	Cross-sectional	Convenience	54	M	Hospital	Blood donors	Serum	EIA (IgG)	22.2%
	WHO (2005a) ¹²⁰	Somalia	Cross-sectional	Convenience	4723	W	Antenatal clinic	Pregnant women	Urine	NAAT	1.7%
	WHO (2005b) ¹²¹	Somalia	Cross-sectional	Convenience	509	W	Antenatal clinic	Pregnant women	Urine	NAAT	1.4%
	Ahmed et al (2018) ¹²²	Sudan	Case-control	Convenience	93	W	Hospital	Healthy pregnant women	Serum	ELISA (IgG)	0
	Ahmed et al (2018) ¹²²	Sudan	Case-control	Convenience	93	W	Hospital	Pregnant women with pre-eclampsia	Serum	ELISA (IgG)	0
	Almroth et al (2005) ¹²³	Sudan	Case-control	Convenience	139	W	Antenatal clinic	Antenatal clinic attendees	Serum	EIA (IgG)	3.6%
	Ortashi et al (2004) ¹²⁴	Sudan	Cross-sectional	Convenience	151	W	Antenatal clinic	Pregnant women	Endocervical and urethral	EIA	19.9%
	Alkayer et al (2017) ¹²⁵	Syria	Case-control	Convenience	21	W	Hospital	Pregnant women	Serum	ELISA (IgG)	4.7%
	Ghazal-Aswad et al (2006) ¹²⁶	United Arab Emirates	Cross-sectional	Multistage cluster sampling	727	W	Clinics	Primary health-care centre and clinic attendees	Endocervical	EIA	2.5%

DFA=direct fluorescent assay. EIA=enzyme immunoassay. ELFA=enzyme-linked fluorescence assay. IFAT=indirect fluorescent antibody test. M=men or sample predominantly of men. MIF=micro-immunofluorescence. NAAT=nucleic acid amplification test. STI=sexually transmitted infection. W=women or sample predominantly of women. *Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).^{127,128} Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.¹²⁸ Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.¹²⁹ †The decimal places of the prevalence measures are as reported in the original report, but prevalence figures with more than one decimal place were rounded to one decimal place. ‡The extracted prevalence measure is for the baseline measurement.

Table 1: Studies reporting *Chlamydia trachomatis* prevalence in general populations in the Middle East and north Africa

by the median difference between year of publication and year of data collection (for studies with complete information). We did meta-regression diagnostics.

Factors associated with prevalence at $p \leq 0.20$ in univariable analysis were eligible for inclusion in the multivariable analysis. In the multivariable model, a $p \leq 0.05$ for any given factor indicated strong evidence for an association with *C trachomatis* prevalence. We did meta-regression models using Stata/SE (version 14).³⁴

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the article. The corresponding author had full access to all the data in the study and had the final responsibility for the decision to submit for publication.

Results

The search identified a total of 1531 citations: 509 through PubMed, 557 through Embase, and 465 through regional and national databases. Of these citations, 302 reports underwent full-text screening after excluding duplicates and screening titles and abstracts. During full-text screening, 228 eligible reports were identified, and 74 were excluded for reasons outlined in figure 1. Hand-searching of reference lists of relevant reports and reviews yielded 14 additional eligible reports. 16 country-level reports were further identified through the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database. Three reports were subsequently excluded. In total, 255 reports contributing 552 prevalence measures

or studies met the eligibility criteria for inclusion, but no incidence measures were identified.

Evidence covered 20 (87%) of 23 countries, encompassing a total of 256769 *C trachomatis* test results (tables 1 and 2; appendix pp 9–14). Iran contributed the largest number of measures or studies ($n=176$), followed by Egypt ($n=89$), Iraq ($n=72$), Saudi Arabia ($n=45$), Pakistan ($n=42$), and Morocco ($n=32$). Most studies assessed current infection ($n=318$), whereas the rest reported different serological measures ($n=211$), such as ever infection (anti-*C trachomatis* IgG; $n=117$). Details of *C trachomatis* testing protocol were specified in 424 (77%) of 552 studies; 320 (75%) of the 424 used commercial assays, 62 (15%) used in-house validated tests, 29 (7%) used culture, and 13 (3%) used a non-validated in-house test.

In general populations ($n=137$), prevalence of current genital infection ranged from 0 to 19.9% with a median of 3.0%, whereas ever infection prevalence ranged from 0 to 37.9% with a median of 4.7% (tables 1 and 3).

In populations at high risk ($n=40$), current infection prevalence in female sex workers ($n=20$) ranged from 0.9% to 72.9% with a median of 8.4%, whereas ever infection prevalence ranged from 19.8% to 100% with a median of 90.0% (tables 2 and 3). In men who have sex with men (including male sex workers and male-to-female transgenders; $n=20$), current infection prevalence ranged from 0 to 8.8% with a median of 1.2% for genital infections and from 3.6% to 18.3% with a median of 6.3% for rectal infections, but no ever infection measure was identified.

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
Populations at high risk										
Kadi et al (1990) ³⁵	Algeria	Cross-sectional	Convenience	44	W	Community	Female sex workers	Serum	MIF (IgG)	100%
El-Sayed et al (2002) ⁴²	Egypt	Cross-sectional	Convenience	52	W	Community	Female sex workers	Urine	NAAT	7.7%
El-Sayed et al (2002) ⁴²	Egypt	Cross-sectional	Convenience	80	M	Community	Men who have sex with men	Urine	NAAT	8.8%
Darougar et al (1983) ³⁰	Iran	Cross-sectional	Convenience	116	W	Community	Female sex workers	Endocervical	Culture	6.9%
Darougar et al (1983) ³⁰	Iran	Cross-sectional	Convenience	154	W	Community	Female sex workers	Serum	MIF (IgM)	29.2%
Darougar et al (1983) ³⁰	Iran	Cross-sectional	Convenience	154	W	Community	Female sex workers	Serum	MIF (IgG)	94.2%
Kassaian et al (2012) ³¹	Iran	Cross-sectional	Convenience	91	W	Mixed	Female sex workers	Serum	ELISA (IgG)	19.8%
Kazerooni et al (2014) ³²	Iran	Cross-sectional	Respondent-driven sampling	278	W	Community	Female sex workers	Vaginal	NAAT	9%
Mirzazadeh et al (2016) ³³	Iran	Cross-sectional	Convenience	1337	W	Community	Female sex workers	Vaginal	NAAT	6%
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	30	W	STI clinic	Women with multiple partners	Endocervical	ELFA	30%
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	30	W	STI clinic	Women with multiple partners	Serum	IFAT (unclear)	36.7%
Bellaji et al (2017) ³⁴	Morocco	Cross-sectional	Convenience	519	W	NGOs	Female sex workers	Endocervical and vaginal	NAAT	20.7%
Morocco Ministry of Health (2008) ³⁵	Morocco	Cross-sectional	Convenience	141	W	STI clinic	Female sex workers	Endocervical and urine	NAAT	22.7%
Morocco Ministry of Health (2011) ³⁶	Morocco	Cross-sectional	Respondent-driven sampling	368	W	Community	Female sex workers in Agadir	Endocervical	NAAT	22.4%
Morocco Ministry of Health (2015) ³⁷	Morocco	Cross-sectional	Respondent-driven sampling	247	M	Community	Men who have sex with men in Agadir	Urine	NAAT	5.4%
Morocco Ministry of Health (2015) ³⁷	Morocco	Cross-sectional	Respondent-driven sampling	252	M	Community	Men who have sex with men in Marrakech	Urine	NAAT	6.5%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	426	W	Community	Female sex workers in Rawalpindi	Endocervical	NAAT	1.7%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	107	W	Community	Female sex workers in Abbottabad	Endocervical	NAAT	0.9%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	195	M	Community	Male sex workers in Rawalpindi (Bantha)	Urine	NAAT	0
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	195	M	Community	Male sex workers in Rawalpindi (Bantha)	Rectal	NAAT	4.7%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	364	M	Community	Male sex workers in Rawalpindi (Khotki)	Urine	NAAT	0
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	364	M	Community	Male sex workers in Rawalpindi (Khotki)	Rectal	NAAT	3.6%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	253	M	Community	Male sex workers in Rawalpindi (Khusra)	Urine	NAAT	0
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	253	M	Community	Male sex workers in Rawalpindi (Khusra)	Rectal	NAAT	9.9%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	83	M	Community	Male sex workers in Abbottabad (Bantha)	Urine	NAAT	1.2%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	83	M	Community	Male sex workers in Abbottabad (Bantha)	Rectal	NAAT	4.9%
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	20	M	Community	Male sex workers in Abbottabad (Khotki and Khusra)	Urine	NAAT	0
Hawkes et al (2009) ³⁸	Pakistan	Cross-sectional	Respondent-driven sampling	20	M	Community	Male sex workers in Abbottabad (Khotki and Khusra)	Rectal	NAAT	6.3%
Khan et al (2011) ³⁹	Pakistan	Cross-sectional	Respondent-driven sampling	730	W	Community	Female sex workers in Lahore	Endocervical	NAAT	7.7%
Osama (2017) ⁴⁰	Pakistan	Cross-sectional	Convenience	2531	M	Drop in centre	Men who have sex with men in Lahore	Unclear	Unclear	35.2%
Rehan et al (2009) ⁴¹	Pakistan	Cross-sectional	Systematic random sampling	383	W	Red-light	Female sex workers in Lahore	Endocervical	NAAT	11%
Rehan et al (2009) ⁴¹	Pakistan	Cross-sectional	Snowball	348	W	Community	Female sex workers in Karachi	Endocervical	NAAT	5.2%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Respondent-driven sampling	395	M	Community	Male sex workers in Lahore	Urethral	NAAT	1.5%
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Snowball	396	M	Community	Male sex workers in Karachi	Urethral	NAAT	1.2%
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Snowball	394	M	Community	Male sex workers in Karachi	Rectal	NAAT	10.4%
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Systematic random cluster sampling	198	M	Community	Hijras in Lahore	Urethral	NAAT	1.5%
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Systematic random cluster sampling	197	M	Community	Hijras in Karachi	Urethral	NAAT	0
Rehan et al (2009) ¹⁴¹	Pakistan	Cross-sectional	Systematic random cluster sampling	197	M	Community	Hijras in Karachi	Rectal	NAAT	18.3%
Znazen et al (2010) ¹⁴²	Tunisia	Cross-sectional	Convenience	188	W	Community	Female sex workers	Endocervical	NAAT	72.9%
Znazen et al (2010) ¹⁴²	Tunisia	Cross-sectional	Convenience	183	W	Community	Female sex workers	Serum	MIF (IgG)	85.8%
Infertility clinic attendees										
Abdel Aleem et al (1996) ¹⁴³	Egypt	Case-control	Convenience	144	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	52%
Abdel Aleem et al (1996) ¹⁴³	Egypt	Case-control	Convenience	104	M	Infertility clinic	Men with unclear infertility diagnosis	Serum	ELISA (IgG)	24%
Abdel Monem et al (2005) ³⁶	Egypt	Case-control	Convenience	150	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	Culture	24%
Abdel Monem et al (2005) ³⁶	Egypt	Case-control	Convenience	150	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	EIA	22.7%
Abdella et al (2015) ¹⁴⁴	Egypt	Case-control	Convenience	50	W	Infertility clinic	Women with idiopathic infertility	Serum	ELISA (IgM)	4%
Abdella et al (2015) ¹⁴⁴	Egypt	Case-control	Convenience	50	W	Infertility clinic	Women with idiopathic infertility	Serum	ELISA (IgG)	36%
Abdella et al (2015) ¹⁴⁴	Egypt	Case-control	Convenience	50	W	Infertility clinic	Women with idiopathic infertility	Endocervical	NAAT	6%
Azab and Hassouna (2008) ¹⁴⁵	Egypt	Cross-sectional	Convenience	70	W	Infertility clinic	Nearly half of women with TFI	Serum	ELISA (IgG)	28.6%
Badary (1996) ³⁸	Egypt	Case-control	Convenience	60	W	Infertility clinic	Women with idiopathic infertility	Endocervical	DFA	33%
Berry and El Shabrawy (1996) ³⁹	Egypt	Case-control	Convenience	70	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	EIA (IgG)	18.6%
Elkayal et al (2015) ¹⁴⁶	Egypt	Case-control	Convenience	100	W	Infertility clinic	Women with mixed infertility diagnosis	Endocervical	ELISA	3%
Elkayal et al (2015) ¹⁴⁶	Egypt	Case-control	Convenience	100	W	Infertility clinic	Women with mixed infertility diagnosis	Endocervical	NAAT	3%
El Sayed et al (1997) ¹⁴⁷	Egypt	Cross-sectional	Convenience	22	W	Infertility clinic	Women with TFI	Serum	MIF (IgG)	81.8%
El Sayed et al (1997) ¹⁴⁷	Egypt	Cross-sectional	Convenience	78	W	Infertility clinic	Women without TFI	Serum	MIF (IgG)	7.7%
Inhorn and Buss (1993) ¹⁴⁸	Egypt	Case-control	Convenience	83	W	Hospital	Majority of women without TFI	Unclear	Unclear	33%
Makled et al (2013) ¹⁴⁹	Egypt	Cross-sectional	Simple random sampling	27	W	Infertility clinic	Women with TFI	Serum	ELISA (IgG)	85.2%
Makled et al (2013) ¹⁴⁹	Egypt	Cross-sectional	Simple random sampling	51	W	Infertility clinic	Women without TFI	Serum	ELISA (IgG)	13.7%
Nada et al (2015) ⁴⁵	Egypt	Case-control	Convenience	100	W	Infertility clinic	Women with idiopathic infertility	Endocervical	NAAT	15%
Sadek et al (1993) ¹⁵⁰	Egypt	Case-control	Convenience	43	W	Infertility clinic	Infertile women in infertile couples with sperm antibodies	Unclear	DFA	18.6%
Sadek et al (1993) ¹⁵⁰	Egypt	Case-control	Convenience	37	W	Infertility clinic	Women partners in infertile couples with sperm antibodies	Unclear	DFA	18.9%
Sadek et al (1993) ¹⁵⁰	Egypt	Case-control	Convenience	62	M	Infertility clinic	Men partners in infertile couples with sperm antibodies	Unclear	DFA	19.4%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Sadek et al (1993) ¹⁵⁰	Egypt	Case-control	Convenience	18	M	Infertility clinic	Infertile men in infertile couples with sperm antibodies	Unclear	DFA	22.2%
Siam and Hefzy (2012) ¹⁵¹	Egypt	Case-control	Convenience	90	W	Gynaecology clinic	Women with idiopathic infertility	Serum	ELISA (IgG)	20%
Siam and Hefzy (2012) ¹⁵¹	Egypt	Case-control	Convenience	90	W	Gynaecology clinic	Women with idiopathic infertility	Urine	NAAT	4.4%
Younis et al (2000) ¹⁵²	Egypt	Cross-sectional	Convenience	30	W	Infertility clinic	Women with TFI	Serum	MIF (IgG)	46.7%
Younis et al (2000) ¹⁵²	Egypt	Cross-sectional	Convenience	14	W	Infertility clinic	Women without TFI	Serum	MIF (IgG)	50.0%
Zaitun and Zaitoun (1990) ¹⁵³	Egypt	Cross-sectional	Convenience	20	W	Infertility clinic	Women with TFI	Serum	Unclear	25%
Zaitun and Zaitoun (1990) ¹⁵³	Egypt	Cross-sectional	Convenience	30	W	Infertility clinic	Women without TFI	Serum	Unclear	3.3%
Zaki (1989) ⁴⁷	Egypt	Cross-sectional	Convenience	100	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	Culture	7%
Zytoon (1994) ¹⁵⁴	Egypt	Cross-sectional	Convenience	75	W	Infertility clinic	Women with mixed infertility diagnosis	Endocervical	Culture	65.3%
Ahmadi et al (2018) ⁴⁹	Iran	Case-control	Convenience	165	M	Infertility clinic	Men with male factor infertility	Semen	NAAT	4.2%
Badami and Salari (2001) ⁵¹	Iran	Case-control	Convenience	125	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	DFA	8.8%
Badami and Salari (2001) ⁵¹	Iran	Case-control	Convenience	125	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	Unclear	20.8%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Infertility clinic	Women with mixed infertility diagnosis	Urine	NAAT	4.8%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	M	Infertility clinic	40% of men had male factor infertility	Urine	NAAT	4.4%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	EIA (IgM)	4%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgA)	0
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	15.6%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	M	Infertility clinic	40% of men had male factor infertility	Serum	EIA (IgM)	1.2%
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	M	Infertility clinic	40% of men had male factor infertility	Serum	ELISA (IgA)	0
Dehghan et al (2017) ⁵⁷	Iran	Case-control	Convenience	250	M	Infertility clinic	40% of men had male factor infertility	Serum	ELISA (IgG)	18%
Golshani et al (2007) ¹⁵⁵	Iran	Cross-sectional	Convenience	200	M	Infertility clinic	Majority of men had male factor infertility	Semen	NAAT	18.0%
Goshayeshi et al (2015) ⁵⁸	Iran	Case-control	Convenience	100	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	NAAT	21.0%
Hajikhani et al (2013) ¹⁵⁶	Iran	Cross-sectional	Convenience	51	W	Infertility clinic	Women with TFI	Endocervical	Culture	3.9%
Hajikhani et al (2013) ¹⁵⁶	Iran	Cross-sectional	Convenience	51	W	Infertility clinic	Women with TFI	Endocervical	NAAT	11.7%
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	32	W	Infertility clinic	Women with TFI	Vaginal	NAAT	9.4%
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	68	W	Infertility clinic	Women with ovarian and other infertility	Vaginal	NAAT	2.9%
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	32	W	Infertility clinic	Women with TFI	Serum	ELISA (IgM)	9.4%
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	68	W	Infertility clinic	Women with ovarian and other infertility	Serum	ELISA (IgM)	4.4%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	32	W	Infertility clinic	Women with TFI	Serum	ELISA (IgG)	0
Joolayi et al (2017) ⁶²	Iran	Case-control	Convenience	68	W	Infertility clinic	Women with ovarian and other infertility	Serum	ELISA (IgG)	0
Kajbaf and Gholamnezhad (1998) ⁶³	Iran	Case-control	Convenience	101	W	Infertility clinic	Women with mixed infertility diagnosis	Endocervical	DFA	7.9%
Kajbaf and Gholamnezhad (1998) ⁶³	Iran	Case-control	Convenience	101	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	17.8%
Kalantar et al (2007) ¹⁵⁷	Iran	Cross-sectional	Convenience	91	W	Infertility clinic	Majority of women had female factor infertility	Serum	ELISA (IgG)	0
Kalantar et al (2007) ¹⁵⁷	Iran	Cross-sectional	Convenience	91	W	Infertility clinic	Majority of women had female factor infertility	Vaginal	NAAT	0
Kamyabi (2009) ⁶⁴	Iran	Case-control	Convenience	35	W	Gynaecology clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	22.9%
Mansour Ghanaie (2014) ¹⁵⁸	Iran	Cross-sectional	Convenience	135	W	Infertility clinic	Majority of women without TFI	Endocervical	NAAT	19.3%
Ministry of Health and Medical Education (2008) ⁵⁷	Iran	Case-control	Convenience	46	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgG)	23.9%
Marashi et al (2014) ⁶⁶	Iran	Case-control	Convenience	150	W	Infertility clinic	Women with idiopathic infertility	Endocervical	DFA	15.3%
Marashi et al (2014) ⁶⁶	Iran	Case-control	Convenience	150	W	Infertility clinic	Women with idiopathic infertility	Endocervical	NAAT	32%
Ministry of Health and Medical Education (2008) ⁵⁷	Iran	Case-control	Convenience	125	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	DFA	8.8%
Ministry of Health and Medical Education (2008) ⁵⁷	Iran	Cross-sectional	Convenience	100	M	Infertility clinic	Men with unclear infertility diagnosis	Unclear	NAAT	9%
Moazenchi et al (2018) ¹⁵⁹	Iran	Cross-sectional	Convenience	1080	M	Infertility clinic	Men with unclear infertility diagnosis	Serum	ELISA (IgA)	4.3%
Moazenchi et al (2018) ¹⁵⁹	Iran	Cross-sectional	Convenience	1080	M	Infertility clinic	Men with unclear infertility diagnosis	Semen	NAAT	10%
Mousavi et al (2014) ⁶⁹	Iran	Case-control	Convenience	104	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	NAAT	4.8%
Nan Bakhsh et al (2008) ¹⁶⁰	Iran	Cross-sectional	Convenience	144	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	11.1%
Nikbakht et al (2008) ¹⁶¹	Iran	Case-control	Convenience	125	W	Infertility clinic	Women with TFI	Unclear	ELISA (unclear)	23.2%
Peivandi et al (2009) ¹⁶²	Iran	Cross-sectional	Convenience	110	W	Infertility clinic	Majority of women with TFI	Serum	MIF (IgG)	24.5%
Rashidi et al (2007) ¹⁶³	Iran	Cross-sectional	Convenience	300	W	Infertility clinic	Women with mixed infertility diagnosis	Unclear	ELISA (unclear)	32.3%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	44	W	Infertility clinic	Women with TFI	Urine	NAAT	4.5%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	190	W	Infertility clinic	Women with ovarian and other infertility	Urine	NAAT	14.2%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	44	W	Infertility clinic	Women with TFI	Serum	ELISA (IgM)	2.3%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	190	W	Infertility clinic	Women with ovarian and other infertility	Serum	ELISA (IgM)	0.5%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	44	W	Infertility clinic	Women with TFI	Serum	ELISA (IgG)	9.1%
Rashidi et al (2013) ⁷¹	Iran	Case-control	Convenience	190	W	Infertility clinic	Women with ovarian and other infertility	Serum	ELISA (IgG)	8.4%
Sadrpour et al (2013) ¹⁶⁴	Iran	Cross-sectional	Convenience	120	M	Infertility clinic	Men with male factor infertility	Semen	NAAT	3%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Sattari et al (2017) ⁷⁵	Iran	Case-control	Convenience	184	W	Infertility clinic	Majority of women without TFI	Serum	ELISA (IgM)	5-4%
Sattari et al (2017) ⁷⁵	Iran	Case-control	Convenience	184	W	Infertility clinic	Majority of women without TFI	Serum	ELISA (IgG)	35-9%
Siahkali and Amini (2018) ¹⁶⁵	Iran	Cross-sectional	Convenience	60	M	Infertility clinic	Men with idiopathic infertility	Semen	NAAT	5-0%
Abid and Al-Zwaid (2015) ¹⁶⁶	Iraq	Case-control	Convenience	61	W	Infertility clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	30%
Ahmed (2012) ¹⁶⁷	Iraq	Case-control	Convenience	47	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	NAAT	29-8%
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	54	W	Infertility clinic	Women with unclear infertility diagnosis	Endocervical	ELFA	9-3%
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	54	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	IFAT (unclear)	11-1%
Ali and Al-Kazaz (2018) ⁸⁵	Iraq	Case-control	Convenience	63	M	Clinic	Men with male factor infertility	Semen	NAAT	17-4%
Al-Kattan and Mohammed (2013) ¹⁶⁸	Iraq	Cross-sectional	Convenience	54	W	Infertility clinic	Women with TFI or adhesions	Serum	ELISA (IgG)	51-9%
Al-Kattan and Mohammed (2013) ¹⁶⁸	Iraq	Cross-sectional	Convenience	67	W	Infertility clinic	Women without TFI or endometriosis	Serum	ELISA (IgG)	29-9%
Dawood (2011) ¹⁶⁹	Iraq	Cross-sectional	Convenience	30	W	Hospital	Women with unclear infertility diagnosis	Serum	ELISA (IgM)	86-6%
Dawood (2011) ¹⁶⁹	Iraq	Cross-sectional	Convenience	30	W	Hospital	Women with unclear infertility diagnosis	Serum	ELISA (IgA)	3-3%
Dawood (2011) ¹⁶⁹	Iraq	Cross-sectional	Convenience	30	W	Hospital	Women with unclear infertility diagnosis	Serum	ELISA (IgG)	53-3%
Dawood (2011) ¹⁶⁹	Iraq	Cross-sectional	Convenience	100	W	Hospital	Women with unclear infertility diagnosis	Endocervical	NAAT	30%
Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	52	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgG)	25%
Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	52	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgM)	42-3%
Ismail and Ali (2012) ⁸⁸	Iraq	Case-control	Convenience	52	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgA)	3-8%
Mohammed et al (2017) ⁹⁰	Iraq	Case-control	Convenience	80	W	Gynaecology clinic	Women with mixed infertility diagnosis	Endocervical	NAAT	13-8%
Mohammed et al (2017) ⁹⁰	Iraq	Case-control	Convenience	80	W	Gynaecology clinic	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	2-5%
Yahya and Al-Siraj (2009) ⁹¹	Iraq	Cross-sectional	Convenience	296	M	Laboratory	Men with unclear infertility diagnosis	Serum	Culture	4-0%
Abusarah et al (2013) ⁹²	Jordan	Case-control	Convenience	81	M	Gynaecology clinic	Men with male factor infertility	Urine	NAAT	4-9%
Al-Ramahi et al (2008) ⁹³	Jordan	Case-control	Convenience	66	W	Infertility clinic	Women with idiopathic infertility	Endocervical	NAAT	3-0%
Al-Ramahi et al (2008) ⁹³	Jordan	Case-control	Convenience	19	W	Infertility clinic	Women with TFI	Endocervical	NAAT	0
Al-Ramahi et al (2008) ⁹³	Jordan	Case-control	Convenience	38	W	Infertility clinic	Women with male factor infertility	Endocervical	NAAT	7-9%
Al-Ramahi et al (2008) ⁹³	Jordan	Case-control	Convenience	29	W	Infertility clinic	Women with ovarian infertility	Endocervical	NAAT	3-4%
Al-Sweih et al (2012) ¹⁰⁰	Kuwait	Case-control	Convenience	127	M	Infertility clinic	Men with unclear infertility diagnosis	Semen	NAAT	3-9%
Radouani et al (1998) ¹⁰⁶	Morocco	Case-control	Convenience	200	M	Infertility clinic	Majority of men had male factor infertility	Serum	MIF (unclear)	21-5%
Radouani et al (1998) ¹⁰⁶	Morocco	Case-control	Convenience	81	W	Infertility clinic	Women with unclear infertility diagnosis	Serum	MIF (unclear)	44-4%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Al Subhi et al (2013) ¹⁷⁰	Oman	Cross-sectional	Convenience	51	W	Infertility clinic	Women with TFI	Endocervical	EIA	5.9%
Al Subhi et al (2013) ¹⁷⁰	Oman	Cross-sectional	Convenience	167	W	Infertility clinic	Women without TFI	Endocervical	EIA	4.8%
Qayum and Khalid-bin-Saleem (2013) ¹⁷¹	Pakistan	Cross-sectional	Convenience	80	W	Gynaecology clinic	Women with unclear infertility diagnosis	Urine	Unclear	7.5%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	69	W	Infertility clinic	Women undergoing IVF in 2000	Serum	ELISA (IgM)	11.6%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	268	W	Infertility clinic	Women undergoing IVF in 2001	Serum	ELISA (IgM)	23.9%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	316	W	Infertility clinic	Women undergoing IVF in 2002	Serum	ELISA (IgM)	33.5%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	399	W	Infertility clinic	Women undergoing IVF in 2003	Serum	ELISA (IgM)	9.3%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	586	W	Infertility clinic	Women undergoing IVF in 2004	Serum	ELISA (IgM)	4.6%
Al-Hindi et al (2010) ¹⁷²	Palestine	Cross-sectional	Convenience	316	W	Infertility clinic	Women undergoing IVF in 2005	Serum	ELISA (IgM)	2.8%
Abdul Jabbar (1990) ¹⁷³	Saudi Arabia	Cross-sectional	Convenience	13	W	Infertility clinic	Women with TFI	Endocervical	DFA	53.8%
Abdul Jabbar (1990) ¹⁷³	Saudi Arabia	Cross-sectional	Convenience	18	W	Infertility clinic	Women without TFI	Endocervical	DFA	11.1%
Abdul Jabbar (1990) ¹⁷³	Saudi Arabia	Cross-sectional	Convenience	34	M	Infertility clinic	Men with unclear infertility diagnosis	Urethral	DFA	26.4%
Alfarraj et al (2015) ¹⁷⁴	Saudi Arabia	Case-control	Convenience	100	W	Infertility clinic	Women with mixed infertility diagnosis	Endocervical	NAAT	8.0%
Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	41	W	Gynaecology clinic	Women with unclear infertility diagnosis	Endocervical	Culture	9.5%
Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	41	W	Gynaecology clinic	Women with unclear infertility diagnosis	Serum	MIF (IgM)	0
Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	41	W	Gynaecology clinic	Women with unclear infertility diagnosis	Serum	MIF (IgG)	16.7%
Kamel (2013) ¹¹⁶	Saudi Arabia	Randomised controlled trial‡	Convenience	640	W	Gynaecology clinic	Women with unclear infertility diagnosis	Endocervical	Culture	12.0%
Kamel (2013) ¹¹⁶	Saudi Arabia	Randomised controlled trial‡	Convenience	640	W	Gynaecology clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgA)	5%
Kamel (2013) ¹¹⁶	Saudi Arabia	Randomised controlled trial‡	Convenience	640	W	Gynaecology clinic	Women with unclear infertility diagnosis	Serum	ELISA (IgG)	8.0%
Sabra and Al-Harbi (2014) ¹⁷⁵	Saudi Arabia	Cross-sectional	Convenience	148	M	Infertility clinic	Men with male factor infertility	Semen	Giemsa stain	8.1%
Almroth et al (2005) ¹²³	Sudan	Case-control	Convenience	81	W	Infertility clinic	More than half of women with TFI	Serum	EIA (IgG)	14%
Alkayer et al (2017) ¹²⁵	Syria	Case-control	Convenience	23	W	Hospital	Women with mixed infertility diagnosis	Serum	ELISA (IgG)	17.1%
Gdoura et al (2001a) ¹⁷⁶	Tunisia	Cross-sectional	Convenience	92	M	Infertility clinic	Men with unclear infertility diagnosis	Urethral	NAAT	18.5%
Gdoura et al (2001b) ¹⁷⁷	Tunisia	Cross-sectional	Convenience	92	M	Infertility clinic	Men with unclear infertility diagnosis	Serum	MIF (IgG)	9.8%
Gdoura et al (2001a) ¹⁷⁶	Tunisia	Cross-sectional	Convenience	92	M	Infertility clinic	Men with unclear infertility diagnosis	Urethral	DFA	4.3%
Gdoura et al (2001a) ¹⁷⁶	Tunisia	Cross-sectional	Convenience	92	M	Infertility clinic	Men with unclear infertility diagnosis	Urethral	Culture	1.1%
Gdoura et al (2001a) ¹⁷⁶	Tunisia	Cross-sectional	Convenience	92	M	Infertility clinic	Men with unclear infertility diagnosis	Urethral	Unclear	8.7%
Gdoura et al (2001b) ¹⁷⁷	Tunisia	Cross-sectional	Convenience	92	W	Infertility clinic	Partners of infertile men	Endocervical	NAAT	26.1%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†
(Continued from previous page)										
Gdoura et al (2001b) ¹⁷⁷	Tunisia	Cross-sectional	Convenience	92	W	Infertility clinic	Partners of infertile men	Serum	MIF (IgG)	17.4%
Gdoura et al (2008) ¹⁷⁸	Tunisia	Cross-sectional	Convenience	104	M	Infertility clinic	Men with male factor infertility	Urine	NAAT	39.4%
Sellami et al (2014) ¹⁷⁹	Tunisia	Cross-sectional	Convenience	85	M	Infertility clinic	Men with unclear infertility diagnosis	Semen	NAAT	15.2%
Women with miscarriage (or abortion of unknown cause)										
Zaki (1989) ⁴⁷	Egypt	Cross-sectional	Convenience	100	W	Gynaecology clinic	Presenting with abortion	Endocervical	Culture	5%
Ahmadi et al (2016b) ¹⁸⁰	Iran	Case-control	Convenience	109	W	Family planning clinic	Spontaneous abortion	Endocervical	NAAT	22.9%
Bagheri and Roghanian (2014) ¹⁸¹	Iran	Cross-sectional	Convenience	70	W	Hospital	Recent or recurrent miscarriage	Vaginal	NAAT	1.4%
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	97	W	Fertility centre	Recent or recurrent miscarriage	Vaginal	NAAT	11.3%
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	97	W	Fertility centre	Recent or recurrent miscarriage	Serum	ELISA (IgA)	2.1%
Bagheri et al (2018) ⁵³	Iran	Case-control	Convenience	97	W	Fertility centre	Recent or recurrent miscarriage	Serum	ELISA (IgG)	4.1%
Jahromi et al (2010) ⁶⁰	Iran	Case-control	Convenience	220	W	Gynaecology clinic	Spontaneous abortion	Endocervical	DFA	25.5%
Massiha et al (2010) ¹⁸²	Iran	Cross-sectional	Convenience	84	W	Hospital	Presenting with abortion	Unclear	Unclear	2.3%
Salari and Badami (2002) ¹⁸³	Iran	Case-control	Convenience	125	W	Hospital	Recurrent abortion	Endocervical	DFA	7.2%
Sisakht et al (2017) ⁷⁶	Iran	Case-control	Convenience	77	W	Gynaecology clinic	Spontaneous abortion	Urine	NAAT	9.3%
Zahirnia et al (2018) ⁷⁸	Iran	Cross-sectional	Convenience	124	W	Gynaecology clinic	Presenting with abortion	Vaginal	NAAT	15.3%
Abdul-Karim et al (2009) ⁸⁰	Iraq	Case-control	Convenience	79	W	Hospital	Presenting with abortion	Serum	ELISA (IgG)	6.4%
Abdulkhudher et al (2014) ⁷⁹	Iraq	Case-control	Convenience	60	W	Antenatal clinic	Recent or recurrent miscarriage	Serum	ELISA (IgM)	38.3%
Abdulkhudher et al (2014) ⁷⁹	Iraq	Case-control	Convenience	60	W	Antenatal clinic	Recent or recurrent miscarriage	Serum	ELISA (IgG)	33.3%
Ahmed (2008) ⁸²	Iraq	Case-control	Convenience	60	W	Hospital	Recurrent miscarriage	Serum	ELISA (unclear)	0
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	89	W	Family planning clinic	Recent or recurrent abortion	Endocervical	ELFA	12.4%
Al-Husseinei et al (2009) ⁸⁴	Iraq	Case-control	Convenience	89	W	Family planning clinic	Recent or recurrent abortion	Serum	IFAT (unclear)	14.6%
Alkhafaf (2013) ⁸⁶	Iraq	Case-control	Convenience	123	W	Hospital	Spontaneous abortion	Serum	ELISA (IgG)	17.1%
Al-Nuaimy and Al-Jandeel (2018) ¹⁸⁴	Iraq	Case-control	Convenience	120	W	Hospital	Recent or recurrent abortion	Endocervical	NAAT	17.5%
Al-Nuaimy and Al-Jandeel (2018) ¹⁸⁴	Iraq	Case-control	Convenience	120	W	Hospital	Recent or recurrent abortion	Serum	ELISA (IgG)	14.2%
Mohammed et al (2012) ⁸⁹	Iraq	Case-control	Convenience	62	W	Gynaecology clinic	Three or more miscarriages	Serum	ELISA (IgM)	16.1%
Mohammed et al (2012) ⁸⁹	Iraq	Case-control	Convenience	34	W	Gynaecology clinic	Less than three miscarriages	Serum	ELISA (IgM)	29.4%
Salman (2016) ¹⁸⁵	Iraq	Cross-sectional	Convenience	184	W	Gynaecology clinic	Presenting with abortion	Serum	ELISA (IgM)	21.2%
Salman (2016) ¹⁸⁵	Iraq	Cross-sectional	Convenience	184	W	Gynaecology clinic	Presenting with abortion	Serum	ELISA (IgG)	8.2%
Hossain (1988) ¹¹⁵	Saudi Arabia	Cross-sectional	Convenience	12	W	Hospital	Recurrent miscarriage	Endocervical	Culture	16.7%

(Table 2 continues on next page)

	Country	Study design	Sampling*	Sample size	Sex	Study context	Population characteristics	Specimen	Assay type	Prevalence†	
(Continued from previous page)											
	Hossain (1988) ¹³⁵	Saudi Arabia	Cross-sectional	Convenience	12	W	Hospital	Recurrent miscarriage	Serum	MIF (IgM)	0
	Hossain (1988) ¹³⁵	Saudi Arabia	Cross-sectional	Convenience	12	W	Hospital	Recurrent miscarriage	Serum	MIF (IgG)	16.7%
Women with ectopic pregnancy											
	Diab (1993) ⁴⁰	Egypt	Case-control	Convenience	30	W	Family planning clinic	Ectopic pregnancy	Serum	EIA (IgG)	30%
	Abdullah (2012) ⁸¹	Iraq	Case-control	Convenience	24	W	Hospital	Ectopic pregnancy	Serum	ELISA (IgM)	4%
	Abdullah (2012) ⁸¹	Iraq	Case-control	Convenience	24	W	Hospital	Ectopic pregnancy	Serum	ELISA (IgG)	45%

DFA=direct fluorescent assay. EIA=enzyme immunoassay. ELFA=enzyme-linked fluorescence assay. IFAT=indirect fluorescent antibody test. M=men or sample predominantly of men. MIF=micro-immunofluorescence. NAAT=nucleic acid amplification test. NGOs=non-governmental organisations. STI=sexually transmitted infection. TFI=tubal factor infertility. W=women or sample predominantly of women. *Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).^{127,128} Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.¹²⁹ Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.¹²⁹ †The decimal places of the prevalence measures are as reported in the original report, but prevalence figures with more than one decimal place were rounded to one decimal place. ‡The extracted prevalence measure is for the baseline measurement.

Table 2: Studies reporting *Chlamydia trachomatis* prevalence in populations at high risk, infertility clinic attendees, women with miscarriage, and women with ectopic pregnancy in the Middle East and north Africa

High prevalence was observed in infertility clinic attendees, for both women and men (n=135), in which current infection prevalence ranged from 0 to 65.3% with a median of 9.2%, whereas ever infection prevalence ranged from 0 to 85.2% with a median of 18.6% (tables 2 and 3). Similarly, high prevalence was observed in women with miscarriage (n=27), in which current infection prevalence ranged from 1.4% to 25.5% with a median of 12.4%, whereas ever infection prevalence ranged from 4.1% to 33.3% with a median of 14.2% (tables 2 and 3).

Table 3 summarises the prevalence for other at-risk populations, and table 2 and the appendix (pp 9–14) include the full data.

The summarised and study-specific risk of bias and precision assessments are shown in the appendix (pp 15–27). Briefly, 166 (30.1%) of 552 prevalence measures were based on samples including 200 participants or more, and were classified as having higher precision. Although convenience sampling was the most common sampling methodology (495 [89.7%] of 552), probability-based sampling methods, such as respondent-driven sampling, are of increasing use for populations at high risk (18 [45%] of 40 studies in female sex workers and men who have sex with men). Almost all studies (524 [94.9%] of 552) specified the type of biological assay used for infection ascertainment (low risk of bias for this domain). Response rate was, however, unclear for 417 (75.5%) of 552 studies. Prevalence studies were overall of reasonable quality; only eight (1.4%) of 552 had high risk of bias in two or more quality domains.

Table 3 shows the meta-analyses' results for the pooled average *C trachomatis* prevalence for each at-risk population, stratified by type of assay used for infection ascertainment. Current infection prevalence was estimated at 3.0%

(95% CI 2.3–3.8) in general populations, 2.8% (1.0–5.2) in populations at intermediate risk, 13.2% (7.2–20.7) in female sex workers, 1.2% (0.2–2.8) for genital infections and 7.7% (4.2–12.0) for rectal infections in men who have sex with men, 11.3% (9.0–13.7) in infertility clinic attendees, 12.4% (7.9–17.7) in women with miscarriage, 12.4% (9.4–15.7) in symptomatic women, and 17.4% (12.5–22.8) in symptomatic men.

Meanwhile, pooled average prevalence of ever infection was estimated at 6.9% (4.3–10.0) in general populations, 1.4% (0.8–2.4) in populations at intermediate risk, 80.9% (43.8–100) in female sex workers, 21.5% (16.3–27.2) in infertility clinic attendees, 12.4% (6.6–19.5) in women with miscarriage, 37.1% (22.4–53.0) in women with ectopic pregnancy, 22.7% (15.4–31.0) in symptomatic women, and 16.9% (9.4–25.8) in symptomatic men (table 3).

Evidence for heterogeneity in *C trachomatis* prevalence estimates was observed; p values for Cochran's Q statistic was <0.0001 in most meta-analyses (table 3). Prediction intervals were generally wide affirming high heterogeneity. I² was also mostly more than 70%, indicating that most variability is due to true differences in effect size across studies rather than chance.

Figures 2 and 3 and the appendix (pp 28–40) summarise the results of subgroup meta-analyses in various sub-populations. These data show the results stratified by sex or by genital versus rectal infection (the latter only for men who have sex with men), for studies reporting current infection prevalence based on NAAT and those reporting ever infection prevalence, as well as by assay type for studies reporting current infection prevalence. Subgroup meta-analyses in infertile populations stratified by infertility diagnosis and by assay type are shown in the appendix (pp 41–42).

Table 4 summarises results of the meta-regression analyses. In the univariable analyses, at-risk population, assay type, sampling methodology, sample size, year of publication, year of data collection, country, response rate, and sex were associated with prevalence at $p \leq 0.2$. Alignment with meta-regression underlying assumption

	Studies (n)	Samples		<i>C trachomatis</i> prevalence (median [range])	Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI])	Heterogeneity measures		
		Tested	<i>C trachomatis</i> positive			Q (p value)*	I ² † (95% CI)	Prediction interval‡
General populations								
Current genital infection								
NAAT	48	25397	748	2.9% (0-15.5)	3.1 (2.2-4.2)	714.3 (p<0.0001)	91.4% (89.4-93.0)	0.0-12.4
Culture	4	4464	55	5.8% (1.0-15.0)	4.3 (0.3-11.4)	22.5 (p<0.0001)	86.6% (67.7-94.5)	0.0-50.9
Other§	23	128013	328	3.5% (0-19.9)	2.4 (1.6-3.4)	722.3 (p<0.0001)	97.0% (96.2-97.5)	0.0-7.2
Overall current genital infection	75	157874	1131	3.0% (0-19.9)	3.0 (2.3-3.8)	2703.5 (p<0.0001)	97.3% (96.9-97.6)	0.0-10.9
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	35	5877	525	4.7% (0-37.9)	6.9 (4.3-10.0)	226.1 (p<0.0001)	86.7% (82.2-90.1)	0.0-30.2
IgM (recent infection)	13	2843	74	1.6% (0-14.0)	1.8 (0.3-3.9)	77.7 (p<0.0001)	84.6% (75.1-90.4)	0.0-12.4
IgA	4	377	12	4.3% (0-40.4)	6.2 (0.0-21.6)	37.8 (p<0.0001)	92.1% (82.9-96.3)	0.0-93.7
Not specified (IgG, IgM, or IgA)	9	1081	61	4.5% (0-14.8)	4.3 (1.9-7.4)	34.5 (p<0.0001)	76.8% (55.7-87.8)	0.0-17.3
Unclear	1	250	8	..	3.2 (1.4-6.2)
Populations at intermediate risk								
Current genital infection								
NAAT	12	2815	69	1.5% (0-38.0)	2.6 (0.8-5.2)	117.4 (p<0.0001)	75.6% (56.0-86.5)	0.0-16.1
Culture
Other§	1	308	15	..	4.9 (2.8-7.9)
Overall current genital infection	13	3123	84	2.0% (0-38.0)	2.8 (1.0-5.2)	127.0 (p<0.0001)	90.6% (85.7-93.8)	0.0-15.8
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	1	1041	15	..	1.4 (0.8-2.4)
IgM (recent infection)
IgA
Not specified (IgG, IgM, or IgA)
Unclear
Populations at high risk								
Female sex workers								
Current genital infection								
NAAT	12	4877	590	8.4% (0.9-72.9)	12.9 (6.5-21.0)	602.1 (p<0.0001)	98.2% (97.6-98.6)	0.0-52.0
Culture	1	116	8	..	6.9 (3.0-13.1)
Other§	1	30	9	..	30.0 (14.7-49.4)
Overall current genital infection	14	5023	607	8.4% (0.9-72.9)	13.2 (7.2-20.7)	611.7 (p<0.0001)	97.9% (97.3-98.3)	0.0-50.9
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	4	472	364	90.0% (19.8-100)	80.9 (43.8-100.0)	209.9 (p<0.0001)	98.6% (97.7-99.1)	0.0-100.0
IgM (recent infection)	1	154	45	..	29.2 (22.2-37.1)
IgA
Not specified (IgG, IgM, or IgA)	1	30	11	..	36.7 (19.9-56.1)
Unclear
Men who have sex with men								
Current genital infection								
NAAT	12	2680	51	1.2% (0-8.8)	1.2 (0.2-2.8)	76.2 (p<0.0001)	85.6% (76.5-91.1)	0.0-9.5
Culture
Other§
Current rectal infection								
PCR	7	1506	129	6.3% (3.6-18.3)	7.7 (4.2-12.0)	40.6 (p<0.0001)	85.2% (71.5-92.3)	0.0-24.9
Overall current infection	19	4186	180	3.6% (0-18.3)	3.0 (1.2-5.4)	231.8 (p<0.0001)	92.2% (89.3-94.4)	0.0-17.9

(Table 3 continues on next page)

	Studies (n)	Samples		<i>C trachomatis</i> prevalence (median [range])	Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI])	Heterogeneity measures		
		Tested	<i>C trachomatis</i> positive			Q (p value)*	I ² (95% CI)	Prediction interval‡
(Continued from previous page)								
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)
IgM (recent infection)
IgA
Not specified (IgG, IgM, or IgA)
Unclear	1	2531	890	..	35.2 (33.3–37.1)
Infertility clinic attendees								
Current genital infection								
NAAT	37	4653	539	8.0% (0–39.4)	10.2 (7.5–13.1)	310.9 (p<0.0001)	88.4% (85.0–91.0)	0.1–31.0
Culture	7	1149	176	9.5% (1.1–65.3)	14.4 (4.8–27.7)	140.8 (p<0.0001)	95.7% (93.3–97.3)	0.0–69.2
Other§	20	1844	203	10.2% (3.0–53.8)	12.3 (8.5–16.5)	112.8 (p<0.0001)	83.2% (75.1–88.6)	0.5–33.7
Overall current genital infection	64	7646	918	9.2% (0–65.3)	11.3 (9.0–13.7)	574.9 (p<0.0001)	89.0% (86.7–90.9)	0.2–33.3
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	37	3608	689	18.6% (0–85.2)	21.5 (16.3–27.2)	531.8 (p<0.0001)	93.2% (91.6–94.6)	0.2–59.7
IgM (recent infection)	17	3145	332	4.6% (0–86.7)	10.2 (5.0–16.8)	435.4 (p<0.0001)	96.3% (95.2–97.2)	0.0–47.0
IgA	6	2302	82	3.6% (0–5.0)	1.8 (0.1–4.7)	54.4 (p<0.0001)	90.8% (82.8–95.1)	0.0–16.4
Not specified (IgG, IgM, or IgA)	5	760	211	23.2% (11.1–44.4)	26.1 (17.9–35.3)	27.5 (p<0.0001)	85.4% (67.8–93.4)	2.6–61.8
Unclear	6	430	73	14.8% (3.3–33.0)	14.9 (7.0–24.8)	30.0 (p<0.0001)	83.3% (65.1–92.0)	0.0–53.3
Women with miscarriage								
Current genital infection								
NAAT	6	597	87	13.3% (1.4–22.9)	12.9 (7.4–19.5)	24.4 (p=0.0002)	79.5% (55.3–90.6)	0.1–38.9
Culture	2	112	7	10.9% (5.0–16.7)	7.1 (0.0–21.8)	2.1 (p=0.1483)	52.1% (0.0–88.0)	..
Other§	3	434	76	12.4% (7.2–25.5)	14.4 (4.9–27.6)	21.9 (p<0.0001)	90.8% (76.1–96.5)	0.0–100.0
Overall current genital infection	11	1143	170	12.4% (1.4–25.5)	12.4 (7.9–17.7)	58.0 (p<0.0001)	82.8% (70.5–89.9)	0.6–34.3
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	7	675	84	14.2% (4.1–33.3)	12.4 (6.6–19.5)	33.5 (p<0.0001)	82.1% (64.2–91.0)	0.0–39.6
IgM (recent infection)	5	352	82	21.2% (0–38.3)	21.2 (11.9–32.2)	16.2 (p=0.0028)	75.3% (39.1–89.9)	0.0–61.8
IgA	1	97	2	..	2.1 (0.3–7.3)
Not specified (IgG, IgM, or IgA)	2	149	13	7.3% (0–14.6)	4.7 (0.0–27.9)	16.1 (p<0.0001)	93.8% (80.0–98.1)	..
Unclear	1	84	2	..	2.3 (0.3–8.3)
Women with ectopic pregnancy								
Current genital infection								
NAAT
Culture
Other§
Overall current genital infection
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	2	54	20	37.5% (30.0–45.0)	37.1 (22.4–53.0)	1.4 (p=0.2418)	27.0%	..
IgM (recent infection)	1	24	1	..	4.2 (0.1–21.1)
IgA
Not specified (IgG, IgM, or IgA)
Unclear
Symptomatic women								
Current genital infection								
NAAT	49	14 398	1123	8.0% (0–68.0)	8.8 (6.2–11.7)	1506.7 (p<0.0001)	96.8% (96.3–97.3)	0.0–35.4
Culture	10	2951	752	12.9% (0.7–69.4)	18.9 (4.1–40.9)	1511.1 (p<0.0001)	99.4% (99.3–99.5)	0.0–97.2
Other§	31	4796	729	14.7% (0–89.3)	16.8 (11.6–22.7)	723.8 (p<0.0001)	95.9% (94.9–96.6)	0.0–55.4

(Table 3 continues on next page)

	Studies (n)	Samples		<i>C trachomatis</i> prevalence (median [range])	Pooled average <i>C trachomatis</i> prevalence (estimate [95% CI])	Heterogeneity measures		
		Tested	<i>C trachomatis</i> positive			Q (p value)*	I ² † (95% CI)	Prediction interval‡
(Continued from previous page)								
Overall current genital infection	90	22145	2604	11.7% (0–89.3)	12.4 (9.4–15.7)	4323.7 (p<0.0001)	97.9% (97.7–98.1)	0.0–52.3
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	23	2377	609	18.2% (2.7–86.0)	22.7 (15.4–31.0)	454.1 (p<0.0001)	95.2% (93.8–96.2)	0.0–67.9
IgM (recent infection)	9	1042	160	3.1% (0.86–0)	13.9 (1.9–33.6)	452.7 (p<0.0001)	98.2% (97.6–98.7)	0.0–91.3
IgA	5	365	93	12.5% (3.7–59.6)	24.9 (6.6–49.6)	98.9 (p<0.0001)	96.0% (93.0–97.7)	0.0–100.0
Not specified (IgG, IgM, or IgA)	7	2530	761	15.0% (11.6–46.6)	23.5 (11.1–38.7)	346.8 (p<0.0001)	98.3% (97.6–98.4)	0.0–80.0
Unclear	6	943	83	7.7% (1.7–30.0)	8.7 (4.3–14.4)	32.2 (p<0.0001)	84.5% (67.9–92.5)	0.0–32.6
Symptomatic men								
Current genital infection								
NAAT	14	7160	726	12.2% (4.2–33.3)	13.9 (8.3–20.6)	488.7 (p<0.0001)	97.3% (96.5–98.0)	0.0–46.4
Culture	5	4744	75	9.3% (0.4–19.6)	8.7 (1.1–21.7)	147.5 (p<0.0001)	97.3% (95.6–98.3)	0.0–72.7
Other§	13	2499	355	27.6% (4.7–52.0)	26.3 (15.3–39.1)	351.7 (p<0.0001)	96.6% (95.4–97.5)	0.0–78.9
Overall current genital infection	32	14403	1156	15.5% (0.4–52.0)	17.4 (12.5–22.8)	1628.1 (p<0.0001)	98.1% (97.8–98.4)	0.0–53.3
Anti- <i>C trachomatis</i> immunoglobulins								
IgG (ever infection)	8	831	164	14.4% (5.1–46.0)	16.9 (9.4–25.8)	69.6 (p<0.0001)	89.9% (82.6–94.2)	0.0–52.8
IgM (recent infection)	3	330	24	3.9% (2.8–12.2)	6.0 (1.4–13.0)	9.3 (p=0.0095)	78.5% (31.0–93.3)	0.0–100.0
IgA
Not specified (IgG, IgM, or IgA)	5	1596	687	46.0% (10.0–49.1)	35.6 (18.5–54.9)	143.3 (p<0.0001)	97.2% (95.4–98.3)	0.0–97.3
Unclear	8	4876	233	14.3% (1.6–76.9)	24.0 (9.2–42.8)	498.1 (p<0.0001)	98.6% (98.1–99.0)	0.0–90.3

A minimum of two studies was necessary to do a meta-analysis. The same population might have contributed different measures for both current and ever infection with *C trachomatis*. NAAT=nucleic acid amplification test. *Cochran's Q statistic is a measure assessing the existence of heterogeneity in effect size of *C trachomatis* prevalence across studies. †I² is a measure assessing the magnitude of between-study variation that is due to differences in effect size of *C trachomatis* prevalence across studies rather than chance. ‡Prediction interval is a measure estimating the 95% CI of the distribution of true effect sizes of *C trachomatis* prevalence measures. §Other assays detecting current infection such as direct fluorescence assays, Giemsa staining, and enzyme-linked immunoassays applied to genital samples.

Table 3: Results of meta-analyses on studies reporting *Chlamydia trachomatis* prevalence in the Middle East and north Africa stratified by at-risk population and *C trachomatis* ascertainment

of normal random effects was confirmed through normal probability plots (appendix p 43).¹⁸⁶ Graphical illustrations of the fitted regression line for year of publication and year of data collection are shown in the appendix (p 44). Only at-risk population, assay type, sample size, country, and sex remained associated with *C trachomatis* prevalence in a multivariable model. No evidence was found for a temporal variation in prevalence (p=0.281 for year of publication), for sampling methodology (p=0.347), or for response rate (p=0.237). This model explained 29.0% of prevalence variation.

Relative to general populations, the adjusted odds ratio (aOR) was 11.28 (95% CI 5.78–22.01) for female sex workers, 7.17 (4.05–12.68) for symptomatic men, 4.93 (1.03–23.52) for women with ectopic pregnancy, 4.16 (1.72–10.08) for men who have sex with men, 3.39 (2.41–4.77) for infertility clinic attendees, 3.47 (2.47–4.87) for symptomatic women, 2.78 (1.57–4.93) for women with miscarriage, and 1.81 (0.79–4.13) for populations at intermediate risk. Other factors associated with *C trachomatis* prevalence were women versus men (aOR 1.61, 95% CI 1.05–2.46), Pakistan versus other Middle East or north African countries (0.39, 0.22–0.69), ever infection (anti-*C trachomatis* IgG; 2.17, 1.54–3.06) and current infection prevalence using assays other than

NAAT or culture versus NAAT (1.47, 1.02–2.13), and studies with higher (≥200 participants) versus lower precision (0.63, 0.48–0.83).

Discussion

We provided a comprehensive assessment of *C trachomatis* epidemiology in the Middle East and north Africa.^{2,3} Unexpectedly, given this region's sexually conservative norms and low observed levels of several viral STIs,^{10,11,187–189} *C trachomatis* current infection prevalence was 3% in the population at large, similar to WHO prevalence estimates for this region of about 3% in 2012³ and about 3.5% in 2016.¹⁹⁰ The prevalence was also in line with WHO estimates for the Western Pacific region (about 4%) and European region (about 3%),¹⁹⁰ where broad *C trachomatis* control programmes, including opportunistic testing, are standard in some high-income countries,^{191–193} but higher than that for South-east Asia region (about 1.5%) and lower than that for the African region (about 5%) and the region of the Americas (about 5.5%).¹⁹⁰ This high prevalence suggests substantial infection and disease burden that needs to be tackled through sexual health and STI-specific programmes, for both women and men. Although these findings were based on a volume of epidemiological evidence, most studies used convenience

sampling (about 90%) or had unclear response rate (>75%). Meta-regression, however, did not identify an effect for these factors on observed prevalence. A summary of this study and its results in Arabic language can be found in the appendix (p 4).

Although infection prevalence in the population at large suggests active transmission networks for *C trachomatis* and other STIs, it might not necessarily reflect prevalent sexual risk behaviours. This outcome might reflect, at least in part, poor access to and utilisation of STI services—there is very limited capacity in the Middle East and north Africa for STI prevention and treatment, not to mention *C trachomatis* screening and broader sexual health programmes. As observed elsewhere, such as in Alaskan Eskimo populations¹⁹⁴ and populations in South Pacific Islands,¹⁹⁵ poor *C trachomatis* diagnosis and specific treatment can result in unusually high prevalence,^{194,196} probably because *C trachomatis* is largely asymptomatic,⁴ and if untreated, shedding can persist even for years,¹⁹⁷ thereby increasing the potential for reinfection within couples¹⁹⁸ and for transmission in the population.

The high prevalence found in populations at high risk such as female sex workers, in context of evidence suggesting strong partial immunity against reinfection,¹⁹⁹ is consistent with the important role of commercial sex networks in infection transmission. Independent evidence supports existence of hidden pockets of high sexual-risk behaviour driving STI incidence in the Middle East and north Africa.^{10,11} Among male STI patients, 77% in Kuwait²⁰⁰ and 80% in Somalia²⁰¹ reported paying a female sex worker for sex, and among migrant workers in Pakistan 22% reported sex with a female sex worker.²⁰² Higher levels of sexual-risk behaviour and emerging HIV epidemics have been also documented among men who have sex with men, male sex workers, and male-to-female transgenders in systematic reviews.^{17,203} Sexual networks, however, remain poorly investigated in the Middle East and north Africa, owing to cultural sensitivities.

The possible role of *C trachomatis* infection in poor reproductive health outcomes remains unappreciated and neglected by the public health establishment in the Middle East and north Africa, despite substantial social and economic implications for women and their families.^{204,205} A main finding of this study is the high current *C trachomatis* prevalence in infertility clinic attendees, with odds of infection three-times higher than in the general population. By contrast, studies among infertility clinic attendees in Europe usually show that current *C trachomatis* infection is uncommon, but serological evidence of past infection, assumed to have resulted in fallopian tube scarring, is common.^{206–209} This finding suggests a role for *C trachomatis* in infertility in the Middle East and north Africa. Indeed, this region appears to have the highest rate of primary infertility worldwide, which remains unexplained.¹² The Middle East and north Africa is also a region where infertility has multiple




Figure 2: Meta-analysis of studies reporting *Chlamydia trachomatis* current infection prevalence assessed using nucleic acid amplification test in the general population in the Middle East and north Africa. Data are stratified by sex. Error bars are 95% CIs.

detrimental sociocultural consequences,²¹⁰ and where several countries have had rapidly declining fertility rates to even below replacement level.^{211,212} The prevalence of current *C trachomatis* infection was also high in women




Figure 3: Meta-analysis of studies reporting *Chlamydia trachomatis* current infection prevalence assessed using nucleic acid amplification test in infertility clinic attendees in the Middle East and north Africa
Data are stratified by sex. Error bars are 95% CIs.

with miscarriage and in pregnant women—similar to that found in pregnant women in low-income and middle-income countries elsewhere.²¹³⁻²¹⁷ This stigmatised and largely asymptomatic infection might not be visible to the public eye, but its reproductive health sequelae are visible, even if not explicitly linked to the underlying cause.

C trachomatis prevalence in women was higher than in men (two-times higher odds). This difference possibly reflects a longer duration of infection for women, considering that infection in men is more symptomatic²¹⁸ (nearly two-times higher prevalence in symptomatic men than in symptomatic women), and therefore more likely to be treated. Ever infection (anti-*C trachomatis* IgG)

prevalence was two-times higher than current infection prevalence, but the epidemiological relevance of ever infection prevalence might be limited given challenges in *C trachomatis* serology interpretation.²¹⁹

The Middle East and north Africa is burdened by *C trachomatis* infection, but the public health response remains rudimentary and far from achieving WHO's Global Health Sector Strategy on STIs.⁸ Evidence for some differences in *C trachomatis* prevalence by country has been reported, but remarkably, no evidence was found for a variation in prevalence over time (1982-2018). Lingering STI stigma prevents those infected from accessing proper health care, including those most at risk. The role of screening and treatment for asymptomatic *C trachomatis* within established programmes, such as for family planning, primary health care, or HIV, needs careful consideration given the cost and uncertain effect on prevalence at modest levels of uptake.²²⁰

Current STI surveillance focused on inefficient routine case reporting is not capturing the reality of the transmission dynamics.²²¹ Although routine case reporting could be improved with more consistency and universality in reporting and emphasis on aetiological approaches,²²¹ its usefulness for a robust long-term evaluation of infection trends is rather limited. Sentinel surveillance of different at-risk populations should be explored, as recommended by the WHO Global Health Sector Strategy on STIs,⁸ to better identify outbreaks or emerging epidemics, strategically direct resources for prevention, treatment, and control, and monitor and evaluate STI programmes.¹ The recent progress in HIV surveillance in the Middle East and north Africa, in the form of repeated rounds of HIV-integrated biobehavioural surveillance surveys,^{222,223} should be extended to STIs.^{221,224}

Our study has important but unavoidable limitations. Quantity and quality of available data varied by country and population, particularly for populations at high risk where most data came from only a few countries—eg, most studies of men who have sex with men were from Pakistan. No data were identified for Afghanistan, Bahrain, Libya, and Yemen. Prevalence levels might not have been strictly representative and might have been affected by publication bias, as suggested by the small-study effect observed. Studies in women with miscarriage might have included women with induced abortions; however, these were not explicitly indicated, possibly for legal reasons, as abortion is illegal in most of the Middle East and north Africa.^{225,226} The wide array of diagnostics used for ascertainment might have also introduced detection bias.

Factors that might have contributed to differences in *C trachomatis* positivity rates across studies include sampling variation and potential selection bias, spatial or temporal variability in prevalence, and possibly unreported underlying comorbidities. This study did not assess other STIs that might have also contributed to infertility, pregnancy-related morbidity, and other health conditions in women with *C trachomatis* infection. Such

potential biases might have contributed to some of the unexplained heterogeneity observed in the prevalence levels. Given potential limitations in the representativeness of the prevalence measures as well as heterogeneity across studies, the calculated pooled prevalence should be interpreted as a pooled average, rather than strictly

	Studies (n)	Samples (n)	Univariable analyses			Variance explained R ²	Multivariable analysis		
			OR (95% CI)	p value	LR test p value*		Adjusted OR (95% CI)	p value	LR test p value†
At-risk population									
General populations	137	168 302	1.00 (ref)	..	<0.0001	19.0%	1.00 (ref)	..	<0.0001
Populations at intermediate risk	14	4164	0.70 (0.32–1.54)	0.374	1.81 (0.79–4.13)	0.157	..
Female sex workers	20	5679	8.99 (4.57–17.71)	<0.0001	11.28 (5.78–22.01)	<0.0001	..
Men who have sex with men	20	6717	0.83 (0.42–1.64)	0.591	4.16 (1.72–10.08)	0.002	..
Infertility clinic attendees	135	17 891	3.77 (2.67–5.31)	<0.0001	3.39 (2.41–4.77)	<0.0001	..
Women with miscarriage	27	2500	3.53 (1.94–6.40)	<0.0001	2.78 (1.57–4.93)	0.001	..
Women with ectopic pregnancy	3	78	8.25 (1.58–43.08)	0.012	4.93 (1.03–23.52)	0.045	..
Symptomatic women	140	29 402	3.74 (2.66–5.26)	<0.0001	3.47 (2.47–4.87)	<0.0001	..
Symptomatic men	56	22 036	5.76 (3.68–9.03)	<0.0001	7.17 (4.05–12.68)	<0.0001	..
Assay type									
NAAT (current infection)	197	64 083	1.00 (ref)	..	<0.0001	7.1%	1.00 (ref)	..	<0.0001
Culture (current infection)	29	13 536	1.92 (1.05–3.50)	0.034	1.10 (0.62–1.95)	0.742	..
Other (current infection)‡	92	137 924	1.90 (1.30–2.79)	0.001	1.47 (1.02–2.13)	0.041	..
Anti-<i>C trachomatis</i> immunoglobulins§									
IgG (ever infection)	117	14 935	2.99 (2.10–4.26)	<0.0001	2.17 (1.54–3.06)	<0.0001	..
IgM (recent infection)	49	7890	1.17 (0.72–1.90)	0.517	0.90 (0.57–1.40)	0.627	..
IgA	16	3141	0.92 (0.42–2.02)	0.836	0.78 (0.39–1.56)	0.481	..
Not specified (IgG, IgM, or IgA)	29	6146	2.81 (1.54–5.13)	0.001	2.25 (1.28–3.97)	0.005	..
Unclear	23	9114	2.53 (1.30–4.94)	0.007	1.49 (0.81–2.75)	0.200	..
Sampling methodology¶									
Non-probability-based sampling	495	227 208	1.00 (ref)	..	<0.0001	3.5%	1.00 (ref)	..	0.347
Probability-based sampling	57	29 561	0.37 (0.24–0.56)	<0.0001	0.80 (0.50–1.27)	0.347	..
Sample size									
<200	386	32 782	1.00 (ref)	..	<0.0001	6.0%	1.00 (ref)	..	0.001
≥200	166	223 987	0.42 (0.32–0.56)	<0.0001	0.63 (0.48–0.83)	0.001	..
Response rate									
≥80%	112	38 732	1.00 (ref)	..	0.187	0.1%	1.00 (ref)	..	0.237
<80% or unclear	440	218 037	0.80 (0.57–1.12)	0.187	0.83 (0.61–1.13)	0.237	..
Year of publication 									
Year of publication	552	256 769	0.96 (0.95–0.98)	<0.0001	<0.0001	4.4%	0.99 (0.98–1.01)	0.281	0.281
Year of data collection									
Year of data collection	552	256 769	0.96 (0.95–0.98)	<0.0001	<0.0001	4.2%
Country									
Other Middle East or north African countries	245	189 529	1.00 (ref)	..	<0.0001	5.2%	1.00 (ref)	..	0.013
Egypt	89	7434	1.58 (1.08–2.31)	0.018	1.05 (0.73–1.51)	0.774	..
Iran	176	38 647	0.80 (0.59–1.08)	0.145	0.90 (0.68–1.19)	0.472	..
Pakistan	42	21 159	0.31 (0.19–0.52)	<0.0001	0.39 (0.22–0.69)	0.002	..
Sex									
Men	133	42 393	1.00 (ref)	..	0.131	0.2%	1.00 (ref)	..	0.029
Women	419	214 376	1.27 (0.93–1.74)	0.131	1.61 (1.05–2.46)	0.029	..

Adjusted R² in the final multivariable model was 29.0%. LR=likelihood ratio. NAAT=nucleic acid amplification test. OR=Odds ratio. *Predictors with p≤0.2 in the univariable model were considered significant. †Predictors with p≤0.05 in the multivariable model were considered significant. ‡Other assays detecting current infection such as direct fluorescence assays, Giemsa staining, and enzyme-linked immunoassays applied to genital samples. §Includes assays such as enzyme-linked immunoassay and micro-immunofluorescence. ¶Non-probability sampling refers to a sampling method in which the data collection process does not allow individuals to have equal chance of being selected; an example is convenience sampling for which individuals are selected on the basis of ease of accessibility (first-come first-served basis).^{127,128} Probability-based sampling refers to a sampling method in which data collection process is based on a random selection of study participants; an example is random sampling from a sampling frame.¹²⁸ Another example of probability-based sampling is respondent-driven sampling, which is a sampling method specifically designed to sample hard-to-reach populations and is based on chain referral with the probability of selection calculated at each step in the network to produce adjusted prevalence estimates.¹²⁹ ||Only year of publication was considered for the multivariable meta-regression analysis because of collinearity with year of data collection.

Table 4: Results of meta-regressions to identify associations and sources of between-study heterogeneity for *Chlamydia trachomatis* prevalence in the Middle East and north Africa

an estimate of the mean prevalence in the considered population or subpopulation.

In conclusion, *C trachomatis* current infection prevalence in the population at large in the Middle East and north Africa is at 3%, similar to other regions, but higher than expected given these countries' sexually conservative norms. The high prevalence (>10%) in infertility clinic attendees and in women with miscarriage, provides suggestive evidence for the potential role of *C trachomatis* in poor reproductive outcomes in the Middle East and north Africa. In the context of very limited programming for sexual health and STIs, our findings highlight an important, yet neglected and poorly recognised infection and disease burden, despite the social and economic impact. There is an urgent need for targeted and culturally appropriate programmes promoting sexual health for different at-risk populations. Tackling this infection with appropriate interventions is essential to control disease sequelae, to address the WHO Global Health Sector Strategy on STIs,⁸ and to accomplish key health Sustainable Development Goals.

Contributors

AS contributed to the study design, did the systematic searches of the literature, selection of studies for inclusion, and the data extraction and data analyses. HC contributed to the study design, double extracted the data, updated the systematic review, and did the data analyses. AS and HC wrote the first draft of the paper. JGH contributed to identification of unpublished data. NL contributed to the data extraction, analyses, and drafting of the Article. LJA-R conceived and led the design of the study, data extraction, data analyses, and drafting of the Article. All authors contributed to discussion and interpretation of the results and to the writing of the manuscript. All authors have read and approved the final manuscript.

Declaration of interests

We declare no competing interests.

Acknowledgments

This study was funded by the National Priorities Research Program (grant number 9-040-3-008) from the Qatar National Research Fund (a member of Qatar Foundation). Infrastructure support was provided by the Biostatistics, Epidemiology, and Biomathematics Research Core at Weill Cornell Medicine-Qatar, Cornell University, Qatar Foundation—Education City, Doha, Qatar. The statements made herein are solely the responsibility of the authors. The publication of this Article was funded by the Qatar National Library. We thank Adona Canlas and Ashwini Deshmukh for their assistance with locating full-text articles, and Ghina Mumtaz and Vajihah Akbarzadeh for their assistance with translating manuscripts from foreign languages.

References

- WHO. Baseline report on global sexually transmitted infection surveillance 2012. Geneva: World Health Organization, 2013. <http://www.who.int/reproductivehealth/publications/rtis/9789241505895/en/> (accessed March 9, 2014).
- Holmes KK. Sexually transmitted diseases, 4th edn. New York: McGraw-Hill Medical, 2008.
- Newman L, Rowley J, Vander Hoorn S, et al. Global estimates of the prevalence and incidence of four curable sexually transmitted infections in 2012 based on systematic review and global reporting. *PLoS One* 2015; **10**: e0143304.
- WHO. Global prevalence and incidence of selected curable sexually transmitted infections: overview and estimates. Geneva: World Health Organization, 2001.
- Mishori R, McClaskey EL, WinklerPrins VJ. *Chlamydia trachomatis* infections: screening, diagnosis, and management. *Am Fam Physician* 2012; **86**: 1127–32.
- Rekart ML, Gilbert M, Meza R, et al. Chlamydia public health programs and the epidemiology of pelvic inflammatory disease and ectopic pregnancy. *J Infect Dis* 2013; **207**: 30–38.
- UN General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development. 2015. <https://sustainabledevelopment.un.org/content/documents/21252030%20Agenda%20for%20Sustainable%20Development%20web.pdf> (accessed June 6, 2017).
- WHO. Global health sector strategy on sexually transmitted infections, 2016–2021. Geneva: World Health Organization, 2016.
- Abu-Raddad LJ, Ghanem KG, Feizzadeh A, Setayesh H, Calleja JM, Riedner G. HIV and other sexually transmitted infection research in the Middle East and north Africa: promising progress? *Sex Transm Infect* 2013; **89** (suppl 3): iii1–4.
- Abu-Raddad LJ, Hilmi N, Mumtaz G, et al. Epidemiology of HIV infection in the Middle East and north Africa. *AIDS* 2010; **24** (suppl 2): S5–23.
- Abu-Raddad LJ, Akala FA, Semini I, Riedner G, Wilson D, Tawil O. Characterizing the HIV/AIDS epidemic in the Middle East and north Africa: time for strategic action. Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project. Washington DC: The World Bank Press, 2010.
- Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S, Stevens GA. National, regional, and global trends in infertility prevalence since 1990: a systematic analysis of 277 health surveys. *PLoS Med* 2012; **9**: e1001356.
- Higgins JPT, Green S, Cochrane Collaboration. Cochrane handbook for systematic reviews of interventions. Chichester: Wiley-Blackwell, 2008.
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- International AIDS Society. Abstract archives of international aids society conferences. 2016. <http://www.abstract-archive.org/> (accessed July 17, 2017).
- Abu-Raddad LJ, World Bank. Characterizing the HIV/AIDS epidemic in the Middle East and north Africa: time for strategic action. Washington DC: World Bank, 2010.
- Mumtaz G, Hilmi N, McFarland W, et al. Are HIV epidemics among men who have sex with men emerging in the Middle East and north Africa? A systematic review and data synthesis. *PLoS Med* 2010; **8**: e1000444.
- Mumtaz GR, Weiss HA, Thomas SL, et al. HIV among people who inject drugs in the Middle East and north Africa: systematic review and data synthesis. *PLoS Med* 2014; **11**: e1001663.
- Harfouche M, Chemaitelly H, Mahmud S, et al. Epidemiology of hepatitis C virus among hemodialysis patients in the Middle East and north Africa: systematic syntheses, meta-analyses, and meta-regressions. *Epidemiol Infect* 2017; **145**: 3243–63.
- Mahmud S, Al-Kanaani Z, Chemaitelly H, Chaabna K, Kouyoumjian SP, Abu-Raddad LJ. Hepatitis C virus genotypes in the Middle East and north Africa: distribution, diversity, and patterns. *J Med Virol* 2018; **90**: 131–41.
- Harfouche M, Chemaitelly H, Kouyoumjian SP, et al. Hepatitis C virus viremic rate in the Middle East and north Africa: systematic synthesis, meta-analyses, and meta-regressions. *PLoS One* 2017; **12**: e0187177.
- Chemaitelly H, Mahmud S, Chaabna K, Kouyoumjian SP, Mumtaz GR, Abu-Raddad LJ. The epidemiology of hepatitis C virus in the World Health Organization Eastern Mediterranean region: implications for strategic action. 2019. World Health Organization Report (in press).
- El-Kettani A, Mahiane G, Bennani A, et al. Trends in adult chlamydia and gonorrhoea prevalence, incidence and urethral discharge case reporting in Morocco over 1995–2015: estimates using the spectrum-sexually transmitted infection model. *Sex Transm Infect* 2017; **44**: 557–64.
- Smolak A, Rowley J, Nagelkerke N, et al. Trends and predictors of syphilis prevalence in the general population: global pooled analysis of 1103 prevalence measures including 136 million syphilis tests. *Clin Infect Dis* 2018; **66**: 1184–91.
- Korenromp EL, Mahiane G, Rowley J, et al. Estimating prevalence trends in adult gonorrhoea and syphilis in low- and middle-income countries with the spectrum-STI model: results for Zimbabwe and Morocco from 1995 to 2016. *Sex Transm Infect* 2017; **93**: 599–606.

- 26 Freeman MF, Tukey JW. Transformations related to the angular and the square root. *Ann Math Stat* 1950; **21**: 607–11.
- 27 Miller JJ. The inverse of the Freeman–Tukey double arcsine transformation. *Am Stat* 1978; **32**: 138.
- 28 Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta-analysis of prevalence. *J Epidemiol Community Health* 2013; **67**: 974–78.
- 29 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; **7**: 177–88.
- 30 Borenstein M. Introduction to meta-analysis. Chichester: John Wiley & Sons, 2009.
- 31 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539–58.
- 32 R core team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing, 2017.
- 33 UN Population Division. World population prospects 2017. 2018. <https://population.un.org/wpp/> (accessed July 10, 2018).
- 34 StataCorp. Stata statistical software: release 14. College Station: StataCorp LP, 2014.
- 35 Kadi Z, Bouguermouh A, Djenaoui T, Allouache A, Dali S, Hadji N. Chlamydial genital infection in Algiers: a sero-epidemiological survey. *Trans R Soc Trop Med Hyg* 1990; **84**: 863–65.
- 36 Abdel Monem AA, Bassyouni MI, Soliman MY, Ahmed SR, Abdel Hafeez HA. *Chlamydia trachomatis* antigen detection in female infertility. *El-Minia Med Bull* 2005; **16**: 104–12.
- 37 Aboul Atta HNE, Ibrahim AA. Role of chlamydia and mycoplasma in the etiology of non-gonococcal urethritis in men. *Egypt J Med Microbiol* 1995; **4**: 355–60.
- 38 Badary MS. Study of the role of cervical chlamydial infection in unexplained infertility and mucopurulent cervicitis. *Egypt J Med Microbiol* 1996; **5**: 431–38.
- 39 Berry ME, El Shabrawy A. *Chlamydia trachomatis* infection and relation to female infertility. *Egypt J Med Microbiol* 1996; **5**: 297–304.
- 40 Diab KM. Gonococcal and chlamydial antibodies in Egyptian women with ectopic pregnancy. *New Egypt J Med* 1993; **8**: 1006–10.
- 41 Draz EI, Hassan AM, Khalil HS, Elomary MA. Evaluation of pelvic inflammatory disease potential in cholinesterase inhibitor pesticide-exposed females. *Environ Sci Pollut Res Int* 2018; **25**: 30818–25.
- 42 El-Sayed N, Abdallah M, Abdel Mobdy A, et al. Evaluation of selected reproductive health infections in various Egyptian population groups in Greater Cairo. Cairo: Egypt Ministry of Health and Population, 2002.
- 43 Mosbah A, Nabiel Y. *Helicobacter pylori*, *Chlamydiae pneumoniae* and *trachomatis* as probable etiological agents of preeclampsia. *J Matern Fetal Neonatal Med* 2016; **29**: 1607–12.
- 44 Mousa A. The association between *Chlamydia trachomatis* and cervical intra-epithelial neoplasia. *Zagazig Med Assoc J* 1990; **3**: 41–46.
- 45 Nada AM, Hassan FM, Al-Azhary NH. Detection of *Chlamydia trachomatis* in patients with unexplained infertility: a case control study. *Egypt J Med Microbiol* 2015; **24**: 35–38.
- 46 Sullam SA, Mahfouz AA, Dabbous NI, el-Barrawy M, el-Said MM. Reproductive tract infections among married women in Upper Egypt. *East Mediterr Health J* 2001; **7**: 139–46.
- 47 Zaki SA. Prevalence of endocervical genital mycoplasmas and *Chlamydia trachomatis* in infertile, abortive and pregnant women in Alexandria. *Bull Alex Fac Med* 1989; **25**: 1031–44.
- 48 Ahmadi A, Khodabandehloo M, Ramazanzadeh R, et al. The relationship between *Chlamydia trachomatis* genital infection and spontaneous abortion. *J Reprod Infertil* 2016; **17**: 110–16.
- 49 Ahmadi MH, Mirsalehian A, Sadighi Gilani MA, Bahador A, Afraz K. Association of asymptomatic *Chlamydia trachomatis* infection with male infertility and the effect of antibiotic therapy in improvement of semen quality in infected infertile men. *Andrologia* 2018; **50**: e12944.
- 50 Ahmadnia E, Kharaghani R, Maleki A, et al. Prevalence and associated factors of genital and sexually transmitted infections in married women of Iran. *Oman Med J* 2016; **31**: 439–45.
- 51 Badami N, Salari MH. Rate of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* in infertile females and control group. *Iran J Public Health* 2001; **30**: 57–60.
- 52 Baghchesaraei H, Amini B, Hossaini M. Prevalence of infection with *Neisseria gonorrhoeae* and *Chlamydia trachomatis* in women visitors of gynecology and obstetrics clinics in Zanjan Province of Iran. *Afr J Microbiol Res* 2011; **5**: 2447–50.
- 53 Bagheri S, Roghanian R, Golbang N, Golbang P, Esfahani MHN. Molecular evidence of *Chlamydia trachomatis* infection and its relation to miscarriage. *Int J Fertil Steril* 2018; **12**: 152–56.
- 54 Behroozi R. Epidemiology of *Chlamydia trachomatis* infections in pregnant women of the hospitals of Tehran University of Medical Sciences. *J Nurs Midwifery Sci* 2001; **1**: 14–20.
- 55 Chamani-Tabriz L, Jedi TM, Zeraati H, et al. A molecular survey of *Chlamydia trachomatis* infection in married women: a cross sectional study on 991 women. *Tehran Univ Med J* 2008; **66**: 498–91.
- 56 Cheraghi M, Rahimi Z, Parsa S. Prevalence of cervical-vaginal infections in the pap-smear samples in Iran. *Glob J Health Sci* 2014; **16**: 201–06.
- 57 Dehghan ML, Aflatoonian A, Talebi AR, Eley A, Pacey AA. Relationship between *Chlamydia trachomatis* and *Mycoplasma genitalium* infection and pregnancy rate and outcome in Iranian infertile couples. *Andrologia* 2017; **49**.
- 58 Goshayeshi L, Vahid Roudsari F, Ghazvini K, Nomani H, Amel Jamehdar S. Pilot prevalence of *Chlamydia trachomatis* by PCR in female infertile referred to study center of infertility in Mashhad. *Iran South Med J* 2015; **18**: 92–99.
- 59 Haghighi Hasanabad M, Bahador A, Mohammadzadeh M, Haghighi F. Prevalence of *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Ureaplasma urealyticum* in pregnant women of Sabzevar, Iran. *Sex Transm Infect* 2013; **89** (suppl 1): A233–34.
- 60 Jahromi AS, Farjam MR, Mogharrab F, et al. *Chlamydia trachomatis* in women with full-term deliveries and women with abortion. *Am J Infect Dis* 2010; **6**: 66–69.
- 61 Javanmard D, Behravan M, Ghannadkafi M, Salehabadi A, Ziaee M, Namaei MH. Detection of *Chlamydia trachomatis* in pap smear samples from south Khorasan province of Iran. *Int J Fertil Steril* 2018; **12**: 31–36.
- 62 Joolayi F, Navidifar T, Jaafari RM, Amin M. Comparison of *Chlamydia trachomatis* infection among infertile and fertile women in Ahvaz, Iran: a case-control study. *Int J Reprod Biomed (Yazd)* 2017; **15**: 713–18.
- 63 Kajbaf MJ, Gholamnezhad A. Prevalence of *Chlamydia trachomatis* antigen and antibody in infertile women in Ahwaz. *Iran Biomed J* 1998; **2**: 45–48.
- 64 Kamyabi Z. The survey of anti *Chlamydia* antibody in infertile women undergoing diagnostic laparoscopy in Afzalipour medical centre, Kerman—Iran. *Int J Gynaecol Obstet* 2009; **107**: S218.
- 65 Khezerdoust S, Hagh Elahi F, Roustaei S, Badami N, Naghizadeh MM, Jafarabadi M. *Chlamydia trachomatis* infection in pregnant women. *J Reprod Infertil* 2009; **10**: 121–28.
- 66 Marashi SM, Moulana Z, Imani Fooladi AA, Mashhadi Karim M. Comparison of genital *Chlamydia trachomatis* infection incidence between women with infertility and healthy women in Iran using PCR and immunofluorescence methods. *Jundishapur J Microbiol* 2014; **7**: e9450.
- 67 Farhoudi B, Kamali K, Rajabpoor Z. Situation analysis of sexually transmitted infections in the Islamic Republic of Iran 2008. Tehran: Ministry of Health and Medical Education, 2008.
- 68 Meidani M. Frequency of *Chlamydia trachomatis* infection in asymptomatic males based on PCR in Tehran, Iran. *Int Urogynecol J Pelvic Floor Dysfunct* 2009; **20** (suppl 3): S460–61.
- 69 Mousavi A, Ramezanzadeh R, Farhadifar F, et al. Detection of *Chlamydia trachomatis* in fertile and infertile women in Sanandaj by PCR. *Iran J Public Health* 2014; **43**: 63.
- 70 Pourabbas B, Rezaei Z, Mardaneh J, Shahian M, Alborzi A. Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections among pregnant women and eye colonization of their neonates at birth time, Shiraz, Southern Iran. *BMC Infect Dis* 2018; **18**: 477.
- 71 Rashidi BH, Chamani-Tabriz L, Haghollahi F, et al. Effects of *Chlamydia trachomatis* infection on fertility; a case-control study. *J Reprod Infertil* 2013; **14**: 67–72.
- 72 Rohi E, Ghasemi K, Kahnemoui Agdam F. Incidence of non-gonococcal infection in childbearing and pregnant women in Ardabil. *Int J Mol Clin Microbiol* 2011; **1**: 71–76.
- 73 Rostami MN, Rashidi BH, Aghsaghloo F, Nazari R. Comparison of clinical performance of antigen-based enzyme immunoassay (EIA) and major outer membrane protein (MOMP)-PCR for detection of genital *Chlamydia trachomatis* infection. *Int J Reprod Biomed (Yazd)* 2016; **14**: 411–20.

- 74 Safdari H, Safdari A, Tahaghoghi S, Yari A, Ghazvini K. Prevalence of *Chlamydia trachomatis* among women with genital infection in northeast of Iran in 2013. *Iran J Obstet Gynecol Infertol* 2015; **18**: 1–6.
- 75 Sattari M, Ghiami Rad M, Ghasemzadeh A, Mohammadoghli Reihan Z. Frequency of anti-*Chlamydia trachomatis* antibodies in infertile women referred to Tabriz Al-Zahra hospital. *Int J Reprod Biomed (Yazd)* 2017; **15**: 17–20.
- 76 Sisakht AJ, Omidifard N, Mohamadkhani N, Karimpoorfard M, Kargar M, Shokripour M. Assessing the presence of *Chlamydia trachomatis* genome in pregnant women with spontaneous abortion using polymerase chain reaction method in Yasuj: first report from southwest of Iran. *J Educ Health Promot* 2017; **6**: 45.
- 77 Yeganeh O, Jjeddi-Tehrani M, Yaghmaie F, et al. A survey on the prevalence of *Chlamydia trachomatis* and *Mycoplasma genitalium* infections in symptomatic and asymptomatic men referring to urology clinic of Labbafnejad Hospital, Tehran, Iran. *Iran Red Crescent Med J* 2013; **15**: 340–44.
- 78 Zahirnia Z, Mansouri S, Saffari F, Mansouri G. Frequency of *Chlamydia trachomatis*, *Mycoplasma genitalium*, and *Ureaplasma urealyticum* isolated from vaginal samples of women in Kerman, Iran. *Arch Clin Infect Dis* 2018; **13**: e65334.
- 79 Abdulkhudher NA, Yousif MG, Sadiq AM. Detection of immunological markers for *Chlamydia trachomatis* and *Trichomonas vaginalis* infection in women with obstetric complications in Najaf, Iraq. *J Al-Qadisiyah Pure Sci* 2014; **19**: 1–25.
- 80 Abdul-Karim ET, Abdul-Muhyemen N, Al-Saadie M. *Chlamydia trachomatis* and rubella antibodies in women with full-term deliveries and women with abortion in Baghdad. *East Mediterr Health J* 2009; **15**: 1407–11.
- 81 Abdullah FT. Study the role of *Chlamydia trachomatis* in ectopic pregnancy in Iraqi women. *J Biotechnol Res Cent* 2012; **6**: 19–25.
- 82 Ahmed DW. Effects of interleukin-2 (IL-2) and interleukin-6 (IL-6) in recurrent spontaneous abortion (RSA). *Iraqi J Pharm Sci* 2008; **17**: 74–79.
- 83 Al-Hamdani SM, Shemran HA, Kadhim HS, Yasser H. Prevalence of anti-chlamydial immunoglobulins in pregnant women in Al-Kadhimiya Teaching Hospital. *J Coll Teach* 2010; **1**: 59–70.
- 84 Al-Husseinei RK, Kadaem KM, Al-Janabi LA, Fathei MiM. ELFA and IFAT techniques to detect chlamydial infections in Baghdad women and its effect on the immunoglobulins level. *J Facult Med* 2009; **51**: 192–97.
- 85 Ali MH, Al-Kazaz AA. Molecular detection of *Chlamydia trachomatis* infection among males with abnormal semen. *Iraqi J Sci* 2018; **59**: 2005–11.
- 86 Alkhafaf D. Prevalence of *Chlamydia trachomatis* in women of Al-Diwaniya city. *J Babylon Univ* 2013; **21**: 1232–38.
- 87 Hwaid AH, Hasan A-RS, Al-Duliami AA. Seroprevalence of anti-*C trachomatis* IgG and IgM antibodies among pregnant women in Diyala province. *J Facult Med Baghdad* 2013; **55**: 348–51.
- 88 Ismail MK, Ali AS. Evaluation of *Chlamydia trachomatis* antibodies in women with infertility. *Al-Mustansiriyah J Sci* 2012; **23**: 21–28.
- 89 Mohammed NAM, Salman AH, Hasan FK. *Chlamydia trachomatis* and recurrent spontaneous abortion in Iraqi pregnant women. *Iraqi J Med Sci* 2012; **10**: 42–46.
- 90 Mohammed IH, Al-Awadei SJ, Saadedin SMK. Molecular diagnosis of *Chlamydia trachomatis* in infertile Iraqi women using real time-PCR and comparison with other methods. *Iraqi J Sci* 2017; **58**: 1437–46.
- 91 Yahya M, Al-Siraj D. Study of genital *Mycoplasma* and *Chlamydia trachomatis* which presence in infertile males. *Al-Taqani* 2009; **22**: A23–31.
- 92 Abusarah EA, Awwad ZM, Charvalos E, Shehabi AA. Molecular detection of potential sexually transmitted pathogens in semen and urine specimens of infertile and fertile males. *Diagn Microbiol Infect Dis* 2013; **77**: 283–86.
- 93 Al-Ramahi M, Mahafzah A, Saleh S, Fram K. Prevalence of *Chlamydia trachomatis* infection in infertile women at a university hospital in Jordan. *East Mediterr Health J* 2008; **14**: 1148–54.
- 94 As'ad A. National AIDS program, final report sexually transmitted infections (STI) prevalence study, Jordan. 2004.
- 95 Awwad ZM, Al-Amarat AA, Shehabi AA. Prevalence of genital chlamydial infection in symptomatic and asymptomatic Jordanian patients. *Int J Infect Dis* 2003; **7**: 206–09.
- 96 Mahafzah AM, Al-Ramahi MQ, Asa'd AM, El-Khateeb MS. Prevalence of sexually transmitted infections among sexually active Jordanian females. *Sex Transm Infect* 2008; **35**: 607–10.
- 97 Jordan Ministry of Health. Prevalence of reproductive tract infections in women attending selected urban OB/GYN clinics in Jordan. 2004.
- 98 Al-Awadhi R, Al-Shaheen A, Al-Juwaiser A, George SS, Sharma P, Kapila K. Prevalence of infectious organisms observed in cervical smears between 1997–2014 at Mubarak Al-Kabeer Hospital, Kuwait. *Sultan Qaboos Univ Med J* 2018; **18**: e324–28.
- 99 Al-Sweih NA, Khan S, Rotimi VO. Prevalence of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* among asymptomatic women attending the Capital Health region clinics in Kuwait. *Sex Transm Infect* 2011; **38**: 793–97.
- 100 Al-Sweih NA, Al-Fadli AH, Omu AE, Rotimi VO. Prevalence of *Chlamydia trachomatis*, *Mycoplasma hominis*, *Mycoplasma genitalium*, and *Ureaplasma urealyticum* infections and seminal quality in infertile and fertile men in Kuwait. *J Andrology* 2012; **33**: 1323–29.
- 101 Deeb ME, Awwad J, Yeretizian JS, Kaspar HG. Prevalence of reproductive tract infections, genital prolapse, and obesity in a rural community in Lebanon. *Bull World Health Organ* 2003; **81**: 639–45.
- 102 Hancali A, Bellaji B, Jennane S, et al. Trend of STIs prevalence among women and men in Morocco between 1999 to 2011. *Sex Transm Infect* 2015; **91**: A155.
- 103 Hulstein SH, Matser A, Alberts CJ, et al. Differences in *Chlamydia trachomatis* seroprevalence between ethnic groups cannot be fully explained by socioeconomic status, sexual healthcare seeking behavior or sexual risk behavior: a cross-sectional analysis in the Healthy Life in an Urban Setting (HELIUS) study. *BMC Infect Dis* 2018; **18**: 612.
- 104 Morocco Ministry of Health. Etude de prévalence IST chez les femmes consultantes en SMI/PF à la Wilaya de Rabat, Rapport final, Programme National de lutte contre les IST/SIDA. Rabat: Morocco Ministry of Health, 2001.
- 105 Abu-Raddad LJ. Additional data points from the Middle East and North Africa HIV/AIDS Epidemiology Synthesis Project database. Tehran: Infectious Disease Epidemiology Group, 2017.
- 106 Radouani F, Takourt B, Ibrahimy S, Sekkat S, Guinet R, Benslimane A. Contribution of *Chlamydia trachomatis* infection to infertility. *Rev Fr Gynecol Obstet* 1998; **93**: 442–46.
- 107 Takourt B, Radouani F, Bencheikroun A, et al. Seroprevalence of *Chlamydia trachomatis* infection in STD consultants in Morocco. *Bull Soc Pathol Exot* 1995; **88**: 194–98.
- 108 Mir AM, Wajid A, Reichenbach L, Khan M. STI prevalence and associated factors among urban men in Pakistan. *Sex Transm Infect* 2009; **85**: 199–200.
- 109 Wasti S, Ashfaq MK, Ishaq R, Hamid R. Prevalence of chlamydial infection in females attending antenatal and family planning clinics in Karachi Pakistan. *Aust N Z J Obstet Gynaecol* 1997; **37**: 462–65.
- 110 Al-Thani A, Abdul-Rahim H, Alabsi E, et al. Prevalence of *Chlamydia trachomatis* infection in the general population of women in Qatar. *Sex Transm Infect* 2013; **89** (suppl 3): iii57–60.
- 111 Alzahrani AJ, Obeid OE, Hassan MI, Almulhim AA. Screening of pregnant women attending the antenatal care clinic of a tertiary hospital in eastern Saudi Arabia for *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections. *Indian J Sex Transm Dis AIDS* 2010; **31**: 81–86.
- 112 Awad NS, Said MM, Mohamed AA, El-Tarras AE. Detection of some sexually transmitted bacterial infection using molecular genetic technique. *World J Med Sci* 2013; **9**: 142–46.
- 113 Bashi SA. Chlamydial infection in Riyadh, Saudi Arabia: a sero-epidemiological survey. *Saudi Med J* 1987; **4**: 387–90.
- 114 Ghazi HO, Daghestani MH, Mohamed MF. Seropositivity of *Chlamydia trachomatis* among Saudi pregnant women in Makkah. *J Family Community Med* 2006; **13**: 61–64.
- 115 Hossain A. Serologic diagnosis of *Chlamydia trachomatis* infections. *Int J Gynaecol Obstet* 1988; **27**: 377–80.
- 116 Kamel RM. Screening for *Chlamydia trachomatis* infection among infertile women in Saudi Arabia. *Int J Womens Health* 2013; **5**: 277–84.
- 117 Massoud M, Noweir A, Salah M, Saleh WA. Chlamydial infection in Riyadh, Saudi Arabia. *J Egypt Public Health Assoc* 1991; **66**: 411–19.
- 118 Ismail SO, Ahmed HJ, Jama MA, et al. Syphilis, gonorrhoea and genital chlamydial infection in a Somali village. *Genitourin Med* 1990; **66**: 70–75.

- 119 Nur YA, Groen J, Elmi AM, Ott A, Osterhaus AD. Prevalence of serum antibodies against bloodborne and sexually transmitted agents in selected groups in Somalia. *Epidemiol Infect* 2000; **124**: 137–41.
- 120 WHO. Central South: the 2004 first national second generation HIV/AIDS/STI sentinel surveillance survey. Geneva: World Health Organization, 2005a.
- 121 WHO. The 2004 first national second generation HIV/AIDS/STI sentinel surveillance survey. Geneva: World Health Organization, 2005b.
- 122 Ahmed MA, Hassan NG, Omer ME, Rostami A, Rayis DA, Adam I. *Helicobacter pylori* and *Chlamydia trachomatis* in Sudanese women with preeclampsia. *J Matern Fetal Neonatal Med* 2018; published online Oct 14. DOI:10.1080/14767058.2018.1536738.
- 123 Almroth L, Elmusharaf S, El Hadi N, et al. Primary infertility after genital mutilation in girlhood in Sudan: a case-control study. *Lancet* 2005; **366**: 385–91.
- 124 Ortashi OM, El Khidir I, Herieka E. Prevalence of HIV, syphilis, *Chlamydia trachomatis*, *Neisseria gonorrhoea*, *Trichomonas vaginalis* and candidiasis among pregnant women attending an antenatal clinic in Khartoum, Sudan. *J Obstet Gynaecol* 2004; **24**: 513–15.
- 125 Alkayer SM, Yazji HN, Khamis A. The prevalence of serum IgG antibody to *Chlamydia trachomatis* in infertile women at the university hospital, Syria. *Res J Pharm Technol* 2017; **10**: 1373–75.
- 126 Ghazal-Aswad S, Badrinal P, Osman NA, Abdul-Khalik S, Raasclou T. Is there a correlation between vaginal chlamydia infection and cervical smear abnormalities? A community-based study in the Al-Ain district, United Arab Emirates. *J Obstet Gynaecol Res* 2006; **32**: 63–67.
- 127 Explorable. Non-probability sampling. 2009. <https://explorable.com/non-probability-sampling> (accessed April 10, 2019).
- 128 Glen S. Probability sampling: definition, types, advantages and disadvantages. 2015. <https://www.statisticshowto.datasciencecentral.com/probability-sampling/> (accessed April 10, 2019).
- 129 Johnston LG. Respondent driven sampling (RDS). 2014. <http://www.lisagjohnston.com/respondent-driven-sampling> (accessed April 10, 2019).
- 130 Darougar S, Aramesh B, Gibson JA, Treharne JD, Jones BR. Chlamydial genital infection in prostitutes in Iran. *Br J Vener Dis* 1983; **59**: 53–55.
- 131 Kassaian N, Ataei B, Yaran M, Babak A, Shoaie P, Ataie M. HIV and other sexually transmitted infections in women with illegal social behavior in Isfahan, Iran. *Adv Biomed Res* 2012; **1**: 5.
- 132 Kazerooni PA, Motazedian N, Motamedifar M, et al. The prevalence of human immunodeficiency virus and sexually transmitted infections among female sex workers in Shiraz, south of Iran: by respondent-driven sampling. *Int J STD AIDS* 2014; **25**: 155–61.
- 133 Mirzazadeh ASM, Khajehkazemi R, Hosseini Hooshyar S, et al. HIV and sexually transmitted infections among female sex workers in Iran: findings from the 2010 and 2015 national surveillance surveys. International AIDS Conference; Durban, South Africa; July 18–22, 2016.
- 134 Bellaji B, Hancali A, Jennane S, et al. Prevalence of *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Trichomonas vaginalis* in female sex workers in Morocco. *Sex Transm Infect* 2017; **93** (suppl 2): A100.
- 135 Morocco Ministry of Health. Etude de prévalence des IST chez les femmes qui consultent pour pertes vaginales et/ou douleurs du bas ventre. Rabat: Morocco Ministry of Health, 2008.
- 136 Morocco Ministry of Health. HIV integrated behavioral and biological surveillance surveys. Rabat: Morocco Ministry of Health, 2011.
- 137 Morocco Ministry of Health. Integrated behavioral and biological surveillance survey among men who have sex with men (MSM) in Agadir, Casablanca, Marrakech, and Tangier. Rabat: Morocco Ministry of Health, 2015.
- 138 Hawkes S, Collumbien M, Platt L, et al. HIV and other sexually transmitted infections among men, transgenders and women selling sex in two cities in Pakistan: a cross-sectional prevalence survey. *Sex Transm Infect* 2009; **85** (suppl 2): ii8–16.
- 139 Khan MS, Unemo M, Zaman S, Lundborg CS. HIV, STI prevalence and risk behaviours among women selling sex in Lahore, Pakistan. *BMC Infect Dis* 2011; **11**: 119.
- 140 Osama M. Rate of STI among selected MSM and transgenders in 5 cities of Pakistan. *Sex Transm Infect* 2017; **93** (suppl 2): A18.
- 141 Rehan N, Bokhari A, Nizamani NM, et al. National study of reproductive tract infections among high risk groups of Lahore and Karachi. *J Coll Physicians Surg Pak* 2009; **19**: 228–31.
- 142 Znazen A, Frikha-Gargouri O, Berrajah L, et al. Sexually transmitted infections among female sex workers in Tunisia: high prevalence of *Chlamydia trachomatis*. *Sex Transm Infect* 2010; **86**: 500–05.
- 143 Abdel Aleem H, Moubasher AED, Abdel Megeed E. Seroprevalence of *Chlamydia trachomatis* among infertile couples in Assiut. *Ain-Shams Med J* 1996; **47**: 759–67.
- 144 Abdella RM, Abdelmoaty HI, Elsherif RH, et al. Screening for *Chlamydia trachomatis* in Egyptian women with unexplained infertility, comparing real-time PCR techniques to standard serology tests: case control study. *BMC Womens Health* 2015; **15**: 45.
- 145 Azab H, Hassouna AA. Evaluation of clinical findings and chlamydia antibody titre as predictors of tubal factor infertility. *Med J Cairo Univ* 2008; **76**: 171–75.
- 146 Elkayal NM, Mahmoud NF, Abdalla S. Detection of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* in Egyptian women suffering from infertility. *Adv Microbiol* 2015; **5**: 769–79.
- 147 El Sayed AF, Al Bakry MM, Zaitoun MM, AbouTalib AF. Comparative study of hysterosalpingography and chlamydia antibody testing in predicting tubal factor infertility. *Ain-Shams Med J* 1997; **48**: 815–25.
- 148 Inhorn MC, Buss KA. Infertility, infection, and iatrogenesis in Egypt: the anthropological epidemiology of blocked tubes. *Med Anthropol* 1993; **15**: 217–44.
- 149 Makled AK, Elkady OS, Swedan KH, Sammour HM, Mohamed EA. Relationship between serum *Chlamydia trachomatis* antibody titer and tubal block in infertile Egyptian women. *Middle East Fertil Soc J* 2013; **18**: 38–41.
- 150 Sadek S, Abo Gabal A, El Hefnawi N. Chlamydial infection and sperm antibodies. *New Egypt J Med* 1993; **8**: 1461–63.
- 151 Siam EM, Hefzy EM. The relationship between antisperm antibodies prevalence and genital *Chlamydia trachomatis* infection in women with unexplained infertility. *Middle East Fertil Soc J* 2012; **17**: 93–100.
- 152 Younis HA, El Sayed SA, Abdel Wahab A, Ayoub AA. Detection of *Chlamydia trachomatis* DNA by polymerase chain reaction in urine of women with tubal infertility as diagnosed by hysterosalpingography and laparoscopy. *Sci J Al-Azhar Med Fac* 2000; **21**: 731–44.
- 153 Zaitun MM, Zaitoun A. *Chlamydia trachomatis* infection in cases of infertility due to tubal factor: a serological study. *Zagazig Med Assoc J* 1990; **3**: 125–33.
- 154 Zytoon MM. The role of *Chlamydia trachomatis* infection in female infertility. *New Egypt J Med* 1994; **11**: 938–42.
- 155 Golshani M, Eslami G, Ghobadloo SM, et al. Detection of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* by multiplex PCR in semen sample of infertile men. *Iran J Public Health* 2007; **36**: 50–57.
- 156 Hajikhani B, Motallebi T, Norouzi J, et al. Classical and molecular methods for evaluation of *Chlamydia trachomatis* infection in women with tubal factor infertility. *J Reprod Infertil* 2013; **14**: 29–33.
- 157 Kalantar SM, Kazemi MJ, Sheikhha MH, Aflatoonian A, Kafizadeh F. Detection of *Chlamydia trachomatis* infection in female partners of infertile couples. *Iran J Fertil Steril* 2007; **1**: 79–84.
- 158 Mansour Ghanaie M. Relative frequency of *Chlamydia trachomatis* infection in infertile women. *Int J Fertil Steril* 2014; **8**: 110.
- 159 Moazenchi M, Totonchi M, Salman Yazdi R, et al. The impact of *Chlamydia trachomatis* infection on sperm parameters and male fertility: a comprehensive study. *Int J STD AIDS* 2018; **29**: 466–73.
- 160 Nan Bakhsh F, Tarvati MR, Yekta Z, Gol MS, Mehrzad Sadaghiani M, Shokri S. Evaluation of the prognostic value of *Chlamydia trachomatis* antibody testing (CAT), HSG and laparoscopy in predicting tubal factor infertility (TFI) in Motahary Hospital, Urmiah, Iran. *Med J Tabriz Univ Med Sci* 2008; **30**: 119–25.
- 161 Nikbakt R, Saharkhiz N, Ghalambor Dezfouli F. Comparison of level of antibodies against *Chlamydia trachomatis* in infertile women due to tubal factors and fertile women. *J Shahid Sadoughi Univ Med Sci Health Serv* 2008; **16**: 10–15.
- 162 Peivandi S, Moslemzadeh N, Gharajeh S, Ajami A. The role of *Chlamydia trachomatis* IgG antibody testing in predicting tubal factor infertility in northern Iran. *Int J Fertil Steril* 2009; **3**: 143–48.

- 163 Rashidi B, Shahrokh Tehraninezhad E, Alaie F. Prevalence of antichlamydia antibody status in infertile patients. *Int J Fertil Steril* 2007; **1**: 15–18.
- 164 Sadrpou P, Bahador A, Asgari S, Bagheri R, Chamani-Tabriz L. Detection of *Chlamydia trachomatis* and *Mycoplasma genitalium* in semen samples of infertile men using multiplex PCR. *Tehran Univ Med J* 2013; **70**: 623–29.
- 165 Siahkali AP, Amiri K. Identification of herpes simplex virus, *Chlamydia trachomatis*, and *Mycoplasma genitalium* in infertile seminal fluid samples using multiplex-PCR in Kerman province, Iran (2016). *Iran J Obstet Gynecol Infertil* 2018; **21**: 69–74.
- 166 Abid A-AJ, Al-Zwaid AJ. Chlamydial antibodies, pro-inflammatory cytokines and bacterial significance among infertile women. *Med J Babylon* 2015; **12**: 943–47.
- 167 Ahmed ST. Detection of *Chlamydia trachomatis* using polymerase chain reaction (PCR). *Al-Mustansiriyah J Sci* 2012; **23**: 35–42.
- 168 Al-Kattan SAA-K, Mohammed RA. Laparoscopic finding and *Chlamydia trachomatis* infection in infertile women. *Med J Tikrit* 2013; **19**: 315–24.
- 169 Dawood SD. *Chlamydia trachomatis* in Iraqi women with infertility. *Iraqi J Biotechnol* 2011; **10**: 89–94.
- 170 Al Subhi T, Al Jashnmi RN, Al Khaduri M, Gowri V. Prevalence of tubal obstruction in the hysterosalpingogram of women with primary and secondary infertility. *J Reprod Infertil* 2013; **14**: 214–16.
- 171 Qayum M, Khalid-bin-Saleem M. Prevalence of *Chlamydia trachomatis* among asymptomatic women. *J Ayub Med Coll Abbottabad* 2013; **25**: 28–30.
- 172 Al-Hindi A, Al-Helou T, Al-Helou Y. Seroprevalence of *Toxoplasma gondii*, cytomegalovirus, rubella virus and *Chlamydia trachomatis* among infertile women attending in vitro fertilization center, Gaza strip, Palestine. *J Egypt Soc Parasitol* 2010; **40**: 451–58.
- 173 Abdul Jabbar HS. Chlamydial infection in Saudi infertile patients. *Bahrain Med Bull* 1990; **12**: 4–7.
- 174 Alfarrag DA, Somily AM, Alssum RM, Abotalib ZM, El-Sayed AA, Al-Mandee HH. The prevalence of *Chlamydia trachomatis* infection among Saudi women attending the infertility clinic in Central Saudi Arabia. *Saudi Med J* 2015; **36**: 61–66.
- 175 Sabra SMM, Al-Harbi MS. An influential relationship of seminal fluid microbial infections and infertility, Taif Region, KSA. *World J Med Sci* 2014; **10**: 32–37.
- 176 Gdoura R, Daoudi F, Bouzid F, et al. Detection of *Chlamydia trachomatis* in semen and urethral specimens from male members of infertile couples in Tunisia. *Eur J Contracept Reprod Health Care* 2001; **6**: 14–20.
- 177 Gdoura R, Keskes-Ammar L, Bouzid F, Eb F, Hammami A, Orfila J. *Chlamydia trachomatis* and male infertility in Tunisia. *Eur J Contracept Reprod Health Care* 2001; **6**: 102–07.
- 178 Gdoura R, Kchaou W, Ammar-Keskes L, et al. Assessment of *Chlamydia trachomatis*, *Ureaplasma urealyticum*, *Ureaplasma parvum*, *Mycoplasma hominis*, and *Mycoplasma genitalium* in semen and first void urine specimens of asymptomatic male partners of infertile couples. *J Andrology* 2008; **29**: 198–206.
- 179 Sellami H, Znazen A, Sellami A, et al. Molecular detection of *Chlamydia trachomatis* and other sexually transmitted bacteria in semen of male partners of infertile couples in Tunisia: the effect on semen parameters and spermatozoa apoptosis markers. *PLoS One* 2014; **9**: e98903.
- 180 Ahmadi A, Khodabandehloo M, Ramazanadeh R, et al. The relationship between *Chlamydia trachomatis* genital infection and spontaneous abortion. *J Reprod Infertil* 2016; **17**: 110–16.
- 181 Bagheri S, Roghanian R. Evaluation of *Chlamydia trachomatis* incidence in women who had abortion in Isfahan. *Iran J Public Health* 2014; **43**: 63.
- 182 Massiha A, Khoshkholgh Pahlaviani MRM, Sedighi Kasmaie P, Vand Yousefi J. Study of the reasons of infectious abortions in the subjects who approached Kamali Hospital, Karaj in Iran. *J Reprod Med Endocrinol* 2010; **7**: 346.
- 183 Salari MH, Badami N. The rate of *Chlamydia trachomatis*, *Mycoplasma hominis* and *Ureaplasma urealyticum* in females and habitual abortion. *ACTA Medica Iranica* 2002; **40**: 79–82.
- 184 Al-Nuaimy WA, Al-Jandeel T. Immunological and molecular study of *Chlamydia trachomatis* as causative agent of abortion in Al-Muthanna province. *Iraqi J Vet Sci* 2018; **42**: 99–104.
- 185 Salman YJ. Chlamydia trachomatous antibodies cross reaction with seropositive *Toxoplasma gondii* and cytomegalovirus among women with abortion and outcomes of congenital abnormalities in Kirkuk City. *Tikrit J Pure Sci* 2016; **21**: 1–5.
- 186 Harbord RM, Higgins JPT. Meta-regression in Stata. *Stata J* 2008; **8**: 493–519.
- 187 Dargham SR, Nasrallah GK, Al-Absi ES, et al. Herpes simplex virus type 2 seroprevalence among different national populations of Middle East and north African men. *Sex Transm Infect* 2018; **45**: 482–87.
- 188 Abu-Raddad LJ, Schiffer JT, Ashley R, et al. HSV-2 serology can be predictive of HIV epidemic potential and hidden sexual risk behavior in the Middle East and north Africa. *Epidemics* 2010; **2**: 173–82.
- 189 Vaccarella S, Bruni L, Seoud M. Burden of human papillomavirus infections and related diseases in the extended Middle East and north Africa region. *Vaccine* 2013; **31** (suppl 6): G32–44.
- 190 Rowley J, Hoorn SV, Korenromp E, et al. Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ* 2019, published online July 4. ID:BLT.18.218651.
- 191 Sonnenberg P, Clifton S, Beddows S, et al. Prevalence, risk factors, and uptake of interventions for sexually transmitted infections in Britain: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *Lancet* 2013; **382**: 1795–806.
- 192 CDC, National Center for Health Statistics. National health and nutrition examination survey data, 1999–2014. <https://www.cdc.gov/nchs/nhanes/continuousnhanes/default.aspx> (accessed Sept 25, 2017).
- 193 Redmond SM, Alexander-Kisslig K, Woodhall SC, et al. Genital chlamydia prevalence in Europe and non-European high income countries: systematic review and meta-analysis. *PLoS One* 2015; **10**: e0115753.
- 194 Toomey KE, Rafferty MP, Stamm WE. Unrecognized high prevalence of *Chlamydia trachomatis* cervical infection in an isolated Alaskan Eskimo population. *JAMA* 1987; **258**: 53–56.
- 195 Walsh MS, Hope E, Isaia L, et al. Prevalence of *Chlamydia trachomatis* infection in Samoan women aged 18 to 29 and assessment of possible risk factors: a community-based study. *Trans R Soc Trop Med Hyg* 2015; **109**: 245–51.
- 196 Stamm WE. *Chlamydia trachomatis* infections of the adult. In: Holmes KK, Sparling FP, Stamm WE, et al, eds. Sexually transmitted diseases, 4th edn. New York: McGraw Hill Medical, 2008: 575–93.
- 197 Molano M, Meijer CJ, Weiderpass E, et al. The natural course of *Chlamydia trachomatis* infection in asymptomatic Colombian women: a 5-year follow-up study. *J Infect Dis* 2005; **191**: 907–16.
- 198 Heijne JC, Herzog SA, Althaus CL, Low N, Kretzschmar M. Case and partnership reproduction numbers for a curable sexually transmitted infection. *J Theor Biol* 2013; **331**: 38–47.
- 199 Omori R, Chemaitelly H, Althaus CL, Abu-Raddad LJ. Does infection with *Chlamydia trachomatis* induce long-lasting partial immunity? Insights from mathematical modelling. *Sex Transm Infect* 2019; **95**: 115–21.
- 200 Al-Mutairi N, Joshi A, Nour-Eldin O, Sharma AK, El-Adawy I, Rijhwani M. Clinical patterns of sexually transmitted diseases, associated sociodemographic characteristics, and sexual practices in the Farwaniya region of Kuwait. *Int J Dermatol* 2007; **46**: 594–99.
- 201 Ismail SO, Ahmed HJ, Grillner L, Hederstedt Issa BA, Bygdeman S. Sexually transmitted diseases in men in Mogadishu, Somalia. *Int J STD AIDS* 1990; **1**: 102–06.
- 202 Faisal A, Cleland J. Migrant men: a priority for HIV control in Pakistan? *Sex Transm Infect* 2006; **82**: 307–10.
- 203 Mumtaz GR, Riedner G, Abu-Raddad LJ. The emerging face of the HIV epidemic in the Middle East and north Africa. *Curr Opin HIV AIDS* 2014; **9**: 183–91.
- 204 Inhorn MC. Quest for conception: gender, infertility and Egyptian medical traditions. Philadelphia, PA: University of Pennsylvania Press, 1994.
- 205 Zurayk H, Sholkamy H, Younis N, Khattab H. Women's health problems in the Arab world: a holistic policy perspective. *Int J Gynaecol Obstet* 1997; **58**: 13–21.
- 206 Eggert-Kruse W, Buhlinger-Gopfarth N, Rohr G, et al. Antibodies to *Chlamydia trachomatis* in semen and relationship with parameters of male fertility. *Hum Reprod* 1996; **11**: 1408–17.

- 207 Eggert-Kruse W, Weltin M, Strowitzki T. Are chlamydial lipopolysaccharide-directed antibodies in seminal plasma or serum clinically significant during investigation of male infertility? *Urology* 2011; **77**: 1101–06.
- 208 Dietrich W, Rath M, Stanek G, Apfalter P, Huber JC, Tempfer C. Multiple site sampling does not increase the sensitivity of *Chlamydia trachomatis* detection in infertility patients. *Fertil Steril* 2010; **93**: 68–71.
- 209 Miron N, Socolov D, Mares M, et al. Bacteriological agents which play a role in the development of infertility. *Acta Microbiol Immunol Hung* 2013; **60**: 41–53.
- 210 Faour M. Fertility policy and family planning in the Arab countries. *Stud Fam Plann* 1989; **20**: 254–63.
- 211 Rashad H, Khadr Z. The demography of the Arab region: new challenges and opportunities. In: Human capital: population economics in the Middle East. Sirageldin I, ed. London: I B Tauris, 2002: 37–49.
- 212 Rashad H, Osman M. Nuptiality in Arab countries: changes and implications. In: Cairo papers in social science. Hopkins NS, ed. Cairo: American University in Cairo Press, 2003; 20–50.
- 213 Bristow CC, Mathelier P, Ocheretina O, et al. *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, and *Trichomonas vaginalis* screening and treatment of pregnant women in Port-au-Prince, Haiti. *Int J STD AIDS* 2017; **28**: 1130–34.
- 214 Cabeza J, Garcia PJ, Segura E, et al. Feasibility of *Chlamydia trachomatis* screening and treatment in pregnant women in Lima, Peru: a prospective study in two large urban hospitals. *Sex Transm Infect* 2015; **91**: 7–10.
- 215 Joseph Davey DL, Shull HI, Billings JD, Wang D, Adachi K, Klausner JD. Prevalence of curable sexually transmitted infections in pregnant women in low- and middle-income countries from 2010 to 2015: a systematic review. *Sex Transm Infect* 2016; **43**: 450–58.
- 216 Nguyen M, Le GM, Nguyen HTT, Nguyen HD, Klausner JD. Acceptability and feasibility of sexually transmissible infection screening among pregnant women in Hanoi, Vietnam. *Sex Health* 2019; **16**: 133–38.
- 217 Wynn A, Ramogola-Masire D, Gaolebale P, et al. Prevalence and treatment outcomes of routine *Chlamydia trachomatis*, *Neisseria gonorrhoeae* and *Trichomonas vaginalis* testing during antenatal care, Gaborone, Botswana. *Sex Transm Infect* 2018; **94**: 230–35.
- 218 Syred J, Engler B, Campbell L, Baraitser P, Sheringham J. Exploration of gender differences of *Chlamydia trachomatis* infection amongst young people reveals limitations of using sexual histories to assess risk in high-prevalence areas. *Int J STD AIDS* 2014; **25**: 564–70.
- 219 Horner PJ, Wills GS, Reynolds R, et al. Effect of time since exposure to *Chlamydia trachomatis* on chlamydia antibody detection in women: a cross-sectional study. *Sex Transm Infect* 2013; **89**: 398–403.
- 220 Hocking JS, Temple-Smith M, Guy R, et al. Population effectiveness of opportunistic chlamydia testing in primary care in Australia: a cluster-randomised controlled trial. *Lancet* 2018; **392**: 1413–22.
- 221 WHO. Strategies and laboratory methods for strengthening surveillance of sexually transmitted infections 2012. Geneva: World Health Organization, 2012. <http://www.who.int/reproductivehealth/publications/rtis/9789241504478/en/> (accessed Aug 9, 2014).
- 222 Bozicevic I, Riedner G, Calleja JM. HIV surveillance in MENA: recent developments and results. *Sex Transm Infect* 2013; **89** (suppl 3): iii11–16.
- 223 Ayoub HH, Awad SF, Abu-Raddad LJ. Use of routine HIV testing data for early detection of emerging HIV epidemics in high-risk subpopulations: a concept demonstration study. *Infect Dis Model* 2018; **3**: 373–84.
- 224 Reintjes R, Wiessing L. 2nd-generation HIV surveillance and injecting drug use: uncovering the epidemiological ice-berg. *Int J Public Health* 2007; **52**: 166–72.
- 225 Hessini L. Abortion and Islam: policies and practice in the Middle East and north Africa. *Reprod Health Matters* 2007; **15**: 75–84.
- 226 Shapiro GK. Abortion law in Muslim-majority countries: an overview of the Islamic discourse with policy implications. *Health Policy Plan* 2014; **29**: 483–94.