

STUDIES ON THE HOST-PARASITE RELATIONSHIP
OF STRONGYLOIDES RATTI IN RATS

A thesis submitted for the degree of Doctor of Philosophy
in the
University of London
(Faculty of Medicine)

by

Redwan Moqbel, B.Sc. (Baghdad), M.Sc. (London)

From the Department of Medical Helminthology,
London School of Hygiene and Tropical Medicine,
Keppel Street, London WC1E 7HT.

November, 1976

I dedicate this thesis
to my mother,
as a token of my
loving devotion

ABSTRACT

Studies were conducted on the host-parasite relationship in Strongyloides ratti infections in rats. During initial infection, rats acquired a strong immunity against this parasite and caused a drop in larval production and the expulsion of adult worms on day 25, ^{post-infection} PI. Infected rats were strongly resistant to reinfection and this immune response lasted for a period of three months, independent of the residual worm population. The immune response during initial infection caused a reduction in the size of adult worms after day 18 PI. Adult worms, which occupied the anterior half of the small intestine in the first 18 days, shifted to the posterior half before they were expelled. The immune response affected the adult worm in the intestine during a primary infection, the migratory larvae in the lung during a secondary infection and the skin stage following multiple infections. In heavily infected rats, larvae disseminated and were detected in the brain tissue.

A full infection with S. ratti induced complete resistance to reinfection, but adult worms were a major source of protective antigens, while migratory larvae did not stimulate an immune response against reinfection. "Normal" worms obtained on days 8 - 16 PI were able to continue their larval production when transferred to new, clean, rats; "damaged" worms obtained on days 18 - 25 PI were unable to do so. Worm damage during initial infection was irreversible. Short non-fertile worms found in rats early after challenge were able to resume their egg production and survive in new hosts. The possibility of adaptation was discussed.

22 day-old worms were examined under the electron microscope and were compared with 10 day-old worms. Damaged worms were smaller in size and showed vacuolation and tissue disorganisation, particularly the gut cells. Peculiar whorls were observed inside the gut lumen of damaged worms, and were suggested to be immune complexes. Precipitates were persistently observed surrounding the mouths of damaged worms. It was proposed that worm damage is instigated by immune complexes plugging the mouth and causing a state of

starvation through inhibition of food intake.

Worm expulsion was demonstrated to be an immunological event. Antibody mediated worm damage, in S. ratti, as an essential step in worm expulsion, was argued. Immune ^{mesenteric lymph node cells} (MLNC) alone, given on days 6 or 12 PI, were capable of inducing worm expulsion. Serum alone repeatedly induced auto-infection. A variety of factors were suggested to explain the action of transferred antibodies. Co-operation between immune MLNC and serum in inducing worm expulsion was only evident when MLNC were given on day 12 PI.

The tissue eosinophils increased in the mucosa of infected intestines immediately before worm expulsion, during initial and challenge infections.

The percentage of degranulating mesenteric mast cells increased during the period of expulsion, and the introduction of histamine into the gut of infected rats caused a significant acceleration of worm damage and expulsion. Aspirin, a potent inhibitor of prostaglandins, delayed the action of expulsion for 5 days suggesting a possible role of prostaglandins in this process.

Corticosteroid treatment caused the long persistence of S. ratti worms in the rats and indicated that worm damage was due to an immune response, and not to the worm's senescence. Levels of tissue eosinophils and degranulated mesenteric mast cells were also suppressed. No auto-infection was demonstrable when corticosteroid treatment commenced with the infection, but it was evident when the treatment started immediately prior to worm expulsion. Corticosteroids suppressed all manifestations of acquired immunity to reinfection. An attempt was made to render rats unresponsive to S. ratti infection by treating with corticosteroids at strategic periods during an initial infection.

ACKNOWLEDGEMENTS

My deep gratitude and appreciation to Professor George S. Nelson for his supervision, guidance and keen interest. My especial thanks to him for his assistance in securing a grant for me. It has been a great pleasure to work in his department.

My grateful appreciation to Dr. David A. Denham to whose tireless efforts, advice and help, throughout the course of this study, I am deeply indebted.

The encouragement, enthusiasm and assistance of Dr. Patrick B. McGreevy has helped immensely during my work.

Not being a pathologist, interpretation of my histological preparations needed comments from experts. For this I am very grateful to Professor George S. Nelson and Dr. Mansour F. Hussein, University of Khartoum.

I would like to record my thanks to Dr. David S. Ellis for the electron microscopy included in this work. Also to Dr. Diane J. McLaren of N.I.M.R. for her advice on fixing worm specimens for the E.M. work.

For her interest and helpful discussion, I would like to thank Dr. Bridget M. Ogilvie of N.I.M.R.

I would like to thank deeply Mr. Hugh Furse for his patience with my demands and his excellent technical help with the histological work. Also to Miss Sue Vickary.

Mr. A. E. C. Harvey of the Wellcome Laboratories, Beckenham, Kent, supplied the original strain of S. ratti.

Working with my colleagues Ted Bianco, Anne Moloney, Pakeer Othuman, Dr. Rosemary Murphy, Gilly Dean, Yohana Matola, Richard Suswillo and others, has been a very joyful, happy and beneficial experience. My thanks to them all for their assistance, companionship and good humour.

My thanks to the staff of the Animal House in the School, especially Marion Swann and Jim McQueen. Also to the staff of the Library.

Sincere thanks are due to my friends, Hazel and Golgaht Mossafai for their help with part of the photography, and Dr. Karim Nasher.

For her excellent work in the typing of this thesis, I am very grateful to Mrs. Elizabeth Manson.

My family, particularly my mother, whose loving encouragement during a period of severe difficulty, has been a constant source of inspiration to me.

To my wife Ann for her love, understanding and unfailing support, I wish to express my heartfelt thanks.

Finally, my sincere thanks to the Wellcome Trust for their generous grant, without which none of this would have been possible.

CONTENTS

	<u>Page</u>
Abstract	1
Acknowledgements	3
Contents	
Chapter 1 : Introduction	6
Chapter 2 : General Materials and Methods	20
Chapter 3 : The parasitological response of rats to primary and challenge infections with <u>Strongyloides ratti</u>	28
Chapter 4 : The effects of the immune response on adult <u>S. ratti</u>	69
Chapter 5 : The role of the different parasitic stages of <u>S. ratti</u> in stimulating the immune response	93
Chapter 6 : Studies on the mechanism of expulsion of <u>S. ratti</u> from the intestines of rats	107
Chapter 7 : Studies on the role of other factors in the process of worm expulsion in <u>S. ratti</u> infections	134
Chapter 8 : Studies on the effect of Betamethasone on the course of <u>S. ratti</u> infections in rats ..	156
Chapter 9 : Summary and Conclusions	187
List of References	191

CHAPTER 1

Introduction

Strongyloidiasis is a generalised term referring to infections by different species of Strongyloides which parasitize the intestinal mucosa of vertebrate hosts, including man. Infections with these soil-transmitted nematodes are cosmopolitan in distribution. A few species, especially those which infect sheep and swine, have veterinary importance as they produce loss of weight, diarrhoea and enteritis. About 3% of the world population is infected with Strongyloides stercoralis. This infection, which is frequently innocuous, is usually acquired by people, such as miners and farmers, whose professions involve contact with soil.

Interest in human strongyloidiasis has increased recently following numerous reports that a mild infection with this parasite can turn into a serious, and often fatal, infection under certain conditions. It has been found that patients asymptotically harbouring this parasite, who are treated with corticosteroids or X-ray therapy, as immunosuppressive agents, developed a sudden and overwhelming infection. In many cases dissemination of larvae occurs and they invade many internal organs causing severe damage and often death (Rogers and Nelson, 1966; Cruz et al., 1966; Willis and Nwokolo, 1966; Civantos and Robinson, 1969; Yim et al., 1970; Fagundes et al., 1971; Bitoun et al., 1972; Cookson et al., 1972; Neeffe et al., 1973; Purtilo et al., 1974).

As experimental studies on S. stercoralis were not possible, the present research work adopted the rodent species Strongyloides ratti as an alternative to investigate the following points:

1. Aspects of acquired resistance which develop in rats following initial, second and multiple infections with S. ratti.
2. The immunogenic stage(s) of infection.
3. Effects of the immune response on adult S. ratti.
4. Mechanisms involved in the immune expulsion of adult S. ratti from the rats.

- 7
5. Alterations in the equilibrium between host and parasite following immunosuppression by corticosteroids.

Taxonomical Review

The members of the suborder Rhabdiata of the order Rhabdiasidea (Yamaguti, 1961) are of particular interest since they manifest a gradual evolutionary trend from free-living organisms to facultative parasites. The main genera of this suborder are placed in two major families: Rhabdiasidae and Strongyloididae. Worms belonging to these two families form a link between free-living and parasitic forms by manifesting the phenomenon of alternations of generations. The free-living adult generation consists of male and female worms, while only parthenogenetic females are found in the parasitic generation.

Family Rhabdiasidae

Nematodes belonging to this family, some of which are considered below, are parasites of reptiles, amphibians and mammals, but they also exhibit free-living activities. Rhabditis strongyloides live in soil, organic matter or water, but their larval forms are also found on the hairs of the perianal region of dogs where they cause itching pustules (Chandler and Read, 1961). Rhabditis pellic was recovered from the human vagina and urine (Chandler and Read, 1961). Another species Diploscapter coronata was found in the aspirated stomach contents of patients suffering from a total lack of gastric hydrochloric acid (Chandler, 1938), and it is possible that this worm is only able to parasitize such patients.

Family Strongyloididae

This family was established by Chitwood and McIntosh (1934)

with only two genera, Strongyloides and Parastrongyloides, as representatives. Chitwood and Chitwood (1950) defined the essential diagnostic criteria for this family, the summary of which is:

"Parasitic generation parthenogenic or dioecious, filariform, stoma cup-shaped or greatly reduced. Females with 2 divergent uteri, ovary reflexed, vulva near middle of body. Live in gastro-intestinal tract of vertebrates".

Species of Strongyloides

Strongyloides spp. were discovered in animals early in the second half of the 19th century, in both sheep and swine. The differentiation of species presented a formidable task since morphologically similar worms were reported from a wide range of vertebrate hosts (Grassi, 1878; Looss, 1911; Bayliss, 1923).

S. stercoralis-like worms were recovered from frogs (Alfieri, 1908) and this led to a temporary belief that the human infection was a zoonosis, with frogs acting as reservoir hosts. Attempts to classify the different species of Strongyloides were made, utilizing various systems and criteria (Chandler, 1925; Sandground, 1925; Henry, 1931; Noto-Soediro, 1933; Desportes, 1945; Basir, 1950). Little (1966a, b) used the following criteria for the specific identification of the various species of Strongyloides:

- (1) The shape of the stoma in the parasitic females viewed en face;
- (2) The shape of the ovary in the parasitic female;
- (3) The developmental stage, or stages, passed in host's faeces.

The first of these criteria was found to be the most consistent and accurate.

Over 40 species of Strongyloides are known to exist, and table

- (1) lists some of the better known species found in vertebrates.

Table 1 : A list of some of the parasitic species
of the genus Strongyloides

<u>PARASITE</u>	<u>HOST</u>
<u>S. stercoralis</u> (Bavay, 1876) Stiles and Hassall, 1903	Man
<u>S. fülleborni</u> Von Linstow, 1905	Old-world primates, not infrequently, man
<u>S. papillosus</u> (Wedle, 1856) Ransom, 1911	Sheep and rabbits
<u>S. oebus</u> Darling, 1911	Rhesus monkeys
<u>S. ransomi</u> Schwarz and Alicata, 1930	Swine
<u>S. myopotami</u> Artigas and Pacheco, 1937	Nutria
<u>S. westeri</u> Ihle, 1917	Horses
<u>S. agouti</u> Griffith, 1940	Agouti and guinea-pigs
<u>S. ratti</u> Sandground, 1925	Rats
<u>S. venezuelensis</u> Brumpt, 1934	Rats
<u>S. lutrae</u> Little, 1966	Otters
<u>S. dasypodis</u> Little, 1966	Armadillos
<u>S. ardea</u> Little, 1966	Herons
<u>S. serpentis</u> Little, 1966	Snakes
<u>S. procynosis</u> Little, 1966	Raccoons
<u>S. physali</u> Little, 1966	Toads
<u>S. elephantis</u> Greve, 1969	Elephants
<u>S. avium</u> Cram, 1929	Fowls and Turkeys

Strongyloides stercoralis:

This species was discovered by Normand, a French Army surgeon, in 1876, from the stools of French soldiers who died from diarrhoea in Cochin-China. At autopsy, minute adult worms were found attached to the mucosa of the ileum, and these were morphologically different from those found in the faeces. The worm of the faeces was named Anguillula stercoralis and the intestinal worm A. intestinalis by Bavay in 1876. Further studies by Grassi (1879), Perroncito (1880) and Leukart (1882) revealed that these two worms were separate stages of the life cycle of one species. The name Strongyloides stercoralis (Bavay, 1876) Stiles and Hassall, 1903 was later adopted. Initial studies on the mode of infection with this parasite were carried out by Zimm (1899), Askanazy (1900), Van Durme (1902) and Ransom (1907). Looss (1905) concluded that the migration of the infective stage of this species was similar to that of Ancylostoma duodenale. Fülleborn (1914), working on experimental S. stercoralis infections in dogs, contributed greatly to our present knowledge of the route of infection and the invasion of the skin by infective filariform larvae. He also proved that their migration, within the host en route to the intestine, was obligatory. These findings were confirmed by Sandground (1926, 1928). Further elucidation of the life cycle in experimental strongyloidiasis was presented by Faust (1933) who described the successive developmental stages of the parasitic phase. The filariform infective larvae penetrated the skin and migrated, via the blood, to the lungs where they developed into postfilariform larvae, which migrated through the trachea and oesophagus to the small intestine of the host and were then called preadolescent females. Adult females were found buried in the mucosa and these produced embryonated eggs which usually hatched into rhabditiform larvae in the lumen of the gut, from where they were carried out with the faeces.

The parasitic male enigma:


Several authors have recognized that parasitic adult females of Strongyloides spp. were parthenogenetic (Rovelli, 1888; Brumpt, 1911; Ransom, 1911; Chitwood and Graham, 1940). However, parasitic males have been found in man and experimentally infected animals (Kreis, 1932; Sotolongo Guerra et al., 1971; Sotolongo Guerra, 1972). Unlike female worms, parasitic males were considered to be temporary residents which were expelled following fertilization of the females (Faust, 1933). On cytological evidence, Sandground (1926) claimed that the worms were hermaphroditic. Graham (1935, 1936, 1938a, 1938b) established infections with S. ratti from a single larva which indicated that parasitic adult females did not require conventional fertilization by "parasitic" males to produce new progeny. No reduction division was noticed by Chang and Graham (1957) during oogenesis in the free living female of S. papillosus, and they proposed that this species was pseudogamous. In pseudogamy the sperm does not penetrate the egg but rather activates it to divide. Bolla and Roberts (1968) investigated gametogenesis in the free living form and the chromosomal complement in both forms of S. ratti. They refuted Sandground's claim that the worms were hermaphroditic and instead confirmed the phenomenon of pseudogamy in the free living forms and parthenogenesis in the parasitic females. However, they did not offer any explanation for the occasional finding of parasitic males in human strongyloidiasis.

Direct and indirect modes of development:

Most Strongyloides spp. follow one of two methods of development outside their hosts, namely direct (homogonic) or indirect (heterogonic) development (Figure 1). This phenomenon suggests that this group of nematodes has only recently evolved to be parasites of vertebrates.

Attempts were made to explain the mechanisms which control these two modes of development. Internal and external environmental conditions, age of the worms and separate biological strains were

Figure 1 : Heterogamy and homogamy in Strongyloides sp.,
 modified from Sandground, 1926



suggested (Leichtenstern, 1899; Darling, 1911a, b; Sandground, 1926). Beach (1935, 1936) stressed that favourable nutritional conditions outside the host induced the rhabditiform larvae to develop into free-living, heterogonic, dioecious adult worms; while unfavourable surroundings induced the direct homogonic path.

Graham (1935, 1936, 1938a, b, 1939a, b) selected pure homogonic and heterogonic lines by breeding from single filariform larvae, but certain homogonic lines produced heterogony and vice versa (Graham, 1940), which indicated that this phenomenon was not, totally, genetically controlled.

As yet, the exact factors, which determine whether or not a strain of Strongyloides spp. develop homogonically or heterogonically, are not fully elucidated. However, it seems that a combination of genetic and environmental factors are responsible for the switch over from one mode of development to another (Smyth, 1962).

Auto-infection in Strongyloidiasis

Most species of Strongyloides possess the rare ability to multiply within the host. Under certain circumstances the rhabditiform larvae moult twice in the lumen of the host, become infective larvae and penetrate the host. This can occur in two ways (Faust and de Groot, 1940):

1. Internal auto-infection: Infective larvae invade the intestinal mucosa and travel, via the portal circulation, to the lungs, migrate to the gut and develop into adult females.
2. External auto-infection: Some rhabditiform larvae moult to become infective larvae near the perianal region and these invade the epithelial lining of the anal sphincter and/or the skin of the perineum, repeating the same migratory cycle as above.

This kind of development might explain how S. stercoralis survives in humans for up to 36 years (Brown and Perna, 1958). It is suggested that auto-infection occurs in people with a depressed cell-mediated immune response (Purtilo et al., 1974). This aspect and the

clinical and pathological picture of human strongyloidiasis will be further discussed in Chapter 8 .

Human strongyloidiasis as a zoonosis:

Natural infections with Strongyloides spp. occur in dogs (Brumpt, 1922; Chandler, 1925) and although the human species S. stercoralis was successfully established in dogs (Tashiro, 1912; Chira, 1918; Sandground, 1928; Faust, 1933), there is insufficient evidence for the existence of a zoonosis between dog and man. However, human infections with the simian species, S. fülleborni occur in Africa (Pampiglione and Ricciardi, 1971a, 1972a), and experimental infections with this species were successfully established in a human volunteer (Pampiglione and Ricciardi, 1971b, 1972a, b).

A few communities on the Fly River in New Guinea harbour an unidentified species of Strongyloides for which no animal host has been found. It has been suggested that this species is probably S. fülleborni, which is interesting since there is no simian fauna on the Island (Knight, 1976, personal communication).

Strongyloides ratti Sandground, 1925:

Grassi and Segre (1887) found a distinct species of Strongyloides in wild mice, Mus sylvaticus and M. decumanus. Hall (1916) included the strain which he recovered from rats with S. papillosus of sheep and rabbits, while Sandground (1925) proposed that this was a new species and named it S. ratti.

The adult females of S. ratti (Plate 1) range between 1.85 - 3.03 mm (average: 2.35 mm). The cuticle has fine transverse striation. The buccal cavity is shallow and surrounded by six minute papillae. The oesophagus is 22-33% of the total body length. The anal orifice is situated 32 - 50 μ from the posterior end of the body, and the vulva 1.27 - 1.73 mm from the anterior end of the body. The ovaries are directly recurrent, bending close to the oesophagus and the posterior end of the intestine. As many as 11 eggs are present in



Plate 1 : Adult female of S. ratti (x 80)



Plate 2 : Egg of S. ratti, phase contrast (x 160)



Plate 3 : Rhabditiform, first stage (L1) larva (x 250)



Plate 4 : Filariform, infective, third stage (L3) larva (x 200)




Plate 5 : Trifurcated tail of third stage larva (x 250)



Plate 1 : Adult female of S. ratti (x 80)



Plate 2 : Egg of S. ratti, phase contrast (x 160)



Plate 3 : Rhabditiform, first stage (L1) larva (x 250)




Plate 4 : Filariform, infective, third stage (L3) larva (x 200)



Plate 5 : Trifurcated tail of third stage larva (x 250)



Plate 1 : Adult female of S. ratti (x 80)



Plate 2 : Egg of S. ratti, phase contrast (x 160)



Plate 3 : Rhabditiform, first stage (L1) larva (x 250)



Plate 4 : Filariform, infective, third stage (L3) larva (x 200)



Plate 5 : Trifurcated tail of third stage larva (x 250)

both uteri. The tail is finely tapered. The eggs (Plate 2) (measuring 47-52 x 28-31 μ) are usually embryonated when leaving the female, and are normally laid in a strand deposited deep in the mucosa, where they hatch. First stage (L1) rhabditiform larvae (Plate 3) average 420 μ in length with a maximum width of 17-21 μ . They are similar to first stage larvae of hookworms except for their short buccal cavity. The third stage (L3) infective filariform larvae (Plate 4) are extremely active. They average 650 μ in length and show two distinctive, equal, parts of the gut; the oesophagus and the intestine. The tip of the tail is characteristically trifurcated (Plate 5).

Life cycle of *S. ratti* in the rat (Figure 2):

Infective larvae enter the body through the hair follicles. Invasion of the cutaneous tissue is rapid and larvae reach the venules and lymphatics within 3 hours, and large numbers of larvae are found in the blood 10 hours after exposure. The majority (Spindler, 1958) arrive in the lungs about 16 hours post-infection (PI) and grow into post-filariform larvae, which migrate to the intestine about 23 hours PI (Abadia, 1963; Wertheim and Lengy, 1965). No moulting occurs during this phase except in rare cases where larvae are trapped in the lungs (Wertheim and Lengy, 1965). Upon reaching the intestine, 34 hours PI, larvae burrow into the mucosa and moult twice at 47 and 74 hours into adult females. Female worms reach their full size 120 hours PI (Wertheim and Lengy, 1965) and egg formation commences at 81 hours. Eggs and larvae can be detected in the intestinal wall and faeces 85-92 hours PI.

Sheldon (1937c) compared subcutaneous, oral, intramuscular, intracolonic, intraperitoneal, intragastric and intravenous routes of *S. ratti* infection. The oral, where larvae were allowed to penetrate the pregastric mucosa, and the subcutaneous routes gave higher yields of adult worms.




Figure 2. Life cycle of *Strongyloides ratti* Sandground, 1925

S. ratti and S. venezuelensis:

Concurrent infections with these two nematodes occur in wild rats (Little, 1961; Wertheim and Lengy, 1964; Araujo, 1967), but they have different distributions along the small intestine (Araujo, 1967). Adult S. ratti mainly occupied the anterior 20 cm of the small intestine but stretched in their distribution as far as 60 cm, distally. They were found buried in the crypts of Lieberkühn where they usually deposited their eggs. S. venezuelensis, on the other hand, occupied the anterior 15 cm of the small intestine, with the majority in the proximal 5 cm. The females of this species were found near the surface of the villi where they laid their eggs. S. ratti reached the intestine about 20 hours before S. venezuelensis (Wertheim, 1970a, b). Wertheim (1970b) concluded that each species, when occurring concurrently, behaved as if present alone and demonstrated its own characteristic distribution in the intestine.

Host specificity of S. ratti

S. ratti was found in 45-70% of sewer rats (Sandground, 1925; Tanabe and Takaishi, 1936; Nashimura, 1943). Amano et al. (1958) and Abe et al. (1960) showed that Rattus norvegicus had a higher prevalence rate of S. ratti than R. rattus.

Attempts to infect mice with S. ratti have yielded a variable degree of success. Brackett and Bliznick (1949) passaged S. ratti through mice 19 times without increasing its infectivity to this host. It appears that different strains of mice react differently to such infections. CFl mice were more susceptible to S. ratti (Goldgraber and Lewert, 1965) than other strains (Sandground, 1925; Sheldon, 1937d; Brackett and Bliznick, 1949).

There are no reports of human infections with this parasite, but there is some evidence that repeated exposure to its infective larvae produces an immediate type hypersensitivity reaction at the site of exposure.

Host sex differences in *S. ratti* infections:

This aspect of the host - parasite relationship in this system is still controversial. Roman (1951) noticed a slightly higher incidence of *S. ratti* in female *R. norvegicus*, *Apodemus sylvaticus* and *Clethrionomys glareolus* but concluded that the sex of the host had no effect on the recovery rate. Schacher and Chee-Hock (1960) and Wertheim and Lengy (1964) obtained equal distribution of *S. ratti* in male and female hosts. Abe et al. (1960) and Wertheim (1963) reported that male rats had a higher rate of infection than females. Katz (1961, 1962, 1963, 1967) investigated this problem and the effects of sex hormones on *S. ratti* infections. He concluded that female rats had a greater worm burden during the first three weeks of infection, while male rats showed a higher burden at the end of 4 weeks, but no definite role of sex hormones was demonstrated.

This introduction has reviewed various features of the different species of *Strongyloides* generally and *S. ratti* particularly. Other more specific aspects of the host - parasite relationship, in this parasite and other related nematodes, are reviewed in their relevant chapters.

CHAPTER 2General Materials and MethodsThe Parasite:

The original culture of S. ratti was obtained from the Wellcome Laboratories, Beckenham, Kent. This strain did not develop free-living adult forms in culture.

Laboratory Hosts:

Random bred, female albino Wistar rats (Tuck & CO.), at 100 - 150 gms (8 weeks old) were used to maintain the parasite, and for most of the experimental work. An inbred stock of the same strain was used for experiments involving passive and adoptive transfer of immunity.

Maintenance of the Parasite:

10 - 12 gms of freshly passed infected faecal pellets were collected once a week by giving a gentle squeeze to the perianal area which stimulated the passing of faeces. Fresh pellets were also collected from the cages. The faecal pellets were weighed and washed 3 - 4 times with distilled water. The faeces were left in a beaker with a few drops of distilled water for 30 - 60 minutes and were thoroughly mixed.

Test-tube/filter paper method for culturing infected faeces:

This method was introduced originally by Harada and Mori (1951) for the diagnosis of hookworms. It was improved by Sasa



Plate 6 : The test tube/filter paper (Harada-Mori) culturing method.



Plate 6 : The test tube/filter paper (Harada-Mori) culturing method.




Plate 6 : The test tube/filter paper (Harada-Mori) culturing method.

et al. (1957) and was recommended by the W.H.O. expert committee for the diagnosis of soil transmitted helminthiasis (W.H.O., 1963). 18 cm x 18 mm test tubes were placed in special racks and 7 ml of distilled water added. 0.5 gm of the moistened faeces was smeared onto 15 cm x 15 mm filter paper strips using a wooden tongue depressor (Lewis). The smeared strip was inserted into the test-tube leaving 7 cm clear at one end with the clean end touching the bottom of the distilled water in the tube (Plate 6). The tubes were sealed with aluminium foil and kept in an incubator at 28°C for 3 - 4 days. During this period the first stage rhabditiform larvae moulted twice and migrated towards the water in the bottom of the tube. The filter papers were removed and the infective larvae collected and concentrated by centrifugation at 1000 rpm for 5 minutes. The larval suspension was stirred with a teflon-coated magnet on a magnetic stirrer (Voss). Aliquots were taken with a Pasteur pipette into a McMaster type counting chamber (Figure 3). If the larval suspension was too concentrated, dilutions of 1:10 or 1:20 were made. A total of 5 sequential counts of live larvae were made in the chamber under a dissecting microscope at x 20 magnification and the number of larvae calculated from the mean of the 5 counts and the volume of the suspension.

Methods of infection:

A. Subcutaneous inoculation: Each infection was of 1000 larvae per rat, in 0.2 ml of distilled water, injected subcutaneously in the inguinal area.

B. Percutaneous infections: The rats were anaesthetized with a mixture of air and Penthrane (Abbott Laboratories). The fur was clipped from the abdomen, or the back, and the animals were placed on a flat surface. The skin was moistened with distilled water and a stainless steel ring (1.5 cm in diameter and 1 cm in depth) placed in position. 1000 infective larvae in 0.5 ml of water were pipetted into each ring (Plate 7). Infective larvae invaded the skin within 2 - 3 minutes, but the rings were left for 10




Figure 3 : McMaster's type counting chamber

minutes. Larvae were rarely recovered from the distilled water inside the rings following this period of exposure. This method was introduced by Smithers and Terry (1965) for exposing rats and other rodents to cercariae of Schistosoma mansoni.

Recovery of adult worms:

The rats were killed by injecting Nembutal (Abbott Laboratories) intraperitoneally. The small intestine was removed, immediately, and slit using a blunt-nosed pair of scissors. The intestine was shaken in a jar of water at room temperature to remove the digesta. This resulted in a loss of 0.5% of the adult worms, but the absence of ingesta made subsequent examinations much easier and more accurate. The intestine was transferred to a jar containing 25 - 30 ml of 0.85% saline and incubated at 37°C for 2 - 3 hours. Subsequently, the intestine was removed from the jar to a ruled petri dish and the adult worms counted under a dissecting microscope (x 15). The lining of the gut was scraped with the edge of a clean microscope slide, the mucosa squashed between two glass plates and the number of undetached worms counted. 2 - 3% of the worms did not detach.

A modified Baermann funnel was used for the mass collection of adult worms. The intestine was slit, washed and placed on a 50 mesh sieve (pore size 300 microns, Endecotts Test Sieves Ltd.) in a funnel filled with saline and left for 3 hours at 37°C. The worms migrated through the sieve to the bottom of the funnel. They were collected and washed twice with fresh saline.

Measuring of worms:

Adult worms were fixed in 10% formal saline and kept in Bijoux bottles. A sample of 30 - 80 worms was pipetted onto a slide and covered with a coverslip. The fresh preparation was examined immediately under x 300 magnification on a Nikon microscope with a built-in Camera Lucida. This magnification was ideal for drawing the lengths of worms on white paper. The overall lengths of drawn



Plate 7 : Percutaneous infection with S. ratti



Plate 8 : Preparation of mesentery for mast cell staining



Plate 7 : Percutaneous infection with S. ratti



Plate 8 : Preparation of mesentery for mast cell staining



Plate 7 : Percutaneous infection with S. ratti




Plate 8 : Preparation of mesentery for mast cell staining

worms were determined with a roller-head map measurer and later converted into millimeters.

Histology:

Relevant tissues were collected and fixed immediately in fixatives depending on the type of staining required. The most frequently used fixative was 10% formol-saline. Bouin's and Carnoy's solutions were used to fix tissues prior to staining with specific dyes.

For the fixation of intestinal tissue, the small intestine was removed and 1 cm segments, each inclusive of a Peyer's patch, were taken. Each segment was slit and placed between two small sheets of glass which were kept together with a rubber band, to avoid curling of the tissue when placed in the fixative. The segments were sectioned longitudinally, passing through the middle of the Peyer's patches. The tissues were dehydrated, cleared and embedded in paraffin wax in a Histokinette (British American Optical Co.). The blocked tissues were sectioned at 5 μ m on a rotary microtome (Jung AG Co., Heidelberg) and stained in Haematoxylin and Eosin (H. & E.), which was used routinely with most of the tissues. Carbol Chromotrope (Lendrum, 1944) was used for staining tissue eosinophils. Sections were stained with Astra-blue (Raymond Lomb) adjusted to pH 0.25 with concentrated HCl (Bloom and Kelly, 1960) and counter stained with Safranin O for intestinal mast cells (Enerback, 1966). Pericapillary mesenteric mast cells were stained for 1 min. in 0.25% acidic Toluidine blue (Smith and Atkinson, 1956) after fixing the stretched mesenteries on a slide (Plate 8) with 10% formol saline for 5 - 10 mins.

Methyl green - pylonin Y was used to stain Peyer's patches in the intestine.

Statistical analysis of data:

The calculation of the mean and standard error was performed

on an Olivetti Programma 101. The results of the majority of experiments are expressed as mean (standard error). Student's 't' test was used to assess the significance of some of the results. Significantly different results had to show a probability P value of less than 0.10.

Other specific materials and methods, used during the course of this study, are presented separately in their relevant chapters.

CHAPTER 3

The parasitological response of rats to primary and challenge infections with Strongyloides ratti

A worm population is expelled much earlier in rodents which have been given a primary ^{nematode} infection and are then challenged with more larvae. This has been shown to occur in Nippostrongylus brasiliensis in rats (Ogilvie, 1969), Trichinella spiralis infections in mice, rats and guinea-pigs (McCoy, 1931, 1940; Roth, 1939; Culberston, 1942; Larsh et al., 1952; Zaiman et al., 1953; Denham, 1968), Trichuris muris infection in mice (Wakelin, 1967) and Trichostrongylus colubriformis in guinea-pigs (Herlich et al., 1956; Herlich, 1963; Rotwell et al., 1973).

The immune response in different hosts, against parasitic nematodes, causes a variety of changes, in addition to the early loss of worms (Ogilvie and Jones, 1973). In some filarial infections, the adult worms show a decline in the number of microfilariae produced in immune hosts (Bertram, 1966; Denham et al., 1972). Microfilariae of Litomosoides carinii remain arrested in the pleural cavity of immune cotton rats and are almost unable to enter the blood circulation (Bagai and Subrahmanyam, 1970). Adult worms are also morphologically affected by the immune response of the host, and show a delay in their migration, moulting and growth and this produces irrevocably stunted worms (Soulsby, 1962; Bertram, 1966; Sinclair, 1970; Jarrett et al., 1971; Denham et al., 1972; Ogilvie and Jones, 1973).

Early work by Sheldon (1937a, b) on experimental S. ratti infections revealed that rats became resistant to reinfection following an initial infection. The development of this resistance in a primary infection caused a sharp drop in the larval output 19 days post-infection and infection was terminated after day 26 (Sheldon, 1937a, c) although a few larvae were still produced. Sheldon (1937a) conferred a marked degree of protection against S. ratti by serial

injections of heat-killed infective larvae, but Griffin (1972) was unable to repeat this result. This sensitization was specific as no resistance against S. ratti was stimulated by heat-killed infective larvae of S. stercoralis (Sheldon, 1939). Thomas (1970) showed that different strains of rats developed similar marked resistance to S. ratti following initial infection and that low-level "trickle" infections caused a delay in the expulsion of the worms for up to 8-10 weeks.

The movement of worms in the gut during the course of infection has been investigated in different rodent nematode systems, e.g. N. brasiliensis and T. spiralis (Brambell, 1965; Larsh et al., 1952). There is no published information on such movement in S. ratti, but the majority of adults occupy the anterior 15-20 cm of the small intestine and, in decreasing numbers, the remaining 45-60 cm, during the first few days after their arrival in the intestine (Abadie, 1963; Wertheim, 1970).

In this chapter an investigation is made of the effects of the rat's response on initial and subsequent infections with S. ratti.

Experiment 3.1:

This experiment was designed to study the development of resistance to primary and secondary infections. Fifty five female Wistar rats were separated into two groups: Group A, 30 rats, were infected subcutaneously with 1000 (\pm 73) L3 each on day 0. Group B, 25 rats, were left uninfected.

Preliminary cultures were made from a random sample of five rats from Group A on days 6, 10, 18, 20 and 26 PI, to monitor larval output during the course of primary infection. On day 30 PI, both groups were treated with thiabendazole (TBZ) at 100 mg/kg for two consecutive days. TBZ is highly effective against human and experimental strongyloidiasis (Franz, 1963; Jorge and Barbosa, 1964; Chaia and Cunha, 1966; Bezjack, 1969). On day 32 PI five rats from group A were autopsied to ascertain the efficacy of TBZ. No adult worms were recovered from any of these rats. On day 34 PI, the remaining rats of groups A and B were challenged with 1000 (\pm 51)

infective larvae of S. ratti per rat. Faecal cultures from both groups were made every two days, commencing on day 6 post-challenge. 5 rats from each group were autopsied on days 10, 14, 18, 21 and 26 after challenge to count the number of adult worms. The results are presented in Tables 2 and 3, and Figures 4 and 5.

During initial infection, larval output sharply increased from 3415 to 7770 L3/gm of faeces between days 6 and 8 PI. There was then a plateau for approximately 8 days followed by a sharp drop in the number of larvae/gm of faeces reaching to 108 L3/gm on day 26 PI. Following a challenge infection, larval output did not exceed 3 L3/gm of faeces.

The highest number of adult worms was recovered on day 14 PI and decreased from then on until it reached a mean of 32 worms per rat on day 26 PI. The number of adult worms recovered following a challenge infection was low with a maximum mean of 20 worms on day 10 PI. The worms were rapidly expelled around day 10 post-challenge and an average number of approximately 2.7 worms was maintained between days 18-26 post-challenge.

The worms recovered from the challenged group on day 10 post-challenge were 54.7% shorter than those recovered from the challenge control on the same day (Figure 6). Following a challenge infection, female worms were unable to produce the expected number of eggs. Adult worms recovered from the challenge control group on day 26 PI, were shorter and less active than those recovered at an earlier stage of the same infection.

Experiment 3.2:

This experiment was a repeat of experiment 3.1, with special attention being paid to the changes in the size of worms during primary and secondary infections, and the timing of the occurrence of these changes. Seventy seven female Wistar rats were divided into two major groups, A and B:

The 31 rats of group A were infected with 1000 (\pm 88) larvae each on day 0, while group B was left uninfected. On day 28 PI, both groups were treated with 2 x 100 mg/kg of TBZ. Three rats from

Table 2, Exp. 3.1:

Mean number of larvae produced during initial and challenge infections with S. ratti

Day post-challenge	Mean (SE) number of larvae/gm of faeces	
	Challenge infection	Initial infection
	Group A	Group B
6	3	3415 (+ 138)
8	2.3	7770 (+ 481)
10	1.83	5733 (+ 129)
12	1	3909 (+ 215)
14	1.5	2866 (+ 129)
16	0.6	2619 (+ 225)
18	0.5	1211 (+ 71.3)
20	1.7	299 (+ 23)
22	0.3	533 (+ 30.8)
24	0.5	202 (+ 23.8)
26	0.5	108 (+ 9.8)

Table 3, Exp. 3.1:

Mean number of adult worms recovered during initial and challenge infections with S. ratti

Days post-challenge	Challenge infection		Initial infection	
	Number of rats	Mean (SE) number of worms	Number of rats	Mean (SE) number of worms
10	5	20 (<u>±</u> 6.4)	5	452 (<u>±</u> 24)
14	5	4.4 (<u>±</u> 2)	5	614 (<u>±</u> 45)
18	5	2.6 (<u>±</u> 1)	5	217 (<u>±</u> 36)
21	5	2.8 (<u>±</u> 1.3)	5	102 (<u>±</u> 10.5)
26	5	2.8 (<u>±</u> 1.1)	5	32 (<u>±</u> 16.5)

Table 3, Exp. 3.1:

Mean number of adult worms recovered during initial and challenge infections with S. ratti

Days post- challenge	Challenge infection		Initial infection	
	Number of rats	Mean (SE) number of worms	Number of rats	Mean (SE) number of worms
10	5	20 (+ 6.4)	5	452 (+ 24)
14	5	4.4 (+ 2)	5	614 (+ 45)
18	5	2.6 (+ 1)	5	217 (+ 36)
21	5	2.8 (+ 1.3)	5	102 (+ 10.5)
26	5	2.8 (+ 1.1)	5	32 (+ 16.5)

Figure 4, Exp. 3.1:

Larval production during the course of initial and challenge infection with *S. ratti*.

- Group A, Initially infected and challenged
- Group B, challenge control.




Figure 5, Exp. 3.1:

Mean number of adult worms recovered from two groups of rats.
●—● initially infected and challenged
○—○ challenge control





Figure 6, Exp. 3.1:

Mean length of worms recovered on day 10 post challenge



group A killed two days later had no worms in their intestines. Both groups were infected with 1000 (\pm 67.5) L3s per rat on day 34 PI (day 0 post-challenge). The lungs of the rats killed on days 1 and 3 were chopped in saline with a pair of fine scissors, and incubated at 37°C for about 3 hours, and the number of larvae counted. Many live and active larvae were recovered from the lungs in group B while only two dead larvae were recovered from the challenged group. The results are presented in Tables 4 and 5 and Figures 7 and 8.

The speed of migration of larvae from the skin to the gut was not the same in the two groups. The few larvae which succeeded in avoiding the immune reaction in the lungs reached the intestine on day 5 post-challenge. The highest number of worms (87.8) was recovered from the challenged group on day 7 post-challenge but the number decreased sharply as expulsion was faster in this group. Only 2 worms were recovered on day 11 post-challenge, and the rats were worm free one day later. After initial infection, the peak number of adult worms was 463 on day 14 PI and this number decreased to 13.8 on day 26 PI.

The fecundity of the worms was calculated by dividing the mean number of larvae produced by each rat, by the mean number of adult worms obtained from rats of the same group killed on the same day. The fecundity of the worms (Figure 9) in the control group reached a peak of 22.8 larvae/worm on day 7 PI and declined after day 18 to a minimum of 1.5 larvae/worm on day 26 PI. The fecundity of the worms in the challenged group was extremely low being 0.091 and 0.215 L3/worm on days 7 and 9 post-challenge, respectively. The adult worms attained their maximum length (2.81 mm) on day 9 PI and there was little or no change until day 14 PI (Figure 10). A significant decrease in the length was seen on day 18 PI and this coincided with the onset of expulsion. The worms reached their shortest length (0.89 mm) on day 26 PI. There was a slight increase in the length of the worms recovered after a challenge infection from 1.28 to 1.43 mm, between days 5 and 7 post-challenge. However, their length decreased soon after and reached a minimum of 0.59 mm on day 11 post-challenge.

The distribution of lengths of adult worms recovered during the

Table 4 Exp: 3.2 Larval production, number of adult worms calculated and number of larvae per worm (= fecundity) during primary and secondary infections

Days	Primary infection			Secondary infection		
	mean (SE) larvae / gm faeces	mean (SE) number of adult worms	larvae per worm	mean (SE) larvae / gm faeces	mean (SE) number of adult worms	larvae per worm
7	5261 (+ 446)	159 (+ 15.5)	22.8	9.09	87.8 (+ 22.1)	0.09
9	4497 (+ 565)	380 (+ 23.8)	7.1	9.16	30.6 (+ 7.21)	0.21
14	2290 (+ 77.4)	463 (+ 14.6)	11.9	-	-	-
18	520 (+ 44.2)	355 (+ 18)	2.17	-	-	-
21	254 (+ 16)	168 (+ 13.7)	2.34	-	-	-
26	23.6 (+ 0.55)	13.8(+ 8.74)	1.50	-	-	-

Table 5 Exp: 3.2 Worm recovery at autopsy during initial and challenge infections and mean (SE) length of recovered worms

Days	Primary infection				Secondary infection			
	number rats/ group	mean (SE) number of adult worms	number measured	mean (SE) length in mm	number rats/ group	mean (SE) number of adult worms	number measured	mean (SE) length in mm
1	3	-	-	-	3	-	-	-
3	3	-	-	-	3	-	-	-
5	5	80 (+ 8.58)	45	2.36 (+ .02)	4	30.2 (+ 8.8)	40	1.28 (+ .03)
7	5	159 (+ 15.5)	34	2.64 (+ .04)	5	87.8 (+ 22.1)	40	1.43 (+ .04)
9	5	380 (+ 23.8)	42	2.81 (+ .05)	5	30.6 (+ 7.21)	31	1.23 (+ .03)
11	5	419 (+ 16.4)	40	2.80 (+ .05)	5	2.0 (+ 0.31)	10	0.59 (+ .05)
12	N.D.	-	-	-	3	0	-	-
14	5	463 (+ 14.6)	47	2.3 (+ .03)	N.D.	N.D.	N.D.	N.D.
18	5	355 (+ 18)	35	1.8 (+ .05)	"	"	"	"
21	5	168 (+ 13.7)	43	1.23 (+ .03)	"	"	"	"
26	5	13.8(+ 8.74)	41	0.89 (+ .03)	"	"	"	"

Figure 7, Exp. 3.2:

Larval production in two groups of rats.
●---● previously infected and challenged
○---○ challenge control.




Figure 8, Exp. 3.2:

Adult worms recovered from two groups of rats.
●—● initially infected and challenged
○—○ challenge control.




Figure 9, Exp. 3.2:

Fecundity (number of larvae per worm) in two groups of rats.
●—● initially infected and challenged
○—○ challenge control.




Figure 10, Exp. 3.2:

Mean length of adult worms recovered from two groups of rats.
□ initially infected and challenged
■ challenge control





Figure 11, Exp. 3.2:

Length distribution of adult worms during the course of initial and challenge infections.



course of initial and second infections (Figure 11) indicates that the worms shrank rather than that there was a selective loss of a population of long adult worms first.

Experiment 3.3:

This experiment was carried out to study the distribution of adult S. ratti in the intestines of infected rats and their migration during initial and challenge infections.

Thirty seven rats were infected with 1000 (\pm 65) L3s each. Groups of four rats were killed on days 9, 13, 18, 20, 22 and 25 PI. The small intestine of each rat was divided into 4 equal segments and the number of adult worms in each part determined. The results are summarized in Table 6 and Figure 12. During the first 18 days, 79% of the worms occupied the anterior segment of the small intestine. By day 13 PI, the worms had begun to move posteriorly. An average of 65% of the worms were recovered from the last segment of the intestine between days 20 and 25 PI compared to 7.3% in the first segment, during the same period. The pattern of worm distribution in the two middle segments, during the course of infection, was similar to the trend in the first and last segments. This migration to the distal part of the intestine coincided with the reduction in the number of adult worms and their shrinkage. No worms were recovered from the lumen of the segments of the gut suggesting that worms migrated through the tissue of the small intestine rather than in the lumen.

The remaining 12 rats were treated with 2 x 100 mg/kg of TBZ and reinfected with 1000 (\pm 81.6) L3 each. The number of worms after challenge was low, and there was an equal distribution in the anterior and posterior segments on day 6. 60% of the worms were found in the last quarter of the intestine on day 8, prior to their expulsion on day 10, while the anterior segment harboured only 16% of the total adult population.

This experiment has shown that adult S. ratti undergo a posterior movement towards the last quarter of the small intestine, prior to their expulsion following initial and challenge infections.


Table 6 Exp. 3.3:

The distribution of adult S. ratti in the small intestine of rats,
during the course of primary and secondary infections

Days PI	Mean (SE) number of adult worms				
	Total	Segment 1	Segment 2	Segment 3	Segment 4
9	400 (+13.5)	333 (+25)	53 (+14.5)	10 (+1.9)	2 (+0.85)
13	647 (+58)	543 (+48)	74 (+10)	20 (+4)	10 (+1)
18	617 (+30)	440 (+25)	89 (+8.5)	64 (+3.4)	24 (+3.4)
20	256 (+66)	37 (+29)	10 (+3)	54 (+16)	153 (+31)
22	194 (+37)	5 (+2.3)	16 (+2.2)	39 (+10)	134 (+33)
25	83 (+10)	60 (+8)	17 (+2)	5 (+1)	1 (+0.5)
Post challenge					
6	36 (+11)	14 (+4)	3 (+1.4)	4.5 (+2.5)	14.5 (+7.6)
8	24 (+8)	4 (+2.8)	1 (+0.5)	5 (+0.8)	14 (+5)
10	0	0	0	0	0

Figure 12, Exp. 3.3:

The distribution of adult *S. ratti* in the intestine of rats during primary and secondary infections.



Experiment 3.4:

The purpose of this experiment was to study the duration of the immune response to S. ratti in rats. Rats initially infected developed very strong resistance to reinfection. However, a few adult worms persisted for up to 100 days (Sheldon, 1937a).

Fifteen rats were divided into 3 groups of 5 rats each. Ten of these rats were infected with 1000 (\pm 93.7) L3/rat while the other 5 were left uninfected to act as challenge controls. After 30 days, 5 infected rats were treated with 2 x 100 mg/kg of TBZ to remove the residual population, while the other infected group of 5 rats was left untreated. Three months later, all the rats were challenged with 1000 (\pm 99.4) L3s/rat. Ten days later, all the rats were autopsied and the number of adult worms in the anterior and posterior halves of the small intestine was counted. Faecal cultures were made on days 6, 8 and 10 post-challenge. The results are presented in Tables 7 and 8 and Figures 13 and 14. There was no significant difference between larval production in the two previously infected groups. The mean number of larvae/gm of faeces was much lower in the two experimental groups than the control group on days 6, 8 and 10. Only 3.5% and 5% of the worms developed in the two previously infected groups, while the challenge control group showed a 53% recovery on day 10 post-challenge and PI, respectively.

Judging from the level of larval production and adult recovery, it was clear that a substantial degree of resistance was maintained in rats previously infected with S. ratti for more than 3 months. The experiment also showed that the residual worm population was not responsible for maintaining this immunity as its removal with TBZ did not eliminate the immune response. The distribution of the worms recovered in the previously infected groups was similar to that observed in a normal challenge infection prior to worm expulsion.

Experiment 3.5:

This experiment was designed to investigate the fate of infective larvae and adult worms following a challenge infection. It also

Table 7 Exp: 3.4 Larval production following challenge after an interval of 3 months, in two groups

Days	Mean (SE) number of larvae / gm of faeces		
	Treated with thiabendazole	Not treated with thiabendazole	Challenge control
6	393 (+ 28.6)	339 (+ 33.3)	2857 (+ 182)
8	55.5 (+ 19.8)	138 (+ 18.8)	7933 (+ 416)
10	17.7 (+ 8.2)	19.4 (+ 8)	6560 (+ 165)

Table 8 Exp: 3.4 Mean number of adult worms recovered on day 10 after challenge in groups reinfected after an interval of 3 months

<u>Reinfected after 3 months</u>	Mean (SE) number of adult worms		
	anterior half	posterior half	total
treated with thiabendazole	14.6 (+ 3.7)	21.2 (+ 1.4)	35.8 (+ 4.4)
not treated with thiabendazole	17 (+ 1.87)	34.2 (+ 3.98)	50.2 (+ 4.88)
challenge control	450 (+ 46)	82.8 (+ 7.3)	533 (+ 47.8)

Figure 13, Exp. 3.4:

- Larval production in three groups of rats.
○—○ reinfected after 3 months interval, treated with TBZ
△—△ reinfected after 3 months interval, not treated with TBZ
●—● challenge control





Figure 14, Exp. 3.4:

Mean number of adult worms recovered from the anterior A, and posterior, P, intestine of three groups, on day 10 post challenge.

■ challenged after 3 months interval, treated with TBZ
▨ challenged after 3 months interval, not treated with TBZ
□ challenge control



studied the effect of the immune response on the larvae during their skin-lung-intestine migration. Fifteen rats were used in this experiment and were divided into the following four groups: Group 1: four rats each received an initial infection followed by 5 challenges of the same inoculum (1000 L3/rat), at intervals of 20, 15, 10, 5 and 5 days, respectively. The first 5 infections were given, percutaneously, on different areas of the back, while the challenge was given percutaneously on a shaven area of the abdomen.

Group 2: four rats, each received an initial infection of 1000 L3s and a challenge of 1000 L3s, percutaneously, on the day the last challenge was given to group 1.

Group 3: four rats, received a percutaneous infection on the same day as the last challenge dose was given to group 1.

Group 4: three rats were left uninfected to act as normal controls.

The following histopathological and parasitological changes were observed:

Skin

Initial percutaneous infections with infective larvae of S. ratti were very fast. The filariform larvae penetrated the skin within 30 minutes. Very few changes were observed in the skin soon after exposure, but there was a slight dilatation of the walls of both capillaries and lymphatics, with no direct evidence that this was due to infection (Plate 9). No eosinophils were seen during the passage of an initial infection through the skin.

Following a second infection, live larvae were demonstrated in the hypodermis with little or no tissue response around them during the first 3 hours post-exposure (Plate 10). After this period, the reaction in the skin changed slightly, with infiltration of lymphocytes, mainly in the deeper layers of the subcutaneous tissue. The reaction was patchy, confined to limited areas of the skin. No neutrophils or eosinophils were detected (Plate 11). Abundant macrophages, and lymphocytes were seen between 4-6 hours post-exposure, with an obvious absence of polymorphs. This indicated that the reaction had developed into a more pronounced humoral type response.




Plate 9 : Skin; 30 minutes after primary exposure (H. & E., x 80)




Plate 10 : Skin; 30 minutes after second exposure (H. & E., x 80)




Plate 9 : Skin; 30 minutes after primary exposure (H. & E., x 80)




Plate 10 : Skin; 30 minutes after second exposure (H. & E., x 80)

After 24 hours, the reaction was dermal with signs of progressive reaction near hair follicles. It appeared that the larvae were capable of evading the skin stage of the immune reaction early after challenge. However, after 24 hours, there was a response to the passage of the infective larvae which manifested itself in a cellular infiltration of lymphocytes and macrophages and total absence of eosinophils (Plate 12).

The histopathological picture changed entirely following 6 infections. The response to the invading population of filariform larvae was immediate and severe (Plate 13). Within the first 30 minutes, the lymphatics were surrounded by neutrophils, eosinophils and mononuclear macrophages. Dead larvae were seen in the subcutaneous tissue, surrounded by many eosinophils, lymphocytes and macrophages (Plate 14). After 24 hours, pronounced oedema and vascular and lymphatic dilatation was evident, as well as a severe infiltration of eosinophils, lymphocytes and macrophages (Plate 15). It was very difficult to quantify the tissue eosinophil response in secondary and multiple infections. However, based on the number of eosinophils per 10 random high power fields, Table 9 was produced to present the difference in this response in the four groups.

Lungs:

Live larvae were frequently seen in histological sections of the lungs of rats infected once (Plate 17) but with little or no reaction surrounding them (Plate 18). A few active larvae were also recovered from chopped lungs after incubation at 37°C in saline. Few lymphocytes, polymorphs and macrophages were seen. Few eosinophils were found in initially infected and normal unexposed rats.

Only dead larvae were recovered from minced lungs during the first day in a secondary infection. Histologically, the lungs developed areas of lymphoid infiltration composed of lymphoblast centres surrounded by small lymphocytes, suggestive of lymphoid hyperplasia (Plate 19). A dead larva was observed in one of these lymphoid centres with a very strong lymphocytic reaction round it (Plate 20). Many other such granulomas were seen developing around fragments of worms. Vast numbers of eosinophils had infiltrated the lung tissue




Plate 11 : Skin; 30 minutes after second exposure, no eosinophil reaction round live larva (Lendrum's, x 250)




Plate 12 : Skin; 24 hours after second exposure, no eosinophil reaction (Lendrum's, x 250)




Plate 11 : Skin; 30 minutes after second exposure, no eosinophil reaction round live larva (Lendrum's, x 250)




Plate 12 : Skin; 24 hours after second exposure, no eosinophil reaction (Lendrum's, x 250)



Plate 13 : Skin, generalised reaction in the hypodermis, 30 minutes after 6th exposure (H. & E., x 80)




Plate 14 : Skin, eosinophil infiltration around dead larva, following 6 exposures (Lendrum's, x 400)




Plate 13 : Skin, generalised reaction in the hypodermis, 30 minutes after 6th exposure (H. & E., x 80)




Plate 14 : Skin, eosinophil infiltration around dead larva, following 6 exposures (Lendrum's, x 400)




Plate 15 : Skin, extensive eosinophil, lymphocyte and macrophage infiltration 24 hours after 6th exposure (Lendrum's, x 400)




Plate 16 : Normal unexposed rat skin (Lendrum's, x 80)




Plate 15 : Skin, extensive eosinophil, lymphocyte and macrophage infiltration 24 hours after 6th exposure (Lendrum's, x 400)



Plate 16 : Normal unexposed rat skin (Lendrum's, x 90)




Plate 17 : Lung following initial infection, live larva (L) within alveoli (H. & E., x 80)




Plate 18 : Lung, live migrating larva in alveoli with very little reaction (H. & E., x 400)




Plate 17 : Lung following initial infection, live larva (L) within alveoli (H. & E., x 80)




Plate 18 : Lung, live migrating larva in alveoli with very little reaction (H. & E., x 400)




Plate 19 : Lung, lymphoid hyperplasia following second infection (H. & E., x 80)




Plate 20 : Lung, dead larva surrounded by small lymphocytes following second infection (H. & E., x 400)




Plate 19 : Lung, lymphoid hyperplasia following second infection (H. & E., x 80)




Plate 20 : Lung, dead larva surrounded by small lymphocytes following second infection (H. & E., x 400)

Table 9, Exp. 3.5:

Tissue eosinophil response in the skin of rats; following 6 infections (group 1);
2 infections (group 2); 1 infection (group 3); and non-infected control (group 4)

Time after last exposure	Group 1	Group 2	Group 3	Group 4
½ hour	++	-	-	-
1 hour	+	-	-	-
1½ hours	+	-	-	-
3 hours	++	-	-	-
4 hours	+++	-	-	-
6 hours	++	+	-	-
24 hours	++++	+	-	-

(Plate 21) and cuffed the blood vessels and the bronchial passages. The multiple granulomas and the infiltration of many granulocytes within the lung tissue was suggestive of pneumonitis.

Following multiple infections no larvae reached the lungs and larvae were killed in the skin. There was a generalized reaction which was manifested by a mild increase in the number of eosinophils and polymorphonuclear cells in the lung tissue.

Small intestine:

There were very few pathological changes in the intestinal tissue following the arrival of worms from the lungs, and during the first two weeks of initial infection. The adult worms embedded their anterior ends in the mucosal crypts and were sometimes seen in the mucosa. The eggs which the adult worms passed were laid in strings of 8-14 eggs inside the submucosa; this was repeatedly seen in squash preparations of the intestinal tissue. Later during an initial infection, infiltration of lymphocytes, plasma cells and eosinophils started, particularly around days 18 PI (Plate 23), when changes in the length and structure of the worms began. Although the Peyer's patches enlarged gradually during the course of infection, few adults or embryonated eggs were seen within the mucosa surrounding the Peyer's patches (Plate 24). This may be due to the gradual movement of the worms from one site to another. There was an increase in the number of eosinophils around the time of worm expulsion (see Chapter 7).

In second infections the few adult worms, which avoided the immune response in the lungs, arrived in the intestine and attached themselves to the mucosa. Egg production was very low, and no egg strings were seen. There was an increase in the number of eosinophils, plasma cells and macrophages in the submucosa of the villi. The Peyer's patches did not increase beyond the size which they reached late during initial infection.

No adult worms were seen in the small intestine following multiple infections.




Plate 21 : Lung, extensive infiltration of eosinophils following a second infection (Iendrum's, x 400)




Plate 22 : Normal unexposed lung (H. & E., x 80)




Plate 21 : Lung, extensive infiltration of eosinophils following a second infection (Lendrum's, x 400)




Plate 22 : Normal unexposed lung (H.& E., x 80)




Plate 23 : Small intestine, eosinophil response around embryonated eggs (E) (Lendrum's, x 250)




Plate 24 : Small intestine, larvae lying between villi and a Peyer's patch (H. & E., x 250)




Plate 23 : Small intestine, eosinophil response around embryonated eggs (E) (Lendrum's, x 250)




Plate 24 : Small intestine, larvae lying between villi and a Peyer's patch (H. & E., x 250)

Experiment 3.6:

In this experiment four different infective doses were given to groups of rats and particular attention was given to the distribution of the developing stages in the host's organs and the percentage of development in the gut.

Twenty two rats were divided into four groups: Group 1, four rats, received 1000 L3 each; Group 2, six rats, received 4000 L3 per rat; Group 3, six rats, received 6000 L3; Group 4, six rats, received 8000 L3. All the surviving rats were killed on day 10 PI, and faecal cultures were prepared from all groups on days 6, 8 and 10 PI.

The results of this study are summarized in Table 10. The group which received 1000 L3s harboured a mean of 457 worms on day 10 PI; a percentage development of 45.7%. The rats which received 4000 L3/rat harboured a mean of 1159 worms which represented 28.9% of the original inoculum.

A similar, proportionately low, recovery rate was observed in the rats which received 6000 L3s each. Four members of this group died on day 3 PI and the two which survived were autopsied on day 10 PI. The four rats which died were emaciated, weak and moribund before their death. They all lost their sight and behaved abnormally.

Live larvae were found in abundance in the lungs and in lesser numbers in the brain and intestines. Larvae were also seen in histological sections of the brains (Plates 25 and 26).

Most of the worms recovered from the intestines of the 3 rats which died on day 3 PI were preadults. A mean of 604 preadults was recovered from the gut which represented 10% of the initial inoculum, while the other two rats, which were autopsied on day 10 PI, harboured a mean of 920 adult worms which represented 15.3% of the original inoculum.

All the rats which received 8000 L3 died within two days following a short period of crisis. There were numerous internal haemorrhages, particularly in the lungs, where many live larvae were seen, but no larvae were recovered from the intestines.

This experiment has shown that in heavily infected rats, larvae disseminated throughout the tissues and the rate of worm

Table 10 Exp: 3.6 Adult worm counts in four groups of rats following initial infection
with 4 different infective doses of S. ratti

	Group 1 Infected with 4000 L3 / rat	Group 2 Infected with 6000 L3 /rat		Group 3 Infected with 8000 L3 / rat	Group 4 Infected with 1000 L3 / rat control
		Died day 3 PI	Killed day 10 PI		
Individual	973	412 (I4)	971	All rats died on	431
counts	1007	743 (I4)	869	day 2 PI	512
	1132	657 (I4)		No larvae were	439
	1731	N.D.		recovered	447
	893				
	1220				
Mean (SE)	1159 (+123)	604 (+99.1)	920 (+51)		457 (+18.5)




Plate 25 : Cerebrum, live larva within brain tissue with marked degree of tissue damage (H. & E., x 250)




Plate 26 : Cerebellum, live larva within brain tissue with no reaction (H. & E., x 80)




Plate 25 : Cerebrum, live larva within brain tissue with marked degree of tissue damage (H. & E., x 250)




Plate 26 : Cerebellum, live larva within brain tissue with no reaction (H. & E., x 80)

development in the gut was low.

Discussion:

The results of these experiments clearly demonstrate that rats produce strong resistance to S. ratti and that this becomes even stronger after multiple infections. Following a primary infection, very few larvae from a challenge infection manage to reach the intestine, and the parasitological and histological study suggests that they are destroyed in the lungs. Their death leads to extensive lymphoid hyperplasia and formation of granulomas in the lung. Migrating larvae, presumably, share antigenic characteristics with adult worms which induce circulating memory cells to recognize them at the lung stage and initiate the drastic changes which lead to the death of the majority of the larvae, and only a few survive and develop into stunted worms in the small intestine.

After a series of infections, the rat responds much more rapidly to challenge and larvae are killed in the skin and do not reach the lungs. Soulsby (1962) suggested that enzymes used by infective larvae to penetrate the skin, act as functional antigens and that the immune response causes an inhibition of these enzymes and this leads to the immobilisation of these larvae. Such inactivation of invading larvae in human strongyloidiasis can result in creeping eruption. Lee and Lewert (1957) described a factor in the serum of immune rats which inhibited the action of skin penetration by S. ratti and Schistosoma mansoni. However, the level of this factor did not increase in hyper-infected rats.

The development of resistance during a primary infection with S. ratti is slower in comparison with that produced by N. brasiliensis and T. spiralis infections, but whether this is due to S. ratti being less immunogenic or not is still undetermined. The effects of the host's response were seen from day 14 PI. The reduction in fecundity, migration away from the initial site of parasitization and expulsion of the adult worms were the most striking features of this response.

Adult N. brasiliensis migrate anteriorly within the intestine

during the course of infection (Chandler, 1935; Brambell, 1965; Alphey, 1970; Connan, 1974). This movement has been attributed to metabolic and immunologic factors, but there is mounting evidence that the evacuation of the worms from a site of preference during early infection, to another part of the intestine, is mainly due to the immune response of the host, particularly the humoral arm (Alphey, 1970; Connan, 1974). Most adult T. spiralis were found in the anterior half of the small intestine in old mice (Larsh and Hendricks, 1949). Migration of these worms down the small intestine was also observed prior to their expulsion (Larsh et al., 1952). Similar worm movement was noted by Campbell (1967) and Duckett (1970) but was attributed to feeding factors and to host strain. Podhajecky (1962) and Denham (1968) could find no evidence for such a movement.

The capacity of previously infected rats to resist challenge in S. ratti infections was maintained for up to three months after the initial worm expulsion, with residual worms playing no role in maintaining this immunity. Similar duration of immune memory was observed in mice infected with T. muris (Wakelin, 1967). Rats initially infected with N. brasiliensis showed a very strong resistance to reinfection after 2 months. Rats challenged 7 months after an initial infection, expelled their worm burden after a short lag and independent of any residual population in the gut (Ogilvie, 1965).

Infective larvae disseminated in the different parts of the host, when introduced in very high numbers. The infective larvae in the brain may have been a reason for the moribund state of these rats and may have contributed towards their ultimate deaths. The number of developing adult worms in the gut, following exposure to a very high infective dose, was low, the majority being found in the lungs and brain.

Massive strongyloidiasis in man occurs occasionally (Brown and Perna, 1958; Yoeli et al., 1963; Cahill, 1967). This usually leads to death due to the penetration, by the infective larvae of S. stercoralis, of different body organs (Walker-Smith et al., 1969; Reiff and DeMoraes, 1972; Purtillo et al., 1974; Ali-Khan and Seemayer, 1975). The involvement of the nervous tissue, particularly the brain, in massive disseminated strongyloidiasis was confirmed

histologically (Neefe et al., 1973), and was attributed to auto-infection with S. stercoralis.

CHAPTER 4The effects of the immune response on
adult *S. ratti*

Chandler (1938) and Ogilvie (1969) have shown that adult *N. brasiliensis*, when transferred to clean rats before exposure to the developing immune response, i.e. at days 6-9, were able to continue egg production at the same level as in a primary infection. Worms which were fully exposed to the immune response, i.e. at day 10-13, did not resume egg production in new, clean, hosts and were rapidly expelled. However, a phenomenon which Ogilvie (1972) called "adaptation" was observed in worms which had developed in immunised rats. Adult worms, when harvested 6-9 days after a challenge infection and transferred to clean recipients, persisted much longer than did normal worms (Chandler, 1936; Ogilvie and Hockley, 1968; Ogilvie, 1972). Although these adapted worms remained stunted, they resumed egg production and their cytological features were restored to normal (Ogilvie and Hockley, 1968). Ogilvie (1972) suggested that adaptation to an immune environment occurred in the lungs. The adaptation of worms is phenotypic and the progeny of adapted worms were similar to the progeny of normal worms (Ogilvie, 1972).

No parallel to this phenomenon of worm adaptation has been recorded outside the rat/*N. brasiliensis* system, although 10 day-old adult *T. spiralis* resumed larval production when transplanted into clean mice (Mougeot and Lancaster, 1973). When 18 day-old worms were transplanted, larval production was not resumed suggesting that they were permanently damaged. A considerable degree of stunting and diminished fecundity of *T. spiralis* females occurred prior to expulsion from the intestine (Denham and Martinez, 1970) and during reinfection (Samrad and Coombs, 1951; Rappaport and Wells, 1951).

Nematodes are exposed to a variety of factors produced by the host, but their cuticle is a very tough barrier (Lee, 1965) and is

impermeable to most substances, including antibodies (Marks *et al.*, 1968; Lee, 1969a). Presumably, the various mediators of the immune response reach the parasite through its orifices.

It is now well established that the humoral components of the immune response act on the worms prior to their expulsion, and cause structural damage particularly to the intestinal cells of the worms (Ogilvie and Jones, 1971, 1973; Love *et al.*, 1976). When these changes occur during the late part of an initial infection with N. brasiliensis, the damage is irreversible (Ogilvie and Hockley, 1968).

The gut of normal N. brasiliensis consists of large cells (Lee, 1969a), the apical surface of which is lined with well-developed microvilli and contain active mitochondria, a large number of ribosomes and Golgi complexes. The gut cells of worms exposed to the immune response are markedly different. The ribosomes are replaced by large vacuoles filled with carbohydrates and lipids (Ogilvie and Hockley, 1968). Similar changes in the structure of gut cells occur in adult Haemonchus placei (Harness *et al.*, 1973).

Within the lumen of the gut of N. brasiliensis, Taliaferro (1940) and Taliaferro and Sarles (1939) repeatedly noted a mass of material which they suggested was an immune precipitate, but this has not been confirmed (Ogilvie and Hockley, 1968; Lee, 1969a). Ogilvie (1964) claimed that these precipitates have no role in the process of worm damage, since similar material was formed round worms incubated in antiserum from rats vaccinated with worm extracts, yet these rats were completely susceptible to a challenge infection.

Externally, precipitates were also seen on worms, in immune hosts, particularly near the mouth (Sarles, 1938; Taliaferro and Sarles, 1939; Oliver-Gonzalez, 1940).

The reproductive system is also adversely affected by the immune response, particularly in females, which results in retardation of their egg production (Lee, 1969b). Female Ostertagia ostertagi develop structural abnormalities of the vulval flap, following multiple infections, due to the host's developing immune response (Michel *et al.*, 1972).

This chapter is concerned with the effects of immunity on S. ratti during primary and secondary infections and the amount of change which is observed in these worms, during initial infection.

Materials and Methods

Transplantation of adult worms:

Adult worms were harvested from infected donor rats using a Baermann funnel filled with saline at 37°C. The worms were counted and the required number collected in 0.1 ml volumes of saline in 1 ml syringes. Recipient rats were anaesthetised with a mixture of air and Penthrane, and the fur of the ventral side was clipped and the duodenum exposed by laparotomy. The worms were injected directly into the duodenum (Plate 27). The abdominal wall was sutured with patent suture silk No. 00 (Archibald and Turner Co.). The skin was joined together with 12 mm Michel clips. The rats recovered from anaesthesia within two minutes of the withdrawal of Penthrane. This procedure was carried out under clean but not aseptic conditions.

Preparation of adult *S. ratti* for examination under light and electron microscope:

Adult worms were collected in the usual way and washed twice with sterile saline. For light microscopy, worms were fixed in Bouin's fluid and left for ½-1 hour and then washed twice for 1 hour in absolute alcohol and cleared twice in Toluene. They were then embedded in paraffin wax in a small glass tube, the worms settled in the bottom and were easily accessible for sectioning when the tube was broken. Sections were cut at 5 µm and stained in haematoxylin and eosin.

For electron microscopy, worms were cut into two using a pair of fine watch-makers forceps, in 3% glutaraldehyde in 0.1 M cacodylate buffer at pH 7.4 and left for two hours, in the presence of calcium ions. The rear ends were discarded, the suspension was centrifuged gently and the worm material embedded in agar. The agar pellet was cut up and the pieces containing the worms were returned to glutaraldehyde for another 19 hours. They were washed 4 times within 21 hours at equal intervals with cacodylate buffer to which was added 0.02 M sucrose and postfixed in 1% osmium



Plate 27 : Intra-duodenal transplantation of adult S. ratti
into an anaesthetized rat



Plate 27 : Intra-duodenal transplantation of adult S. ratti
into an anaesthetized rat

tetroxide in Millonig's buffer (Millonig, 1961) at pH 7.4 for 2 hours at 4°C. After this they were washed 3 times in distilled water (15 minutes each) and fixed in 0.5% solution of uranyl acetate containing 45 mg/ml of sucrose (de Harven, 1967) at pH 5.0 for 1 hour at 4°C.

Cross sections were cut on a Huxley (Cambridge) ultramicrotome, and were collected on Formvar coated copper grids stained with lead citrate (Venable and Coggeshall, 1965), before examination in a Zeiss Em 9A electron microscope.

Experiment 4.1:

This experiment was designed to study the fate of worms of different ages transplanted into new recipients. A total of 118 female Wistar rats (120 gm in weight) were used for this experiment. Three sets of donor and recipient rats were prepared. The timing of the infections was such that all donor rats were killed on the same day.

Donor group 1: Forty rats were infected with 3000 L3/rat, reinfected 30 days later with 3000 L3 each, killed 7 days later and their adult worms collected in a Baermann funnel. Sixteen groups of 300, 7 day-old, worms from "immune" rats in 0.1 ml of saline were inoculated, intraduodenally, into 16 rats of the recipient group A.

Donor group 2: Ten rats were infected with 3000 L3/rat and killed on day 10 PI. Sixteen groups of 300, 10 day-old, worms were injected intraduodenally into 16 recipient rats of group B.

Donor group 3: Thirty rats were infected with 3000 L3/rat and killed on day 22 PI. Sixteen groups of 300, 22 day-old, worms were injected, intraduodenally, into 16 recipient rats of group C.

Ten worms from each donor group were measured before transplantation. Worms from donor group 1 had a mean length of 1.45 (\pm 0.05) mm, those of group 2 had a mean length of 2.87 (\pm 0.03) mm and those of group 3 a mean length of 1.01 (\pm 0.06) mm, respectively.

Four rats from each recipient group were killed on days 5, 7, 12 and 15 post-transplantation and faecal cultures were made on the same days. The worms were counted and measured. The results are

presented in Table 11 and Figures 15 and 16.

The worms transferred from resistant hosts into normal recipients (group A) were able to re-establish themselves and commence egg production. However, the larval production was not very high and was maintained for only 10 days. An average of 22.8 worms was recovered on days 5 and 7 post-transplantation and decreased to an average of 11.3 worms on days 12 and 15 post-transplantation. The average length of these worms was 1.33 mm. The 10 day-old worms transplanted from an initial infection into recipient rats of group B continued their egg production on a normal level with the output on day 5 post-transplantation being 3080 larvae/gm of faeces. Larval production dropped on day 12 post-transplantation and ceased on day 15 post-transplantation. The average length of these worms followed the pattern of a normal infection and dropped from 2.3 mm on day 5 post-transplantation to 0.91 mm on day 15 post-transplantation. The 22 day-old worms in recipient rats of group C did not recover after transplantation, and the majority of the worms were expelled by day 5 post-transplantation. No larvae were recovered from the faeces of this group.

This experiment demonstrated that worms, recovered early during challenge infection, can resume producing larvae without any increase in their overall length. This might indicate that damage of worms during challenge infection is reversible. Also the experiment showed that worms recovered late during initial infection could not establish themselves in new hosts and hence, the damage was irreversible. Thus worms obtained on days 8 - 16 PI were considered to be "normal" with 10 day-old worms being typical. Worms recovered between days 18 - 25 PI were considered "damaged", with 22 day-old worms being typical. Worms collected early during a challenge infection were damaged as shown by their reduced size, but were able to recover.

Table 11 Exp: 4.1

Mean larval production, number and lengths of adult worms during four different periods

A: received 300 worms from a challenge infection

B: received 300 worms from day 10 PI

C: received 300 worms from day 22 PI

Day post transpl- antation	Mean (SE) number of larvae / gm faeces			Mean (SE) number of adult worms			Mean (SE) length of adult worms (mm)		
	A	B	C	A	B	C	A	B	C
5	1866 (+ 237)	3074 (+ 257)	0	25.2 (+ 3.56)	182 (+ 10.3)	1.25 (+ .62)	1.4 (+ .04)	2.3 (+ .09)	.88(+ .04)
7	960 (+ 88)	650 (+ 118)	0	20.2 (+ 3.94)	151 (+ 26)	0	1.38 (+ .03)	1.4 (+ .05)	-
12	54.9(+ 12)	26.6 (+ 5.5)	0	10.2 (+ 1.37)	92.7 (+ 8.35)	0	1.30 (+ .05)	1.11 (+ .04)	-
15	0	0	0	12.5 (+ 2.53)	2.5 (+ 1.65)	0	1.27 (+ .05)	0.91 (+ .06)	-

Figure 15, Exp. 4.1:

- Mean faecal larval production in three groups of rats.
- A. received 300 worms from challenge infection.
 - B. received 300, 10 day-old worms from initial infection
 - C. received 300, 22 day-old worms from initial infection.





Figure 16, Exp. 4.1:

Mean number of adult *S. ratti* recovered from three groups. ■ Group A received 300 worms from challenge infection; □ Group B received 300, 10 day-old worms from initial infection; ■ Group C received 300, 22 day-old worms from initial infection.



Experiment 4.2:

Studies were made on the morphological changes in S. ratti during the course of an initial infection. Worms were taken from a group of 5 rats on days 10 and 22 PI. The two types of worms were divided into two lots for fixation for light and electron microscopy.

The "normal" worms had a mean length of $2.83 \pm .03$ mm with a range of 2.55 - 3.21 mm, while the damaged worms had a mean length of $1.1 \pm .06$ mm with a range of 0.55 - 1.9 mm.

Light microscopy:

The difference in size between "normal" and "damaged" worms under the light microscope was very obvious. The majority of the sections of normal adult worms (Plate 28) had a consistent elliptical shape with most of the internal organs staining densely with H. & E., while the duplicated reproductive tract was usually darkly stained with haematoxylin. The gut showed no vacuolation. The damaged worms (Plate 29) were irregular in shape with all organs staining faintly with H. & E. The reproductive tract was smaller and stained lightly with haematoxylin. The gut showed patches of vacuolation surrounding the lumen which was not patent in most sections. The area under the cuticle and the circular muscles was always vacuolated and unstained.

Electron microscopy:

No studies on the ultrastructure of S. ratti have been reported in the literature. All the observations reported below on the two types of worms, are compared with N. brasiliensis. The observations on the structural and morphological changes in these worms were concentrated on the anterior third and changes in the gut and surrounding tissues were considered. The most obvious difference at EM level, as by light microscopy, between the 10 and 22 day-old




Plate 28 : C.S. Normal adult
S.ratti (H. & E., x 400)



Plate 29 : C.S. Damaged adult
S.ratti (H. & E., x 400)



Plate 30 : Low power electron micrograph of section of a 10 day-old
S. ratti (x 2000)



Plate 31 : Low power electron micrograph of section of a 22 day-old
S. ratti (x 2000)



Plate 28 : C.S. Normal adult
S.ratti (H. & E., x 400)



Plate 29 : C.S. Damaged adult
S.ratti (H. & E., x 400)



Plate 30 : Low power electron micrograph of section of a 10 day-old
S. ratti (x 2000)




Plate 31 : Low power electron micrograph of section of a 22 day-old
S. ratti (x 2000)

worms, was that of the overall size (Plates 30 and 31). There were no apparent differences in the cuticle or the underlying musculature, which was typical of nematodes (Lee, 1969). However, beneath the muscle layer, the damaged worms often showed vacuolated spaces or disorganised tissues which were not found in normal worms (Plates 32 and 33). There were other differences between the two kinds of worms such as; a tendency for the damaged worms to contain bigger interstitial spaces (Plates 32 and 33), a reduced amount of rough endoplasmic reticulum in cells around the gut and an increased number of lysosomes (Plates 34 and 35), many of which would appear to be forming secondary lysosomes with cell lipids.

It was very difficult to quantify these findings, which were made on sections cut from over 50 worms. However, there were two observations that, like the size difference, were invariably noted. The first was that the gut of young worms was open (Plates 36 and 37) and usually full of a variety of material; while the guts of the damaged worms were either tightly closed or only just patent. The second difference was that the contents of the gut of damaged worms contained peculiar whorls which were persistently seen within the lumen and could represent lipoproteins, or immune complexes (Plates 38 and 39). They were never found in the guts of normal worms.

The microvilli of both 10 and 22 day-old worms were similar in size and numbers, though the damaged cells beneath them showed reduced activity (Plates 40 and 41). To sum up, apart from the general signs within the damaged worms suggestive of malnutrition, the only invariable findings which occurred in all specimens examined were an overall difference in size and the presence of whorls within the closed guts of damaged worms.

Experiment 4.3:

The study below was conducted to identify and endeavour to define the plugs of precipitates observed, repeatedly, near the peri-oral area of worms (Plates 42 and 43) during initial and challenge infections, as well as the peculiar whorls,^{seen} under the EM,



Plate 32 : Electron micrograph of section through anterior third
of a 10 day-old *S. ratti*. C = cuticle; G = Golgi complex;
M = mitochondria (x 29,000)




Plate 32 : Electron micrograph of section through anterior third of a 10 day-old S. ratti. C = cuticle; G = Golgi complex; M = mitochondria (x 29,000)



Plate 33 : Electron micrograph of section through anterior third of a 22 day-old S. ratti. V = vacuolated space (x 30,000)



Plate 33 : Electron micrograph of section through anterior third of a 22 day-old S. ratti. V = vacuolated space (x 30,000)




Plate 34 : Electron micrograph of a section through the gut cells of 10 day-old S. ratti. ER = endoplasmic reticulum (x 23,000)




Plate 35 : Electron micrograph of a section through the gut cells of 22 day-old S. ratti. L = lysosomes (x 23,000)




Plate 34 : Electron micrograph of a section through the gut cells of 10 day-old S. ratti. ER = endoplasmic reticulum (x 23,000)




Plate 35 : Electron micrograph of a section through the gut cells of 22 day-old S. ratti. L = lysosomes (x 23,000)




Plate 36 : Electron micrograph of the gut contents of 10 day-old S. rattus. il = intestinal lumen (x 16,000)



Plate 37 : Electron micrograph of the gut of 22 day-old S. rattus, note the non-patent lumen (x 16,000)




Plate 36 : Electron micrograph of the gut contents of 10 day-old *S. rattii*. il = intestinal lumen ($\times 16,000$)




Plate 37 : Electron micrograph of the gut of 22 day-old *S. rattii*, note the non-patent lumen ($\times 16,000$)




Plate 38 : Electron micrograph of the peculiar whorls observed within the intestinal lumen of 22 day-old S. ratti. Lp = lipoprotein structures; Mv = microvilli (x 75,000)




Plate 39 : Electron micrographs of similar whorls (x 75,000)



Plate 38 : Electron micrograph of the peculiar whorls observed within the intestinal lumen of 22 day-old *S. ratti*. lp = lipoprotein structures; mv = microvilli (x 75,000)




Plate 39 : Electron micrographs of similar whorls (x 75,000)




Plate 40 : Electron micrograph of a section through the gut of 10 day-old S. ratti, to show microvilli.
Mv = microvilli (x 23,000)




Plate 41 : Electron micrograph of a section through the gut of 22 day-old S. ratti (x 23,000)




Plate 40 : Electron micrograph of a section through the gut of 10 day-old S. ratti, to show microvilli.
Mv = microvilli (x 23,000)




Plate 41 : Electron micrograph of a section through the gut of 22 day-old S. ratti (x 23,000)

to a specific immune reaction against S. ratti, and a non-specific inflammatory response which lasts for a short period and can expel and damage other intestinal nematodes when introduced subsequently, e.g. N. brasiliensis and T. spiralis (Louch, 1962; Kazacos and Thorsen, 1975; Kazacos, 1976; Mogbel and Moloney, 1976, unpublished observations). The fact that only 5% of the worms transferred from a challenge infection established does not substantiate the conclusion that adaptation occurred in this system. On the other hand, the persistence of these worms in the gut for 15 days after transplantation might suggest otherwise. This aspect of the relationship between worms from challenge and clean rats is worth pursuing. If adaptation was confirmed, S. ratti would be only the second nematode to manifest this phenomenon, N. brasiliensis being the first.

The changes observed in the gut cells of damaged worms were basically similar to those observed in N. brasiliensis and T. spiralis (Ogilvie and Hockley, 1968; Love *et al.*, 1976), particularly with regard to the large number of vacuoles and the distorted endoplasmic reticulum. However, the lipid droplets described in the gut cells of N. brasiliensis by Januar (1966) and Ogilvie and Hockley (1968), were not seen in damaged S. ratti worms.

The cuticle of damaged worms was intact, which indicated that antibodies which caused internal disorganisation entered the nematode through the body orifices, and most probably, the mouth.

The invariable discovery of peculiar whorls in the lumen of the gut of damaged worms could be a confirmation of observations by Sarles (1938) and Taliaferro and Sarles (1939) in N. brasiliensis, under light microscopy. It is possible that these bodies are ingested immune complexes (Ab/Ag), but due to the minute size of the lumen of the gut, no specific fluorescence was detectable under the UV microscope. It is doubtful whether these bodies, even if they are immune complexes, have any role in worm damage, since immune complexes act by activating complement, and through their chemotactic character, they can be involved in opsonisation. It is unlikely that this can occur within the worm. These bodies may have been formed following the ingestion of anti-worm antibodies

Similar changes occur in other parasitic nematode systems (Soulsby, 1962; Ogilvie and Hockley, 1968; Ogilvie and Jones, 1971, 1973; Ogilvie^{and worms}, 1976; Love et al., 1976).

Normal S. ratti transferred before being fully exposed to the immune response of rats, were able to continue their egg production. Worms taken from primary infections, after day 20, and those from the early stages of a secondary infection, were the same length. Worms which were transferred after day 22 PI were unable to resume their egg production and did not survive in non-immune hosts. This indicated that, as with N. brasiliensis (Ogilvie and Hockley, 1968), the damage to S. ratti from late initial infection was irreversible.

However, adult S. ratti collected 7 days after a challenge survived for 15 days in non-immune rats, three days longer than normal worms did, and showed preliminary manifestations of "adaptation" as described by Ogilvie and Hockley (1968) and Ogilvie (1972). These worms did not regain their original length but managed to resume egg production, with high fecundity, for a substantial length of time, similar to the transferred normal worms.

It is possible that adaptation may occur in S. ratti during a challenge infection, however this, in the light of the results obtained in experiment 4.1, is still arguable.

As mentioned (see introduction to this chapter), adaptation of N. brasiliensis was suggested to occur during the L3/L4 moult in the lungs (Ogilvie, 1972) where there was little response against the migrating larvae. This evasion of the immune response led to the establishment of worms which were capable of surviving and resuming egg production when transferred to naive rats (Ogilvie and Wilson, 1976).

As far as S. ratti is concerned, it has been established that no moulting occurs in the lungs of infected rats (Wertheim and Lengy, 1968). Moreover, there is a severe response against the pulmonary stages of a challenge infection (see Chapter 3), resulting in the establishment of very few stunted worms in the gut. The possibility that adaptation in S. ratti might occur during the L3/L4 moult in the gut, cannot be excluded, but the few worms which manage to evade the strong immune response in the lungs may face a more severe reaction upon their arrival in the gut. This is due



Plate 42 : Anterior end of adult, 10 day-old *S. ratti*, with no plug of precipitate around the mouth (x 400).




Plate 43 : Anterior end of adult, 22 day-old *S. ratti*; note the peri-oral precipitate (x 400).



Plate 42 : Anterior end of adult, 10 day-old *S. ratti*, with no plug of precipitate around the mouth (x 400).



Plate 43 : Anterior end of adult, 22 day-old *S. ratti*; note the peri-oral precipitate (x 400).



Plate 44 : 10 day-old S. ratti showing fertile embryonated eggs (E) within the uterus (U). V = vulva (x 400)



Plate 45 : 22 day-old S. ratti showing absence of eggs (phase contrast x 400)




Plate 44 : 10 day-old *S. ratti* showing fertile embryonated eggs (E) within the uterus (U). V = vulva (x 400)



Plate 45 : 22 day-old *S. ratti* showing absence of eggs (phase contrast x 400)

inside the gut lumen of damaged worms.

Normal (10 day-old) and damaged (22 day-old) worms were washed three times in phosphate buffer saline (PBS) with continuous agitation, and then run through a direct fluorescent antibody test (FAT). The washed worms were incubated in fluorescein isothiocyanate (FITC) conjugated to goat anti-rat immunoglobulin (Wellcome Reagents Ltd.) directly, at a working dilution of 1:20 and 1:40 in 37°C incubator for 30 minutes. The excess FITC was washed off with PBS and the worms were mounted on slides in buffered glycerine. The slides were examined under an Ultra-violet (UV) microscope (Nikon).

The normal worms had no precipitates near their mouth or any other orifice, and the anterior part showed no fluorescence. Damaged worms had precipitates which varied in shape and size, and they fluoresced strongly in the direct FAT at both dilutions.

For the study of the peculiar bodies within the lumens of damaged worms, normal and damaged worms were washed three times in sterile PBS after being killed by cooling. Sections of adults were prepared by freezing 300 worms of each age in an embedding medium for frozen tissue specimens (Ames O.C.T. Compound) on to a metal block. Sections were cut at 6 μ m in a cryostat microtome at -20°C and were attached to coated multispot slides (C. A. Hendley & Co., Essex). The sections were left in a dessicator at 4°C overnight and were subsequently fixed with acetone for 5 seconds immediately before running direct FAT, using FITC conjugated to goat anti-rat immunoglobulin. Even under high magnification, no fluorescence was seen within the very small damaged worms.

Discussion:

The experiments in this chapter have demonstrated that adult S. ratti are affected, at both macro- and micro-levels, by the immune response of rats during the course of primary and secondary infections. The length of adult worms, which decreased with the progress of initial infection (see Chapter 3), was accompanied by internal cyto-pathological, as well as morphological, changes.

which react with internal metabolic secretions and form complexes within the lumen of the gut.

It is likely, however, that these bodies are associated with the plugs of precipitate repeatedly seen attached to the mouth of damaged worms in vivo. The fluorescence of the latter structures in a direct FAT suggests they were immune complexes. Similar plugs were observed in other nematodes particularly in vitro (Sarles, 1938; Taliaferro and Sarles, 1939; Lawler, 1940; Oliver-Gonzalez, 1940; Lukashenko, 1960). However, their role in worm damage is still controversial and worth pursuing, particularly since we know very little about the mechanism(s) of damage and worm deterioration. Ogilvie (1964) suggested that these precipitates were not significant in protection as far as the host was concerned, but they were recognised as being antibody/antigen complexes (Ogilvie and Jones, 1973). It is very likely that worms bring about the immune attack by their own activities. The early production of antibodies to the metabolic presence of the worms controls the production and modulation of the enzymatic activity of the worms, particularly that of acetylcholinesterase (AChE) (Jones and Ogilvie, 1972) which has been shown to be antigenic (Sanderson, 1969; Lee, 1970; Edwards et al., 1971; Jones and Ogilvie, 1972). Soon after AChE is released, specific Ab/Ag complexes develop near the site of production.

It is proposed here that worm damage may be partially instigated by the development of precipitates internally and around the mouth which blocks the worm's alimentary canal and prevents the usual food uptake leading to starvation. The changes within the worms reflect this, as most of the cellular contents were consumed by the worms through the action of lysosomes.

It is essential that, as a follow up to this work, studies should be conducted on the metabolic changes which occur in adult S. ratti during the course of infection similar to those shown in N. brasiliensis by Henney et al. (1971), as well as the activities of AChE in S. ratti which have been shown to be significant in many other nematodes (Rothwell et al., 1973; Ogilvie et al., 1973; Bremner et al., 1973; McLaren et al., 1974).

The role of the different parasitic stages of *S. ratti*
in stimulating the immune response

Different methods have been employed to determine the immunogenic stages of worm infections. These included irradiation of different stages in order to arrest their growth, particularly to sterilise infective larvae to prevent them reaching sexual maturity (Levin and Evans, 1942; Hendricks, 1950, 1952; Gould *et al.*, 1955; Hartung and Becht, 1960; Jarrett *et al.*, 1960; Zaiman *et al.*, 1961; Jennings *et al.*, 1963; Miller, 1964, 1966). Using this method, Kim (1957) concluded that the intestinal phase of *T. spiralis* had a significant immunogenic role.

An attempt was made by Chala and Murta (1967) and followed by Rego and Kahtalian (1972), to immunise rats against irradiated third stage infective larvae of *S. ratti*, with varying degrees of success (see discussion).

Chemically abbreviated infections have also been used to determine which stages were immunogenic. This has proved to be a convenient technique since anthelmintics are available which affect one or more stages of different helminths. This method has been extensively used for the determination of the immunogenic stages of *T. spiralis* (reviewed by Zaiman, 1970), *T. muris* (Wakelin, 1969), *Oesophagostomum radiatum* (Roberts *et al.*, 1963) and *Haemonchus placei* (Roberts and Keith, 1959a, b).

Sensitization of the host with a specific stage of the infection by transplantation, into the host, of these stages was also carried out in this context. This method was used by Ogilvie (1965) to investigate the role of adult worms in stimulating an immune response in rats against *N. brasiliensis*. Muscle larvae of *T. spiralis* were put in diffusion chambers and were implanted in rats to investigate their immunogenic role (Despondier, 1971).

Using each of these methods or a combination of more than one (Danham, 1966b), it was shown that adult *T. spiralis* and

N. brasiliensis are important immunogenic stages (Campbell et al., 1963; Campbell, 1965; Denham, 1966 b; Ogilvie, 1965). More protection was obtained by transplanting female N. brasiliensis (Ogilvie, 1965). Both males and females stimulated the same degree of immunity in T. spiralis infection (Anderson and Leonard, 1940) and in Schistosoma mansoni infection in monkeys (Smithers, 1962).

Stage specificity has been demonstrated in T. spiralis infections and a "dual-antibody" hypothesis suggested to support it (Oliver-Gonzalez, 1941). Recently, James and Denham (1975) reiterated this theory and demonstrated a stage specific immune response in T. spiralis in mice.

Immunity elicited by adult N. brasiliensis worms was not stage specific, since larvae developing in rats previously infected with adult worms alone were affected by the immune response (Ogilvie, 1965 a). In normally infected rats, reinfection affected both the larvae during their tissue migration and the adult worms in the gut (Ogilvie, 1965 b).

No stage-specificity was demonstrated in T. muris infections in mice. Using chemically abbreviated infections, the immune reaction was elicited by, and acted against, all stages of infection (Wakelin, 1969).

In the following study the role of both infective larvae and adult S. ratti in the induction of acquired immunity in rats is investigated using chemically abbreviated infections and transplantation of adult worms. Thiabendazole (TBZ) is very effective against the intestinal stages of S. ratti, particularly the parthenogenetic females, but has no effect on the migratory larval stages (Chais, 1963; Persand and Grant, 1972).

Experiment 5.1:

In this experiment, chemically abbreviated infections were used to assess the role of the migratory stages of infection with S. ratti in inducing protection against reinfection. Fourteen female Wistar rats were divided into two groups: The first group was infected with 1000 (\pm 94) L3 on day 0. TBZ was administered per os at 100 mg/kg

on days 4, 5, 6 and 7 PI to eliminate intestinal worms as soon as they arrived in the gut, and to confine the infection to the migratory stage. The rats of the second group acted as challenge controls and were treated with TBZ on the same days. On day 8 PI, two rats from the first group were killed. No worms or larvae were recovered from their intestines showing that TBZ had eliminated the infection. On day 9 PI the two groups were challenged with 1000 (\pm 101) L3/rat. All rats of both groups were killed 12 days later. The results of the worm counts are shown in Table 12.

There was no statistically significant difference between the number of adult worms recovered from the two groups.

Experiment 5.2:

In this experiment, another approach was used to determine the role of infective larvae in inducing immunity to reinfection. Three injections of 2000 freeze-thaw killed L3s were given intravenously to a group of five rats at two days intervals. The experimental and the challenge control group (5 rats) were then challenged with 1000 (\pm 78) viable infective larvae. The rats of both groups were killed on day 10 PI and the number of adult worms determined. The results are shown in Table 13. There was no significant difference between the numbers of adult worms recovered in both groups.

The two previous experiments suggest that there was no significant role of the migratory stages of S. ratti in stimulating resistance to reinfection.

Experiment 5.3:

The object of this experiment was to determine the role of transplanted adult S. ratti in inducing a protective immune response. Ten 120 gm rats were infected with 3000 L3 each. They were killed

Table 12 Exp: 5.1: Adult worm count after a challenge infection
with S. ratti following chemically
abbreviated sensitization to the migratory
stages.

Sensitized Group	Challenge control
370	682
363	608
658	613
568	521
498	391
376	333
472 (\pm 50.3)	524 (\pm 56)

Sensitized group vs Challenge control N. S.

Table 13 Exp: 5.2: Adult worm count after a challenge infection with S. ratti, following sensitization by serial IV injections with Freeze-thaw killed L3s.

Sensitized Group	Challenge control
631	498
599	531
500	507
612	473
587	510
585 (\pm 22.6)	503 (\pm 9.4)

Sensitized group vs Challenge control N.S .

and the adult worms collected ten days later. 300 adult worms in 0.4 ml volume of saline were transplanted, intraduodenally, into each of 16 rats. A control group of 16 rats was infected with subcutaneous injections of 300 L3/rat on the same day as the donor rats were initially infected. One other group of 4 rats was left uninfected to act as challenge controls. Four rats were killed from both the group given adult worms and the normal infection group on days 5, 10 and 15 post-transplantation and the number of adult worms were counted. On day 15 post-transplantation, the remaining 4 rats from the experimental, normal infection and uninfected control groups were treated with a single dose of TBZ at 200 mg/kg. Three days later all the rats were challenged with 1000 (\pm 107) L3s. Rats of all three groups were killed on day 10 post-challenge and the number of adult worms counted. The results are summarized in Table 14. In this experiment very few transplanted worms (5.1% on day 5, 20% on day 10 and 5% on day 15 post-transplantation) established themselves in clean rats. The normal infection control showed a development rate of 59% on day 15 PI. On day 10 post-challenge, the group of rats which were sensitised to 300 adult worms only harboured 448 worms, while the group which received a subcutaneous infection of 300 infective larvae harboured 79.2 worms. The challenge controls had a mean of 411 worms on the same day.

No protective immunity was stimulated by the transplantation of 300 worms, but a full infection with 300 infective larvae produced a substantial degree of resistance to reinfection.

Experiment 5,4:

The protocol of this experiment was similar to that of the previous one with the exception that the number of transplanted adult worms was increased to 700 per rat in 0.1 ml volume of saline. Ten recipient rats received these worms intraduodenally, while the infection control group was infected with a normal subcutaneous infection of 700 L3s on the day the donors were initially infected. Four rats were left uninfected to act as a challenge

Table 14 Exp: 5.3 Mean adult worm recovery in three groups:

A: received an adult infection only (300 transplanted intraduodenally)

B: normally infected with 300 infective larvae (same day as the donor rats)

C: challenge control group

Days post - transplantation	Days PI	A adult infection only (300)	B normal infection (300 L3)	C challenge control
5	15	15.5 (<u>+</u> 4.19)	177 (<u>+</u> 13.5)	-
10	20	60 (<u>+</u> 22.2)	95.2 (<u>+</u> 7.44)	-
15	25	9.25 (<u>+</u> 4.23)	15.2 (<u>+</u> 3.7)	-
Day 10 post challenge		448 (<u>+</u> 46.6)	79.2 (<u>+</u> 11.7)	411 (<u>+</u> 44.9)

control group. Four rats from the experimental group were sacrificed on day 5 post-transplantation and they harboured a mean of 281 (\pm 32) worms. Faecal cultures made from the two infected groups showed similar levels of larval production on days 0, 3, 6, 10 and 15 post-transplantation (equivalent to days 10, 13, 16, 20 and 25 PI). The rats in all three groups were given a single dose of TBZ at 200 mg/kg per os 15 days after transplantation. No worms were recovered from two rats killed two days after anthelmintic treatment. Four days after TBZ all rats were challenged with 1000 (\pm 89.9) L3s. The results are presented in Tables 15 and 16 and Figures 17 and 18. Faecal cultures from all infected groups made on days 6 and 8 post-challenge revealed no larvae in the rats given either a normal infection or transplanted adult worms, whilst the normal control group had an average of 5850 L3/gm of faeces.

All rats were killed 10 days after challenge and the adult worms counted. There was a recovery of 4.5% from the group "immunised" with adult worms and 3.5% from the group previously given a full infection. The challenge control harboured 41.6% of the inoculum and these worms were of normal length, while those of the other previously sensitised groups were stunted.

In this experiment, the transplanted worms established themselves better than in the previous experiment and there was a highly significant difference between the recovery rate in the group which was sensitised with adult worms alone and in the challenge control group. Rats normally infected with a full infection manifested a strong degree of resistance to reinfection.

This experiment suggests that adult *S. ratti* are a major source of protective antigens, but a full infection was more immunogenic.

Discussion:

Due to the complex nature of the antigens produced by the different stages of nematode infections, it is very difficult to identify the precise immunogenic stage of the life cycle of these worms.

Table 15 Exp: 5.4 Mean larval production in three groups during initial and challenge infections

A: received transplanted adult infection only (700 adults)

B: received normal infection (700 adults)

C: challenge control

Day	Mean (SE) number of larvae		
	A adult infection only	B normal infection	C challenge control
10	4080 (+ 792)	6693 (+ 478)	0
13	2150 (+ 220)	4165 (+ 143)	0
16	720 (+ 168)	1874 (+ 155)	0
20	204 (+ 39.2)	458 (+ 59.3)	0
25	33.3(+ 22.2)	192 (+ 27)	0
reinfection on day 30			
36	0	0	5569 (+ 421)
38	0	0	6122 (+ 282)

Table 16 Exp: 5.4

Adult worm recoveries in rats autopsied 10 days after challenge with 1000 L3 / rat in three groups:

A: initially infected with 700 adult worms only

B: initially infected with 700 L3 (normal course)

C: challenge control

	Group A	Group B	Group C
Individual worm recoveries	31	23	413
	71	28	393
	59	51	473
	21	40	388
Mean (SE) adult recovery	45.5 (\pm 11.7)	35.5 (\pm 6.27)	416.75 (\pm 19.5)
Number of worms measured	39	29	44
Length of worms (mm)	1.39 (\pm .03)	1.22 (\pm .02)	2.79 (\pm .03)

Figure 17, Exp. 5.4:

- Larval production in three groups of rats.
●—● infected with 700 transplanted worms only
○—○ infected with 700 infective larvae
■—■ challenge control





Figure 18, Exp. 5.4:

Worm length and number of adult worms recovered 10 days after challenge in three groups.

- initially infected with 700 adult worms only and challenged
- ▨ initially infected with 700 L3/rat and challenged
- challenge control



The results of the experiments in this chapter have suggested that adult S. ratti were more immunogenic than infective and migratory larvae.

Soulaby (1958) suggested that L3/L4 moult was the first major source of protective antigens. In S. ratti, this moult occurs in the intestine (Wertheim, 1970). It is possible that TBZ either affected the larvae immediately after their arrival at the intestine, or following their moult to L4 and L5. The time lapse between moulting and evacuation of these worms was, perhaps, not sufficient for the antigens produced to be recognised by the immune system.

Repeated injections of freeze-thaw killed infective larvae of S. ratti conveyed no immunity to reinfection. It was hoped that, since these were injected IV, they would be arrested in the lungs and subsequently produce a response against larvae of a challenge infection. Freeze-thaw killed new born larvae of T. spiralis produced no noticeable immunity whereas viable new born larvae, which were allowed to develop naturally into muscle larvae, produced good immunity (Despondier, 1971).

It seems that sensitisation with non-living nematode material is not effective in protection against a challenge. Live stages and their metabolic products are required to induce a significant immune response. Only Sheldon (1937a) reported successful immunisation of rats against S. ratti using heat-killed infective larvae.

Chala and Murta (1967) were able to demonstrate a degree of immunisation following 3 injections of 10,000 irradiated larvae of S. ratti/S. venezuelensis at 15 day intervals. On the other hand, 50-54,000 irradiated S. ratti larvae conferred no immunity against challenge in rats (Rego and Kahtalian, 1972). Stankiewicz et al. (1971) successfully immunised rabbits with UV-irradiated infective larvae of S. papillosum and these larvae did not develop to maturity.

In order to assess accurately the role of infective larvae only, in stimulating protective immune response, it is vital that no adult worms reach the intestine. Adult worm antigens, whether they are shared with the larval stages or not, undoubtedly interfere with the development of the immune response against larvae. Sinaki and Krzysztofowicz (1971) immunised rabbits against S. papillosum

by exposing them to repeated doses of intact infective larvae at different intervals. It seemed that immunisation was obtained when the challenge was introduced after a period which allowed the development of adult stages in the intestine.

Although a full infection with S. ratti induced the most effective immune response against reinfection (see also Chapter 3), adult worms alone were capable of producing similar resistance. 10 day-old S. ratti transferred from donors into clean recipients survived just as if they were in their original hosts.

The worms transplanted during the first experiment did not convey any protective immunity. This was probably due to the low number used and the relatively large amount of saline used during transfer.

No stage specificity was observed in S. ratti infection since immunity produced by an initial infection acted on all stages of infections (see Chapter 3) and it is possible that transplanted adult worms alone also produced similar effects.

CHAPTER 6Studies on the mechanism of expulsion of *S. ratti*
from the intestines of rats

It is now well established that worm expulsion from the host's intestine is the consequence of an immunological reaction and is not due to the worm's senescence, as worms can survive for a long time with a steady level of fecundity when transferred from one rat to another (Ogilvie and Hockley, 1968) and fertile worms persist in hosts treated with immunosuppressive drugs (Ogilvie, 1965b).

With a classical sense of prediction, Taliaferro and Sarles (1939) and Taliaferro (1940) concluded that the process of worm elimination in *N. brasiliensis* infections in rats was mediated by a humoral response with subsequent secondary cellular co-operation. Some 25 years later, it was suggested that local anaphylaxis was the major event which led to worm damage (Mulligan et al., 1965; Urquhart et al., 1965; Barth et al., 1966). Since then, this view has been drastically altered in favour of Taliaferro and Sarles' hypothesis of a more complicated sequence of events. It is now generally believed that nematodes are eliminated from the gut, in rodent parasitic systems, through intricate mechanisms, which involve both areas of the immune response.

The development of active immunity to *N. brasiliensis* and its subsequent expulsion from the host appears to be thymus-dependent (Wilson et al., 1967; Ogilvie and Jones, 1967). Neonatally thymectomised and adult thymectomised rats treated with anti-thymocyte serum (ATS), harboured more adult *N. brasiliensis* than sham-operated controls (Kelly, 1972; Keller and Keist, 1972). Similar results were obtained by treating infected rats with ATS daily (Kelly, 1972).

Neonatal thymectomy, anti-lymphocyte serum (ALS) and whole-body irradiation suppressed the expulsion of *T. muris* in mice, suggesting that this response was associated with thymus-derived lymphocyte population (T. cells) (Wakelin and Selby, 1974b).

ATS prevented rats and mice from developing protective

immunity against T. spiralis (Ruitenber*g et al.*, 1974; Walls *et al.*, 1973). On the other hand, antibodies against T. spiralis develop in both thymectomised and intact mice (Ljungström and Ruitenber*g*, 1976). Ljungström and Ruitenber*g* (1976) suggested that thymus-dependent and thymus-independent antibodies were formed during infection with T. spiralis, but that the latter were not protective and may have acted as blocking antibodies.

Larsh and his co-workers (Larsh *et al.*, 1952; Larsh and Race, 1954; Larsh *et al.*, 1956) emphasised the role of the inflammatory process before and during T. spiralis expulsion. Cortisone treatment of mice infected with this worm considerably delayed the expulsion of adult worms (Coker, 1956a; Markell, 1958). Coker (1956b) suggested that expulsion required a primary antibody action with secondary cellular co-operation. However, Campbell (1968) demonstrated that non-steroid anti-inflammatory drugs had no effect on expulsion and concluded that intestinal inflammation was not a cardinal part of the mechanism of worm expulsion.

There has been some controversy concerning the role of antibodies in worm expulsion. Passive transfer of immunity with serum was employed early in experimental nematode infections, with a varying degree of success. Although immune serum conferred immunity to N. brasiliensis (Sarles and Taliaferro, 1936; Chandler, 1938b; Taliaferro, 1940), Ogilvie and Jones (1968) showed that only a few of their pools of immune serum were protective against this parasite.

Immunity to T. spiralis was transferred successfully by serum on many occasions resulting in decreased worm burdens (Culberston and Kaplan, 1937, 1938; Oliver-Gonzalez, 1941; Culberston, 1942, 1943; Dorin, 1946; Hendricks, 1953; Mills and Kent, 1965), but this was not always consistent (McCoy and Bond, 1941; Danham, 1969; Larsh *et al.*, 1970).

Antibodies of the class IgG1 have been demonstrated following active T. spiralis infection or immunisation against stichosome granules (Despondier and Muller, 1970; Crandall and Crandall, 1972). Stichocytes ^{and are} contain immunogenic granules found in the anterior third of T. spiralis (Despondier and Muller, 1970). Passive immunity from mother to baby, via milk, was also reported in mice infected with T. spiralis (Duckett *et al.*, 1972; Perry, 1974).

It is now generally thought that the major function of antibodies is to induce worm damage (reviewed by Ogilvie and Jones, 1971, 1973; see Chapter 4), but there has been considerable conflict over the role played by the various types of cells in worm expulsion.

Transfer of immune spleen cells had little or no effect in conveying immunity from sensitised to non-sensitised animals against the intestinal stages of nematodes (Hunter and Leigh, 1961; Ogilvie and Jones, 1968). However, mesenteric lymph node cells (MLNC), which are closely associated with the infected gut, instigated the abrupt elimination of an initial infection with N. brasiliensis (Keller and Keist, 1972; Dineen et al., 1973a, b; Kelly et al., 1973; Dineen and Kelly, 1973a; Ogilvie and Love, 1974), T. spiralis (Larsh et al., 1964a, b; Larsh, 1967; Love et al., 1976; Wakelin and Lloyd, 1976b) and T. muris (Selby and Wakelin, 1973; Wakelin, 1975a).

Jones and Ogilvie (1971) proposed that expulsion of N. brasiliensis from rats was achieved in two sequential steps. The first, antibody-mediated, step damaged the worms structurally (Ogilvie and Hockley, 1968; Lee, 1969a) and metabolically (Edwards et al., 1971). The second step involved components of a non-specific inflammatory process, which correlated with a rise in mast cell infiltration and amine levels (Barth et al., 1966; Keller, 1970), but it was later shown that inflammation and mast cells do not play an important role in this step, in N. brasiliensis infections (Keller, 1971; Keller and Keist, 1972; Kelly and Ogilvie, 1972).

Keller and Keist (1972) showed that functionally intact lymphocytes were necessary for the second step of worm expulsion. In order to assess the overall role of cellular components in immunity against N. brasiliensis, Dineen et al. (1973b) studied the effects of transferred sensitised and non-sensitised lymphocytes on transplanted, damaged, worms in both irradiated and non-irradiated recipients. Sensitised cells caused a significant reduction in the worm burden in both groups of recipients, which demonstrated the definite role of a committed population of lymphocytes in this reaction.

Kelly et al. (1973) investigated the effect of sublethal and

lethal whole-body irradiation on the capacity of sensitised MINC to expel damaged worms, in syngeneic rats. A substantial degree of immune competence was demonstrated in sublethally irradiated rats, as measured by the production of plaque forming cells (PFC) against sheep red blood cells. This suggested that in non-irradiated and sublethally irradiated rats, the capacity for active participation in worm expulsion was present.

Dineen and Kelly (1973b) transferred ^{immune} MINC and normal bone marrow cells to rats exposed to a lethal dose of irradiation. These components caused a significant reduction in the number of adult worms recovered five days later. They concluded from this that after antibody damage, the expulsion of worms required two major components. These being sensitised T. cells, which require the co-operation of a non-specific population of myeloid, radio-sensitive, bone marrow-derived cells which are able to undergo rapid regeneration.

In the host-parasite relationship between mice and T. spiralis, immunity was transferable by sensitised cells from the peritoneal exudate and lymph nodes (Larsh *et al.*, 1964a, b). From these experiments, Larsh and his group claimed that delayed hypersensitivity (DH) was involved in the process of worm elimination.

In a recent review, Larsh and Race (1975) have reiterated their emphasis on the causal relation of DH to the expulsion of adult T. spiralis, and totally excluded the participation of a humoral response in this phenomenon. Love *et al.* (1976) used the experience derived from their previous work on N. brasiliensis infections, in both rats and mice to study the expulsion of T. spiralis from rats. The picture was very similar to that of N. brasiliensis in that both antibodies and sensitised MINC were required to bring about worm expulsion. Structural damage of adult worms was also observed prior to their expulsion and when the host received antiserum or cells. Similar results were recently obtained in mice by Wakelin and Lloyd (1976a, b). In T. colubriformis infections in the guinea-pigs, cellular transfer of immunity has been successful (Wagland and Dineen, 1965; Dineen and Wagland, 1966), leading to the expulsion of these worms before they reach maturity. Less success was achieved with passive transfer of antibodies

(Connan, 1972). Ogilvie and Jones (1973) proposed that passive or adoptive immunisation against T. colubriformis in the guinea-pig had an enhancing effect on the active immune response which is already developing in this abnormal host. In T. cell-deprived guinea-pigs, the infection was long lived (Dineen and Adams, 1971).

Immunity against T. muris was transferable through both antiserum and sensitised cells (Selby and Wakelin, 1973) though the former was more effective. Further work by Wakelin (1975) revealed that a sequential action of antibody-mediated responses and lymphoid cell-mediated immunity was required for the expulsion of T. muris. Although this field requires further elucidation, there is no doubt that work on N. brasiliensis, T. spiralis, T. colubriformis and T. muris has shown that a multi-step mechanism is involved in the process of worm expulsion (Ogilvie and Jones, 1971; Love et al., 1976; Wakelin and Lloyd, 1976; Rothwell and Dineen, 1973; Wakelin, 1975).

The experiments reported in this chapter are an attempt to analyse the role played by the cellular and humoral response in the elimination of S. ratti from rats. This was achieved by passive and adoptive transfer of immunity from competent donors to non-sensitised rats.

Collection of serum:

25 days PI rats were anaesthetized with Nembutal and bled, by cardiac puncture, using 5 ml syringes and 1½ inch x 21 gauge needles. The blood was allowed to clot at room temperature for 1 hour then at 4°C for three to four hours. The samples were pooled and then stored in 1 ml aliquots at -70°C. In serum transfer each rat received a total volume of approximately 5 ml of serum. Each experiment had its pool of serum obtained from a group of donors on day 25 PI.

Collection of sensitised immune competent lymphocytes:

Donor rats were infected with 3000 L3 each and killed on day 25 PI. The enlarged mesenteric lymph nodes were removed and kept in medium 199 at pH 7.2 on ice and were then gently teased apart and pushed through a sterile stainless steel tea-strainer with a 10 ml syringe plunger. The suspension of lymphocytes was pipetted through a sterile glass wool column to eliminate the adipose cells and the macrophage population. The MLNC suspension was collected on ice by washing the column twice with chilled medium 199. The cells were concentrated by centrifugation at 1000 rpm for 4 minutes at 4°C. The cells were washed 3 times with medium 199 and the volume of the suspension made up to the required amount prior to injection. After dilution, lymphocytes were counted in a "Neubauer" haemocytometer. The viability of the MLNC was determined by the trypan-blue exclusion test.

Recipient rats received 2×10^8 MLNC each intravenously (IV) through the lateral caudal vein with a 1 ml graduated disposable syringe fitted with a $\frac{1}{4}$ inch x 25 gauge needle. When the MLNC were injected IV, rats were restrained in specially prepared perspex boxes. Three different sizes of boxes were made for different sizes of rats (Plate 46). The rats' tails were accessible for IV inoculation (Plate 47).

Experiment 6.1:

This experiment was designed to establish whether sensitised lymphocytes cause an early expulsion of the worm population, whether immunised serum causes the expulsion of the worm burden and to determine whether immunised serum and immunised lymphocytes cooperate to induce expulsion of adult S. ratti.

50 inbred female Wistar rats were infected with 2000 L3 each and killed on day 25 PI when serum and MLNC were collected. The average viability of MLNC was 70% and the concentration of cells was adjusted so that 2×10^8 viable cells were given to each recipient rat. The mean worm burden of the donor rats, on day 25,




Plate 46 : Three different sizes of restraining boxes for rats



Plate 47 : Intravenous injection of MNC into the caudal vein of a rat




Plate 46 : Three different sizes of restraining boxes for rats



Plate 47 : Intravenous injection of MNC into the caudal vein of a rat.

counted in a sample of four rats from each donor group was 11.7 (\pm 6.1). Ten weeks-old inbred female Wistar rats were used as recipients.

The recipient rats were divided into the following major groups and each rat infected with 1000 L3 on day 0:

Group 1: This group was divided into 3 sub-groups; and all rats were infected on day 0 with 1000 (\pm 39) infective larvae.

- A: Received immune serum on days 3, 6, 9 and 12 PI, and immune MLNC on day 6 PI (5 rats)
- B: Received immune MLNC on day 6 PI (5 rats)
- C: Normal infection control (4 rats)

Group 2: This was also divided into three sub-groups of rats which were infected on day 0 with 1000 (\pm 86) infective larvae.

- A: Received immune serum on days 3, 6, 9 and 12 PI and immune MLNC on day 12 PI (5 rats)
- B: Received immune MLNC on day 12 PI (5 rats)
- C: Normal infection control (4 rats)

Group 3: This contained two sub-groups of rats which were infected on day 0 with 1000 (\pm 96) infective larvae.

- A: Received immune serum only on days 3, 6, 9 and 12 PI (5 rats)
- B: Normal infection control (5 rats)

Faecal cultures were made from all groups on days 8, 12 and 13 PI and all the recipient rats autopsied on day 16 PI. This day was chosen because the worms do not show signs of "damage" before day 18 PI and expulsion usually commences after day 22, and thus it was a good monitor of the effects of the transferred immune components on the course of infection.

The results of this experiment are presented in Table 17 and Figures 19 and 20.

Table 17 Exp: 6.1 Effects of the transfer of immune MNC and serum from sensitised donors to recipients at different intervals, during the course of infection with *S. ratti*

Group	Days of transfer		Mean (SE) number of larvae/gm faeces on days:			Mean (SE) number adult worms on day 16	Percentage recovery	Number of worms measured	Mean (SE) length of worms (mm)
	Serum total/ rat 5.3 ml	MNC viability 70%	8	12	15				
A1	3,6,9,12	6	12692 (+1293)	2187 (+191)	866 (+124)	681 (+161)	68.1%	50	1.33 (±.01)
A2	-	6	8272 (+560)	3970 (+247)	2146 (+238)	166 (+64.4)	16.6%	50	2.92 (±.05)
A3	-	-	4028 (+225)	4596 (+430)	2566 (+169)	372 (+26.5)	37.2%	40	2.85 (±.04)
B1	3,6,9,12	12	4714 (+522)	2554 (+338)	3306 (+233)	145 (+25.6)	14.5%	50	1.79 (±.03)
B2	-	12	2852 (+208)	1772 (+222)	1976 (+298)	96.8 (+30.1)	9.6%	34	2.72 (±.04)
B3	-	-	2960 (+220)	2748 (+263)	2821 (+225)	424 (+26.9)	42.4%	38	2.82 (±.04)
C1	3,6,9,12	-	13546 (+621)	7414 (+378)	755 (+171)	1562 (+218)	156 %	40	1.66 (±.03)
C2	-	-	3087 (+231)	5270 (+238)	3200 (+191)	403 (+27.9)	40.3%	35	2.36 (±.03)

The effect of immune MLNC only:

On day 16 PI, a mean of 166 worms was recovered from the rats given immune MLNC on day 6, while the control group harboured a mean of 372 worms, a reduction of 45%. The worms collected from the recipient group were normal in length (2.92 mm) and their larval production on days 8, 12 and 15 was similar to the normal control group.

Group B2 which received immune MLNC alone on day 12 PI had a mean of 97 worms on day 16 compared to the control group mean of 424 worms. The worms of this recipient group were also of normal length (2.72 mm), and their larval production was still relatively high on day 15 PI.

It seems that immune MLNC when transferred on day 6 or 12 PI were very effective in causing a reduction in the number of intestinal worms.

The effect of immune serum only:

Following passive transfer of immune serum on days 3, 6, 9 and 12 PI, a mean of 1563 adult S. ratti were recovered from rats of group C1 on day 16 PI. The control group harboured a mean of 403 worms on the same day. The size of the original inoculation was 1000 (\pm 96) L3 and, thus, it seems that there was the possibility of auto-infection in the recipient rats. The worms collected from group C1 on day 16 PI were small with a mean length of 1.66 mm while the control worms were 2.36 mm. The number of larvae/gm in the recipient group showed a maximum number of 13,546 L3/gm on day 8 PI but dropped sharply to 755 L3/gm on day 15 PI when the control group produced a mean of 3200 L3/gm.

Effect of immune MLNC and immune serum:

The two groups which received a combination of both immune MLNC and serum, produced different results depending on the time

Figure 19, Exp. 6.1:

The effect of passive and adoptive transfer of immunity on the percentage of worm recovery on day 16 PI.








Figure 20, Exp. 6.1:

Larval production in three major groups of rats, A, B, and C. \blacktriangle — \blacktriangle A₁ received immune serum and MLNC on day 6 PI; \triangle — \triangle A₂ received MLNC alone on day 6 PI; \blacksquare — \blacksquare B₁ received immune serum and MLNC on day 12 PI; \square — \square B₂ received MLNC only on day 12 PI; \circ — \circ C₁ received immune serum only; \bullet — \bullet Control groups.



of MLNC transfer. Group A1, which received immune serum on days 3, 6, 9 and 12 PI and immune MLNC on day 6 PI, harboured a mean of 681 worms on day 16 PI with a range of 1122 - 238 worms. The control group harboured 372 worms on the same day. The initial inoculum was 1000 (+ 39) L3. This recovery rate also suggested auto-infection. When immune MLNC were given on day 12 PI in conjunction with four injections of immune serum in group B1, the picture was different. A very significant reduction of the worm burden was encountered on day 16 PI (mean 145 worms) while the control group harboured 424 worms on the same day. The mean length of the worms recovered from the recipient group was below normal (1.79 mm) but larval production on days 8, 12 and 15 PI was similar to the control group.

This experiment suggested that immune MLNC alone, whether given early (day 6 PI), or during the middle of infection (day 12 PI), were capable of causing a reduction in the worm burden and accelerated the expulsion of normal looking worms. Immune serum alone appeared to induce auto-infection. Immune MLNC and immune serum caused expulsion only when immune MLNC were introduced on day 12 PI. Auto-infection also occurred following transfer of immune serum and MLNC on day 6 PI.

Experiments 6.2, 6.3, 6.4, 6.5, and 6.6:

These experiments were repeats of the previous experiment and the same type of animals and procedures were used in transferring immune serum and MLNC.

The protocol of these experiments and their results are summarized in Table 18 and are illustrated collectively in Figure 21.

Figure 22 summarizes the results of the experiments on adoptive and passive transfer. In this figure, the normal infection control group is expressed as 100%. Thus any figure appreciably over 100 indicates auto-infection and any figure appreciably below 50 indicates worm expulsion.

Table 18

Summary of five repeats of serum and cell transfer of immunity to rats infected with S. ratti

Exp:	Groups (no. rats)	Days of transfer		Mean (SE) number of adult worms	Mean (SE) length of worms (mm)	% of development	Statistical significance
		Serum	MINC (viability %)				
6.2	A (4)	-	6 (73%)	181 (+32.2)	2.81 (+.04)	18.1%	A vs B P<.001
	B (4)	-	-	297 (+30.5)	2.83 (+.04)	29.7%	
6.3	A (5)	-	6 (78%)	234 (+47.2)	2.79 (+.03)	23.4%	A vs C P<.01
	B (5)	3,6,9,12	6 (78%)	380 (+104)	2.1 (+.06)	38%	B vs C P<.01
	C (5)	-	-	577 (+71.6)	2.68 (+.04)	57.5%	
6.4	A (4)	-	12 (51%)	401 (+25.3)	2.8 (+.03)	40.1%	A vs C N.S.
	B (4)	3,6,9,12	-	1170 (+100)	1.7 (+.06)	117 %	B vs C P<.001
	C (4)	-	-	681 (+45.7)	2.63 (+.05)	68.1%	
6.5	A (5)	-	12 (81%)	63.2 (+9.8)	2.73 (+.05)	6.3%	A vs C P<.001
	B (5)	3,6,9,12	-	174 (+14.6)	1.6 (+.06)	17.4%	B vs C P<.01
	C (4)	-	-	257 (+27)	2.66 (+.03)	25.7%	
6.6	A (5)	3,6,9,12	12 (71%)	191 (+20.5)	2.65 (+.04)	19.1%	A vs C P<.001
	B (5)	3,6,9,12	-	906 (+230)	1.71 (+.06)	90.6%	
	C (4)	-	-	430 (+46.9)	2.8 (+.03)	43%	

Figure 21

Summary of five repeats of passive and adoptive transfer of immunity to rats infected with *S. ratti*.






Figure 22

Pooled results of experiments on adoptive and passive transfer of immunity to S. ratti.

- A : Normal infection control = 100
B : Received immune MLNC on day 6 PI = 45
C : Received immune MLNC on day 12 PI = 43.3
D : Received immune serum on days 3, 6, 9, 12 PI = 221
E : Received immune serum and MLNC on day 6 PI = 123
F : Received immune serum and MLNC on day 12 PI = 39



It was evident from experiments 6.1 - 6.6 that transferred sensitised MLNC, whether on day 6 or 12 PI, significantly decreased the number of worms on day 16 PI. On the other hand, immune serum alone induced a significant increase in the number of adult worms. Rats which received immune MLNC on day 6 PI in conjunction with immune serum usually showed signs of auto-infection. However, results of experiment 6.3 indicated a decrease in the number of adult worms. Immune serum and MLNC given on day 12 PI consistently resulted in a significant degree of worm expulsion.

Experiment 6.7:

This experiment was designed to determine whether immune MLNC, injected intraperitoneally 12 days after challenge, could induce immunity. One group of rats received 4×10^8 viable MLNC intraperitoneally from donor rats infected 25 days previously. One group received 2×10^8 MLNC intravenously and another group 2×10^8 MLNC plus immune serum on days 3, 6, 9 and 12 PI. Another group received neither cells nor serum. The results are presented in Table 19 and Figure 23.

The group which received MLNC IP had similar numbers of adult worms to the normal group (354 vs 455). The group given MLNC IV had fewer worms than the controls (94 vs 455). The group given MLNC and serum had rather fewer worms than the controls (247 vs 455).

These results suggest that injecting sensitised MLNC intraperitoneally did not convey immunity to the recipient group.

Experiment 6.8:

This experiment was designed to explore the activity of immune serum and/or cells when transferred to a naive group of rats before or on the day of infection.

Each rat in group A received 2×10^8 immune MLNC, IV,

Table 19

Exp: 6.7





Comparison between two routes of transferring (MLNC) to recipient rats


(IV) and (IP) and effect of serum and lymphocyte transfer on the course of infection

Group	Days of transfer		Injection route	Number MLNC	Mean (SE) number of worms on day 16 PI	% development
	Serum	MLNC (70% viable)				
A	-	12	IV	2×10^8	94.4 (+ 29)	9.4%
B	-	12	IP	4×10^8	354 (+ 32.9)	35.4%
C	3,6,9,12	12	IV	2×10^8	246 (+ 15.3)	24.6%
D	-	-		-	455 (+ 5.4)	45.5%

A vs D P < .001 : B vs D N.S. : C vs D P < .01 : A vs B P < .001

Figure 23, Exp. 6.7:

Percentage of worm recovery on day 16 PI from four groups of rats.  A received MINC, IV;  B received MINC on day 12 IP;  C received serum and MINC on day 12;  D normal infection control.



(viability 70%) one day before infection and 1.5 ml of immune serum on the day of infection. Group (B) received MLNC alone on day -1, and group (C) received immune serum only on day 0. The fourth group (D) was infected on the same day but was not given any MLNC or serum. All the rats in the four groups were killed on day 8 PI, and the number of adults counted. The results are summarized in Table 20 and Figure 24. The results indicated that the best effect of transferred immune components was obtained when immune MLNC were transferred on day -1 and immune serum on day 0. The number of worms recovered on day 8 PI was the lowest among the other groups (54 ± 9). However, MLNC alone on day -1 were also capable of causing a reduction in the number of worms following transfer. Immune serum in this experiment did not transfer protective immunity to rats in group (C). This experiment demonstrated that immune MLNC can sensitise recipient rats and produce protective immunity against an initial infection. Immune MLNC and immune serum can co-operate to protect the recipient rats effectively.

Discussion:

The results of the previous experiments have suggested that expulsion of S. ratti is due to an immune reaction in rats. It is inevitable that these results should be compared with results, on the mechanisms of immune expulsion, obtained in N. brasiliensis, P. muris and T. spiralis which have been studied extensively. As mentioned before, a multi-step mechanism has been proposed for worm expulsion in these parasitic systems. The results in this chapter have shown that the necessity of worm damage and the effect of transferred immune serum on the course of S. ratti infection (see later) did not conform with results obtained in other systems. New, and sometimes controversial, results have been presented here which require careful interpretation.

It seems that MLNC collected on day 25 PI from initially infected donors contained sufficient committed lymphocytes which transferred the immune response to recipients.

Table 20

Exp: 6.8 Number of adult worms in four groups following
passive and adoptive transfer of immunity

Group	Days of transfer		Mean (SE) number of adult worms	Percentage of worm development
	Serum	MLNC (viability 70%)		
A	0	- 1	54 (+ 9)	5.4%
B	-	- 1	115 (+ 8.83)	11.5%
C	0	-	268 (+ 14.5)	26.8%
D	-	-	304 (+ 10.9)	30.4%


A vs D P < .0005 :

B vs D P < .0005 :

C vs D P < .001

Figure 24, Exp. 6.8:

Percentage of worm development in four groups.
A received immune MLNC on day -1 and immune serum on day 0; B received immune MLNC on day -1 only; C received immune serum on day 0 only; D normal infection control.



The effect of the transfer of immune MLNC only to syngeneic rats on day 6 PI was intriguing. The worm burden of recipient rats was reduced following this transfer and the worms had lengths similar to those of the control group. It should be remembered that only worm length was used as the criterion for damage. Accordingly, worms were considered to have been expelled when they showed no superficial signs of damage. Since it has been shown that worm damage was a prerequisite for the expulsion of N. brasiliensis and T. spiralis (Ogilvie and Jones, 1973; Love et al., 1976; Wakelin and Lloyd, 1976b) it is possible that worm damage is not required in the process of S. ratti expulsion.

It is important to remember that the amount of worm damage required by the host to induce the second step of worm expulsion, has not been defined as yet. It is possible that there is a minimum amount of damage and that this is sufficient to induce an effective attack by the cellular step. The worms which were expelled following the transfer of MLNC on day 6 might have fulfilled this minimum requirement, which was not detectable on a macro-level. Moreover, it is possible that these expelled worms were shorter than those recovered on the day of autopsy.

The other important fact is that although mesenteric lymph nodes (MLNs) contain a high proportion of T. cells (Raff, 1970), a substantial proportion of these lymphocytes are B. cells. Despite the use of glass wool columns to purify MLNC suspensions from the majority of B. cells and macrophages, some B. cells would have been transferred with the MLNC suspension. Cochrane and Dixon (1962) have demonstrated that the transfer of cells initiates the formation of antibodies in recipients. Thus, such transferred humoral response may have cooperated with the recipient's own response and caused a degree of damage sufficient to allow the second step to act. Crandall and Crandall (1972) demonstrated the presence of IgG1 on cells from MLNs of mice infected with T. spiralis. This is interesting since this immunoglobulin is involved in protective immunity to N. brasiliensis (Jones et al., 1970).

A gap of 10 days between cell transfer and autopsy, in groups which received MLNC on day 6, was sufficient for MLNC to mount a successful attack on the intestinal worm population. The

transferred MLNC possibly acted by producing relevant lymphokines causing the proliferation of specific and by-stander lymphocytes, as well as specific antibody-producing cells, which together cooperated to eliminate the majority of worms.

When immune MLNC were transferred on day 12 PI, leaving a period of 4 days before autopsy, expulsion was evident. The worms had normal lengths and no apparent signs of damage were observed. The argument regarding the validity of damage mentioned earlier applies here as well.

It appears that cooperation between the transferred MLNC and the recipients' own immune system was successful in bringing about a decrease in worm burden within 4 days.

The effects of transfer of immune serum, at 4 intervals, on the course of S. ratti infection was unique. The recovery of worms which exceeded in number that of the original inoculum indicated the development of auto-infection. Auto-infection did not occur during the course of normal infection, but did occur under certain states of immune depletion (see Chapter 8). When immune serum was introduced to infected rats on day 3, 6, 9 and 12 PI, auto-infection occurred repeatedly (with the exception of experiment 6.5) and worms recovered were short and less fecund. The short worms were either damaged through the action of antibodies as described in N. brasiliensis (Ogilvie and Jones, 1973) or they were new young worms which arrived in the intestine as a result of auto-infection.

In human strongyloidiasis, recent evidence indicated that auto-infection occurred in patients with depleted cell mediated immunity (CMI) either due to infections which cause such depletion, e.g. Lepromatous Leprosy and Tuberculosis, or following immunosuppressive therapy (see Chapter 9) (Purtillo *et al.*, 1974). If this was true in rats, then transferred antibodies were causing some state of specific or generalized depletion which led to auto-infection.

It can be suggested that a proportion of these transferred antibodies was not protective and acted as blocking (enhancing) antibodies, possibly similar to those demonstrated in certain tumour-bearing individuals (Hellström and Hellström, 1969, 1970; Hellström

et al., 1969). Since these antibodies, in tumour immunology, may be antibody/antigen complexes (Sjögren et al., 1971), they may operate a feed-back mechanism which inhibits the production and proliferation of active specific lymphocytes.

The mechanisms by which antibodies render the cellular arm of the immune response ineffective are still undetermined. Such depletion must be effective on both the T. cells in the gut and associated lymph nodes, as well as the circulating population, so as to allow the infective stages to re-migrate through the various tissues during auto-infection, without any resistance.

The effects of these antibodies on the course of S. ratti infection may be attributed to mechanisms similar to those suggested by Rickard and Outteridge (1974) for the depletion of the OMI response in hyperinfection with Taenia pisiformis in rabbits. Antibodies produced by the infected rabbits, coated the surface of the cysts of this parasite and inhibited the action of OMI on them (Rickard, 1974).

The possibility that a factor, produced by the interaction of worms and antibodies, kills lymphocytes, similar to that suggested in T. spiralis infection (Faubert and Tanner, 1975; Faubert, 1976), also exists.

The blocking effects of these antibodies might be a contributory factor in the tenacity of S. stercoralis in some individuals, which have been attributed to repeated auto-infection. It would be very interesting to study the level of antibody production in those individuals as well as the state of their specific and general OMI response.

Lawler (1940) passively transferred immunity against S. ratti to a group of rats by injecting them with immune plasma from multiply infected rats. The degree of protection which he obtained was only partial and was similar in rats given immune plasma before or after the introduction of the infection. This is, obviously, in sharp contrast to the results obtained in my experiments.

It was not possible to assess fully the role of antibodies in inducing worm damage in this system due to the consistent development of

auto-infection. However, when serum did not cause auto-infection, worms were short.

In one experiment (6.5), immune serum alone did not cause auto-infection, but instead produced a significant drop in the number of adult worms. This can be attributed to the different pool of serum used (Ogilvie and Jones, 1968).

Rats which received immune serum and MINC on day 6 PI showed a high rate of worm development, with a few rats showing auto-infection. On the other hand, those which received serum and MINC on day 12 PI showed the best expulsion rate. These two results are difficult to interpret. It is possible that antibodies affected the worms and the MINC which followed on day 12 acted immediately and caused expulsion. It seems that this would be the ideal approach in bringing about worm elimination by these transferred components. Why this serum, which came from the same pool and was given at exactly the same intervals, did not appear to have caused auto-infection on this occasion, is not clear. Antibodies may have inhibited the host's own cells (and probably the transferred MINC on day 6) from recognising the antigens (masking) or possibly produced other factors, as discussed previously.

This aspect of blocking antibodies and their causal effects on auto-infection in this system needs to be pursued. No such immunological enhancement has been demonstrated in other helminthic infection, but preliminary observations suggest that in some parasitic infections, antibodies are produced in abundance while the CMI response is ineffective (Ogilvie and Wilson, 1976).

MINC, when given through the IP route did not convey immunity compared to the IV route. Undoubtedly, a few of these cells, when put in IP, reach the MINS, but the majority are absorbed and may reach other organs which are not involved in the reaction against the intestinal worms.

MINC given alone one day before infection gave a substantial degree of protection. The action of these lymphoid cells can be aimed against either the migratory larvae, the L4 stage, the adult worms or all three. Love (1975) suggested a quantitative difference between the immunological requirements for the expulsion of larvae and adult N. brasiliensis. The transferred MINC may have lingered

about until L3 of S. ratti moulted to L4 or the subsequent stage, before mounting an attack. On the other hand, it is possible that the cells may have infiltrated to the lungs and affected the migration of the invading larvae by developing lymphoid centres similar to those observed during second infection with S. ratti (see Chapter 3). There was no evidence of larval detention in the lungs in rats infected with N. brasiliensis and given immune serum and MNC (Love et al., 1974). Love (1975) suggested that fourth-stage larvae of N. brasiliensis were expelled from the intestine by the action of MNC before moulting to the fifth-stage.

MNC transferred one day before infection and serum on the day of infection also gave good protection. Serum alone gave no protection, but produced no auto-infection either. Whether immune serum co-operated in the process of reducing the worm burden on day 8 PI, is still unclear.

In conclusion, it is difficult to propose a multi-step mechanism in the expulsion of S. ratti from rats as described in other systems (Ogilvie and Love, 1974; Wakelin, 1975; Love et al., 1976; Wakelin and Lloyd, 1976b). However, it is clear that immune MNC are important in inducing worm expulsion, whether given on day 6 or 12 PI. Co-operation between immune serum and MNC was evident only when MNC were given on day 12 and not 6. Immune serum presented a new parameter as far as the relationship between antibodies and worms is concerned.

It is important that studies should be conducted on transfer of immunity to infected, irradiated rats, in order to present conclusive evidence on the role of lymphoid cells in the process of worm expulsion. However, this might prove to be difficult in this system since irradiation causes immune depletion and can subsequently lead to the development of auto-infection.

CHAPTER 7Studies on the role of other factors in the process
of worm expulsion in *S. ratti* infections

In many nematode infections of rodents, a variety of additional factors have been implicated in the humoral and cellular reactions against the living worms, among them mast cells, basophils and eosinophils.

Although mast cells and basophils have been studied extensively, there is still uncertainty as to their role in the expulsion of helminthic infections (Murray et al., 1971a, b, c; Wells, 1962; Whur, 1966, 1967; Rothwell et al., 1974b). The trigger for the production of histamine and other substances, from mast cells and basophils, is the reaction between IgE and an antigen. This immunoglobulin increased during helminthic infections, particularly prior to the expulsion of the intestinal worms (Ogilvie, 1967; Ogilvie and Jones, 1969, 1971). Simultaneously with the rise in the amount of IgE, there was an increase in the infiltration of mast cells and/or basophils near the site of infection (Murray et al., 1971a, b, c). Analysing the role of anaphylaxis in worm expulsion in *N. brasiliensis*, Urquhart et al. (1965) and Barth et al. (1966) proposed what they called the "leak lesion" hypothesis. It suggested that the production of amines by the mast cells of the gut attracted anti-worm antibodies towards the worms where they caused damage. Murray et al. (1971c) demonstrated that worm expulsion did not occur when infected rats were treated with the anti-amine drug, promethazine. However, mounting evidence against this hypothesis has been presented. Production of IgE occurred in rats which were not resistant to infection, and in rats immunized against worm extracts, although they showed no immunity to re-infection (Ogilvie, 1967). Also, many pools of antisera had high titres of IgE but were not protective (Ogilvie and Jones, 1967, 1968). Changes induced by the immune response in adult *N. brasiliensis* whether on metabolic or

structural levels occurred before the infiltration of mast cells into the villi (Henney et al., 1971; Keller, 1971; Murray et al., 1971c; Whur, 1966). Confirmation of this came from work on neonatal and lactating rats in which worm damage was brought about by a humoral response independent of mast cells (Ogilvie and Hockley, 1968; Keller, 1971; Kelly and Ogilvie, 1972). Damaged worms were expelled swiftly in irradiated and T. cell reconstituted rats long before mast cells appeared (Keller and Keist, 1972). The anti-amine drugs used by some workers (Campbell et al., 1963; Murray et al., 1971; Rothwell et al., 1971) to prolong infection with T. spiralis and N. brasiliensis, were found to be immunosuppressive (Henney et al., 1972) which might account for the delay in expulsion (Kelly and Dineen, 1972; Gudson et al., 1972; Keller and Ogilvie, 1972). The final argument against the "leak lesion" hypothesis was based on histopathological evidence, where expulsion occurred prior to a significant increase in mast cells in the intestinal tissue (Wells, 1962; Whur, 1966; Keller, 1971; Kelly and Ogilvie, 1972).

As far as N. brasiliensis is concerned, it appears that amines are secondary in their role in worm damage and expulsion. However, a definite role for amines in the process of worm expulsion in T. colubriformis in guinea-pigs has been firmly established (Rothwell et al., 1971; Rothwell et al., 1974a, b; Jones et al., 1974).

Preliminary evidence implicated tissue eosinophils in the second cell-mediated step of worm expulsion. Tissue eosinophils have been linked with the time of expulsion (Wells, 1962; Hogarth-Scott and Bingley, 1971; Kelly and Ogilvie, 1972) and during the migratory stages of helminthic infections (Opie, 1904; Cornwell, 1962; Zaiman et al., 1963).

The current knowledge of the role of eosinophils in immune response is still limited. It is well established that the specific action of eosinophils is thymus-dependent (Walls et al., 1971, 1973). They act through receptors for immune adherence and possibly the Fc portion of IgG (Kay, 1974). Eosinophils can be attracted to the site of infection by a cellular response, through the production, by sensitized lymphocytes, of an "eosinophil chemotactic factor,

E.C.F.", which attracts these granulocytes towards the site of antigenic stimulation (Cohen and Ward, 1971). There is some evidence for the existence of a relationship between the distribution of eosinophils and mast cells. Histamine released by mast cells is chemotactic for eosinophils (Riley, 1959; Archer, 1963, 1965; Fernex, 1968). Also, antibody-antigen complexes can be chemotactic for eosinophils which phagocytose these particles on contact (Archer and Bosworth, 1961; Sabesin, 1963; Litt, 1964). Eosinophils played a definite role in mediating an antibody-induced damage to schistosomulae (Butterworth *et al.*, 1974, 1975). Whether the cells cooperating during the finalization of worm expulsion in these nematode infections, are eosinophils or macrophages, is still under investigation. Macrophages are not directly involved in this last step (Keller and Keist, 1972), but have an essential role in the early induction of the immune response.

Although complement is not important during the process of worm elimination in *N. brasiliensis* (Jones and Ogilvie, 1971), its direct role in the final stages of expulsion should not be dismissed.

A new mediator of inflammatory reaction has also been implicated in the final stages of the immunological expulsion of *N. brasiliensis*. Kelly *et al.* (1974) suggested that pharmacologically active products of different tissue cells might have a role in the final step of worm expulsion. They suggested that lymphokines and prostaglandins may play a significant role as non-specific mediators, since they are widely spread in different tissues of the body, particularly the intestine. They presented evidence that prostaglandins from rat semen, which had a smooth muscle stimulating activity, caused expulsion of *N. brasiliensis* from rats. Prostaglandins are unsaturated hydroxy-fatty acids, found in both sexes and are present in most body tissues (Smith *et al.*, 1974). The administration of synthetic prostaglandins, particularly of the class E, was very effective in causing rapid worm expulsion (Dineen *et al.*, 1974). Potent inhibitors of prostaglandins, such as aspirin and dextropropoxyphene hydrochloride, prevented the expulsion of *N. brasiliensis* when administered daily during the course of infection (Kelly *et al.*, 1974).

The experiments in this chapter are preliminary investigations

into the role of eosinophils, mast cells and their derivatives and prostaglandins in the process of S. ratti expulsion from rats.

Method for counting tissue eosinophils in intestine:

Intestinal villi of rats, which are tongue-shaped in longitudinal sections, are separated, at their base, by the gland crypt (Reynolds et al., 1967; Nordstrom et al., 1968). The sections of intestines were stained with Carbol Chromotrope. For counting purposes, the villi were divided into "villus-crypt" units (VCU) and the number of eosinophils lying within each villus, between two gland crypts and its lamina propria, were counted (Jarrett et al., 1968; Kelly and Ogilvie, 1972). Ten VCU were used for each standard count and the mean of six standard counts was determined. Tissue eosinophil counts were expressed as the number of cells/10 VCU.

Mesenteric mast cells (see Chapter 2) were counted in 50 randomly selected microscopic fields (x 16) and the percentage of degranulated mast cells was determined.

Total white blood cell (W.B.C.) count:

The freshly collected blood was sucked up to the 0.5 mark on a white cell haemocytometer pipette. This was diluted 1 in 20 with a fluid consisting of the following:

- 1 ml of glacial acetic acid
- 99 ml of distilled water
- 4 drops of toluidine blue

Cell counts were performed in a "bright-line" improved Neubauer haemocytometer.

Blood eosinophil count:

Fresh blood was diluted 1:10 in cold Discombe's fluid.

This consisted of the following:

5 volumes of 1% aqueous Eosin Y

5 volumes of Acetone

90 volumes of distilled water

In both previous counts, the total number of cells in all 8 corner squares was counted and the number of eosinophils or W.B.C. calculated as follows:

Total number of cells counted x 25 = Number of cells per
cu.mm. of blood.

Experiment 7.1:

In this experiment, levels of eosinophils, in the peripheral blood and intestinal lamina propria, were compared and the percentage of mast cell degranulation, during the course of initial and challenge infections with S. ratti, was determined.

No success was achieved with staining intestinal mast cells using either Astra-Blue or Acid Toluidine Blue with tissues fixed with Carnoy's or Formol-saline. However, mesenteric mast cells were easily stainable with Acid Toluidine Blue and were, thus, used to assess mast cell activity during the course of infection.

Sixty four female Wistar rats were divided into 2 groups. 28 rats were infected with 1000 (\pm 43) larvae each and 36 rats were left uninfected. Four rats from both groups were killed on days 8, 14, 18, 23 and 28 PI and the percentage of degranulating mesenteric mast cells and the number of tissue eosinophils counted. The intestine was divided into anterior and posterior halves and the number of adult worms determined. The peripheral blood picture was examined every three days from day 3.

On day 30 PI, the 8 members of the first group were each reinfected with 1000 (\pm 56) larvae, simultaneously with 8 rats from the control group. This left 8 rats to act as uninfected controls. 4 rats were killed from each of these groups on day 5 and 8 post-

Table 21

Exp: 7.1

Changes in the tissue eosinophil level in the intestinal mucosa
and the percentage of degranulation of mesenteric mast cells
during and initial and a challenge infection

Days PI	Mean (SE) total number of worms		% of worm development	Mean (SE) total number of eosinophils in 10 V.C.U./rat		% degranulation of mesenteric mast cells	
				infected	non-infected	infected	non-infected
8	298	(+21.6)	29.8%	108 (+ 8.04)	94.8 (+ 8.49)	31.6	21.7
14	325	(+36.9)	32.5%	139 (+ 18.2)	125 (+ 6.15)	54.5	19.8
18	349	(+13.6)	34.9%	134 (+ 8.72)	88.5 (+ 1.80)	63.3	27.9
23	187	(+12.2)	18.7%	251 (+ 16.4)	71.8 (+ 6.07)	75.5	26.6
28	46.5	(+16.6)	4.6%	194 (+ 6.63)	72.8 (+ 5.95)	52.8	23.1
Post-challenge							
5	38.5	(+6.17)	3.8%	119 (+ 14.2)	72 (+ 2.89)	62.5	21.9
8	18.5	(+ 4.57)	1.8%	248 (+ 13.4)	73 (+ 3.43)	70	22.6

Table 21

Exp: 7.1


Changes in the tissue eosinophil level in the intestinal mucosa
and the percentage of degranulation of mesenteric mast cells
during and initial and a challenge infection

Days PI	Mean (SE) total number of worms		% of worm development	Mean (SE) total number of eosinophils in 10 V.C.U./rat		% degranulation of mesenteric mast cells	
				infected	non-infected	infected	non-infected
8	298	(+21.6)	29.8%	108 (+ 8.04)	94.8 (+ 8.49)	31.6	21.7
14	325	(+36.9)	32.5%	139 (+ 18.2)	125 (+ 6.15)	54.5	19.8
18	349	(+13.6)	34.9%	134 (+ 8.72)	88.5 (+ 1.80)	63.3	27.9
23	187	(+12.2)	18.7%	251 (+ 16.4)	71.8 (+ 6.07)	75.5	26.6
28	46.5	(+16.6)	4.6%	194 (+ 6.63)	72.8 (+ 5.95)	52.8	23.1
Post-challenge							
5	38.5	(+6.17)	3.8%	119 (+ 14.2)	72 (+ 2.89)	62.5	21.9
8	18.5	(+ 4.57)	1.8%	248 (+ 13.4)	73 (+ 3.43)	70	22.6

Figure 25, Exp. 7.1:

Changes in the levels of tissue eosinophils degranulating mesenteric mast cells and number of adult worms in two groups of rats.

- percentage of adult development in rats
- percentage of degranulating mesenteric mast cells in infected group
- percentage of degranulating mesenteric mast cells in non-infected group
- ▲—▲ number of tissue eosinophils as mean in 10 VCU in infected group
- △—△ number of tissue eosinophils as mean in 10 VCU in non-infected group.



challenge and the levels of peripheral and tissue eosinophils, mast cell degranulation and adult worms determined.

The results are summarized in Table 21 and Figure 25. There was an increase in the level of tissue eosinophils from 134, on day 18 PI, to 251 cells/10 VCU on day 23 PI, which coincided with the period immediately before worm expulsion. The infiltration of eosinophils was extensive in both the mucosa and the Peyer's patches (Plates 48 and 49). Tissue eosinophils showed another sharp increase to 248 cell/10 VCU on day 8 post-challenge (Plate 50) while the uninfected control (Plate 51) maintained an average of 85 cells/10 VCU throughout the experiment.

The percentage of degranulating mesenteric mast cells (Plates 52 and 53) increased from day 14 PI and reached a maximum on day 23 PI, after which it dropped slightly. A similar increase was observed on day 8 post-challenge.

In the initially infected group, there was a sharp rise in the total W.B.C. count (Figure 26) to 20,000 cells/mm³ between days 21-28 PI. The levels of blood eosinophils demonstrated peaks on days 6-12 and 28-34 PI. Following challenge, the total W.B.C. count rose sharply to 18,000 cells/mm³, but this was not paralleled by any rise in the level of blood eosinophils.

Experiment 7.2:

In this experiment, the role of a histamine derivative, on the course of S. ratti infection, was investigated.


Ten rats were each infected with 1000 (\pm 93) larvae. Histamine diphosphate (Koch-Light Laboratories Ltd.) was administered by intubation, twice daily, to 5-rats on days 12, 13, 14 and 15 at 100 mg/kg of body weight, while the other 5 rats were given equal volumes of saline. All the rats were killed on day 16 PI and the number of adult worms, in the anterior and posterior intestine, and larvae/gm of faeces determined.

The results are presented in Table 22 and Figure 27. Rats which received histamine harboured a mean of 317 worms on day 16, compared to 575 worms in the untreated group. This difference was

Figure 26, Exp. 7.1:

Blood eosinophilia and total white blood cell count in two groups of rats.

- eosinophil count in infected group.
- eosinophil count in normal rats.
- △—△ total W.B.C. count in infected group.
- ▲—▲ total W.B.C. count in normal rats.






Plate 48 : Section through villus/crypt units showing infiltration of eosinophils prior to S. ratti expulsion (Lendrum's, x 250)




Plate 49 : Infiltration of eosinophils within a Peyer's patch at the end of an initial S. ratti infection (Lendrum's, x 250)




Plate 48 : Section through villus/crypt units showing infiltration of eosinophils prior to S. ratti expulsion (Lendrum's, x 250)




Plate 49 : Infiltration of eosinophils within a Peyer's patch at the end of an initial S. ratti infection (Lendrum's, x 250)




Plate 48 : Section through villus/crypt units showing infiltration of eosinophils prior to S. ratti exulsion (Lendrum's, x 250)




Plate 49 : Infiltration of eosinophils within a Peyer's patch at the end of an initial S. ratti infection (Lendrum's, x 250)




Plate 50 : Tissue eosinophils in villi of rats challenged with S. ratti (Lendrum's, x 250)




Plate 51 : Eosinophils in a normal uninfected rat (Lendrum's, x 250)




Plate 50 : Tissue eosinophils in villi of rats challenged with S. ratti (Lendrum's, x 250)




Plate 51 : Eosinophils in a normal uninfected rat (Lendrum's, x 250)




Plate 52 : Intact mesenteric mast cells (Acid Toluidine Blue,
x 400)



Plate 53 : Degranulated mesenteric mast cells (Acid Toluidine
Blue, x 400)




Plate 52 : Intact mesenteric mast cells (Acid Toluidine Blue,
x 400)




Plate 53 : Degranulated mesenteric mast cells (Acid Toluidine
Blue, x 400)

Table 22 Exp : 7.2: Result of autopsy on day 16PI of two groups: one treated with histamine and the other a normal infection.


	Infected treated	Infected non-treated
Number of rats	5	4
Total number of worms	317 (+ 69)	575 (+ 33)
Mean (SE) number worms in anterior intestine	287 (+ 61)	535 (+ 34)
Mean (SE) number worms in posterior intestine	29.6 (+ 61)	40 (+ 4.3)
Larvae / gm of faeces	1077 (+ 64.6)	15,300 (+ 437)
Fecundity larvae/worm	4.07 (+ 0.24)	26.6 (+ 1.4)
Mean overall length of adult worms	1.81 (+ 0.03)	2.77 (+ 0.02)

Infected treated vs Infected non-treated P < .01

Figure 27, Exp. 7.2:

Effect of histamine treatment on the recovery of adult worms on day 15 PI and their fecundity

■ infected, histamine treated
□ normal infection



statistically significant. There was no difference in the distribution of adult worms between the two groups. Worms recovered from histamine treated rats, were stunted and showed diminished fecundity (4.07 vs 26.6 larvae/worm).

This experiment suggests that histamine may play a part in the elimination of S. ratti infection from rats, but has no effect on the distribution of adult worms in the intestine.

Experiment 7.3:

In this experiment, the effect of the prostaglandin inhibitor, aspirin, on the expulsion of S. ratti, was noted.

Sixteen rats were infected with 1000 (\pm 69) larvae each. Aspirin powder B.P. (Acetylsalicylic acid, John Bell and Croydon) was administered to 8 rats twice daily at 3 mg/kg, per os, for the entire period of experiment. 8 rats were treated with equal volumes of saline alone, on the same occasions. Faecal cultures were made on days 14, 22 and 24 PI. Four rats from each group were autopsied on days 28 and 30 PI and the number of adult worms, in the anterior and posterior intestine, counted. The results are presented in Table 23 and Figures 28 and 29.

The aspirin-treated group harboured more adult worms, on day 28 and 30 PI, than the saline-treated group. This indicated that a relative, but significant, delay in worm expulsion occurred in rats treated with aspirin. Aspirin also had no effect on the distribution of worms in the intestine. There were more larvae per gram of faeces produced by the rats treated with aspirin, on day 24 PI, than the control group.

This experiment suggests a possible role for prostaglandins in the final step of S. ratti expulsion, as indicated by the effect of a potent inhibitor.

Table 23, Exp. 7.3:

Result of autopsies on two different days of two groups:

A: infected and treated twice daily with Aspirin

B: normal, non-treated group

Day	Group	Mean (SE) total number of worms	Mean (SE) number adult worms recovered	
			anterior intestine	posterior intestine
28 PI	A	114 (<u>±</u> 52.2)	4 (<u>±</u> 1.08)	110 (<u>±</u> 52.1)
	B	24.7 (<u>±</u> 3.49)	2 (<u>±</u> .40)	22.5 (<u>±</u> 3.17)
30 PI	A	38.7 (<u>±</u> 7.25)	1 (<u>±</u> .70)	37.7 (<u>±</u> 6.62)
	B	0.25 (<u>±</u> 0.25)	0	0.25 (<u>±</u> 0.25)

Figure 28, Exp. 7.3:

Effect of aspirin on the number of adult worms
■ anterior intestine; □ posterior intestine.
A = treated with aspirin, B = treated with saline.





Figure 29, Exp. 7.3:

Effect of aspirin on the level of larval production.
○—○ treated with aspirin; ●—● normal control.



Discussion:

The results of the previous experiments suggest that factors, other than antibodies and cells, might be involved in the expulsion of S. ratti.

The rise in the level of eosinophils in the lamina propria of infected intestines, coincided with the onset of worm expulsion. This also occurred in rats infected with N. brasiliensis (Wells, 1962; Hogarth-Scott and Bingley, 1971; Kelly and Ogilvie, 1972).

Blood eosinophils, on the other hand, increased in two separate peaks before and after the expulsion. This result was in agreement with those obtained by Bailenger and Caroenac (1974) in rats infected with S. ratti. The decline in the number of blood eosinophils can be correlated with the rise in tissue eosinophils.

The activities and properties of eosinophils and their involvement in immune reactions are not fully elucidated (Review by Archer, 1970). Eosinophils are associated with cellular immune reactions and, thus, can be expected to appear in the intestinal tissue during the second step of worm expulsion.

Eosinophils are attracted to the site of inflammation by a variety of factors (see introduction, ^{to} this chapter). The duration of life of blood eosinophils is brief and they can leave the circulation very rapidly (Archer, 1970) and it is generally known that there are 4-5,000 tissue eosinophils to each circulating one (Delanie and Garraty, 1975).

The presence of eosinophils near the site of infection, particularly close to the time of expulsion, might indicate that the cooperation between antibodies and cells can be substantiated by the activities of eosinophils. Drastic alteration of the intestinal environment, through the production of pharmacologically active substances by eosinophils or other myeloid cells (e.g. mast cells) can lead to the evacuation of worms. However, the chronological order of this reaction is still not determined.

Direct reaction between eosinophils and worms in vivo has not been demonstrated. However, sensitised eosinophils to N. brasiliensis or T. spiralis readily adhere to these worms in vitro, particularly to their anterior ends (McKenzie, 1976,

personal communication). The significance of this reaction in vivo is yet to be evaluated.

The rise in the W.B.C. count during the period of worm expulsion was interesting, but no definite interpretation can be offered except for the possible involvement of other types of leucocytes in this response. Normal rats have a high percentage of lymphocytes in their peripheral blood (about 75-80%) (Cameron and Watson, 1949; also see Table 26 in Chapter 8). The significance of this, as far as worm expulsion is concerned, is still unclear.

No explanation can be offered regarding the lack of success in staining the intestinal mast cells, which was unfortunate, since this might have shed direct light on their role during worm expulsion. The level of degranulation of the perivascular mesenteric mast cells might be of relevance to the activities of these cells during the course of infection. These cells are close to the site of infection and are supplied and drained by the same vascular network as the intestine. Mesenteric mast cells showed an increased level of degranulation at the onset of worm expulsion and following a challenge infection. This degranulation might be due to the sensitization of these cells through the circulation of antibodies and worm antigens within the mesenteries and the production of mast cell degranulating factor (Uvnas and Wold, 1967) by the worms.

The administration of histamine, intragastrically, to rats infected with S. ratti, suggested that this amine might be involved in the process of worm elimination. The worms, which were recovered from histamine-treated rats, developed signs of early damage. Whether histamine attracted homocytotropic antibodies (Hogarth-Scott and Bingley, 1971), which in turn caused morphological changes and diminished fecundity, is still doubtful.

The role played by histamine, in worm expulsion, in vivo, has been shown in guinea-pigs infected with T. colubriformis. Infusion of histamine and 5-hydroxytryptamine directly in the intestine caused rapid expulsion of these worms (Rothwell et al., 1974a). Injection of reserpine, an antagonist of 5-HT, caused a delay in the expulsion of T. colubriformis (Rothwell et al., 1974b). Bailanger and Faraggi (1975b), using reserpine, prevented the expulsion of S. ratti from rats and maintained the infection at a high level, but they attributed

this delay to a state of hypercorticoasteronaemia (Ballenger and Faraggi, 1975a).

When infected rats were treated with the analgesic drug, aspirin, the expulsion of S. ratti, which usually occurs on day 25 PI, was delayed for 5 days. The faecal larval count was higher in the treated group than in the controls.

Aspirin, which is a potent inhibitor of prostaglandins, both in vivo and in vitro (Vane, 1971; Aiken, 1972; Smith and Temple, 1973), caused a delay in the expulsion of N. brasiliensis and prevented the drop in the egg production (Dineen et al., 1974).

Results obtained by the use of antagonists and inhibitors to determine the role of pharmacologically active substances, have been treated with reservations (Ogilvie and Jones, 1973). Some amine antagonists, which caused delay in worm expulsion, had anti-lymphocytic activities (Henney et al., 1972; Kelly and Dineen, 1972b). However, Dineen et al. (1974) showed that administration of aspirin to rats infected with N. brasiliensis had no suppressive effects on their lymphocytic activity.

Recent studies on the role of prostaglandins in the expulsion of N. brasiliensis suggested that they were capable of causing rapid expulsion associated with a reduction in the egg/gm counts (Dineen et al., 1974; Kelly et al., 1974).

Prostaglandins interfere with both humoral and cellular reactions (Piper and Vane, 1969; Søndergaard and Greaves, 1970). In the intestine, where they are found in the tissue and fluids, they affect peristalsis (Bernet, 1972). It was suggested that prostaglandins also contribute towards environmental changes in the gut which makes it difficult for the worms to survive (Dineen et al., 1974).

The trigger for the increase in the level of prostaglandins in the intestine, during expulsion, is still not known. However, it has been suggested, recently, that eosinophils might be involved in the production of these substances (Kay, 1976). If this was ascertained in nematode infections, the interaction between these two factors, during the period of expulsion, could be better understood.

The clinical significance of aspirin treatment to individuals infected with S. stercoralis, and possibly other nematode infections, should be investigated.

The present study presented a preliminary evidence that tissue eosinophils, histamine and prostaglandins might play some role in the process of S. ratti expulsion from rats.

CHAPTER 8Studies on the effect of Betamethasone on the course
of S. ratti infections in rats

Corticosteroids have a remarkable, and as yet ill-defined, capacity to immunosuppress and act as anti-inflammatory agents (Allison, 1965). Their inhibitory action on the nonspecific inflammatory response in a wide range of diseases is often beneficial. They are also useful in assisting with the control of allografts. However, they can be very harmful due to their non-specific action on the adoptive immunological mechanisms which are necessary for the natural defence against viral, bacterial and parasitic infections and which lead to opportunistic infections.

Corticosteroids have a variety of effects on delayed hypersensitivity (DH) (Medawar, 1969). These include their action on the number of circulating lymphocytes and monocytes (Boggs *et al.*, 1964; Thompson and van Furth, 1970), their effect on vascular activity, which causes the inhibition of diapedesis of the different cellular components of the inflammatory reaction (Ashton and Cook, 1952; Ebert, 1952), their inhibitory action on lymphocytopoiesis through their involutionary and cytotoxic effect (Gyllenstein, 1962; Humphrey and White, 1970) and their inhibitory action on the activities of macrophages (Gabrielson and Good, 1966), mast cells (Cavallero and Braccini, 1950, 1951) and eosinophils (Archer, 1956). Casey and McCall (1971) working with BCG, demonstrated the suppressive effects of corticosteroids on DH, both *in vivo* and *in vitro*. They concluded that their effects were on the afferent (adaptation) stage of DH rather than the efferent (expression) stage. Corticosteroids also affect antibody response during the presence of antigen (Darrach, 1959; Berglund, 1962).

Corticosteroids have been widely used in a variety of studies on resistance to helminthic infections, e.g. to study the effects of immunosuppression on initial and subsequent infections in those host-parasite relationships characterized by a strong immune

reaction against the parasite (Stoner and Godwin, 1953, 1954; Weinstein, 1953, 1955; Coker, 1954, 1955, 1956a, b; Markel and Lewis, 1957; Markel, 1958; Campbell and Collette, 1962; Campbell, 1963; Ogilvie, 1965; Tewari, 1970; Wakelin, 1970; Hopkins *et al.*, 1972; Moss, 1972). Also to introduce parasites into naturally resistant hosts (Roman, 1956; Bezubik, 1960; Cross, 1960, 1963; Harley and Gallicchio, 1970) to inhibit the action of the immune response of hosts immunized with antigens from worms (Avlavidou, 1970) and to induce a state of immunological tolerance (Wakelin and Selby, 1974; Behnke, 1975). Corticosteroids depress the immune response to parasites, allowing them to persist in the host beyond the normal period.

The persistence of adult worms in animals treated with cortisone has been reported in many different nematode (Weinstein, 1953, 1955; Ogilvie, 1965; Stoner and Goodwin, 1953, 1954; Coker, 1955, 1956; Markell and Lewis, 1957; Markell, 1958; Cross, 1960; Campbell and Collette, 1962; Campbell, 1963, 1968; Wakelin, 1970, 1974; Avlavidou, 1970), trematode (Tewari, 1970) and cestode infections (Hopkins *et al.*, 1972; Moss, 1972), and this was usually accompanied by a rise in the egg or larval production.

Corticosteroid treatment suppressed acquired immunity following an initial infection with *N. brasiliensis* and a second infection was successfully established (Ogilvie, 1965). Mice treated with cortisone acetate during a primary infection with *T. muris* did not expel the developing larvae within the usual 3 weeks (Campbell, 1963; Wakelin, 1967) and the immune response against a challenge infection was suppressed (Wakelin, 1970). Prolonged cortisone treatment of rats infected with *T. spiralis* suppressed the acquisition of immunity to a primary infection and rendered immunised animals non-immune to a challenge infection (Coker, 1956a; Markell and Lewis, 1957).

Wakelin and Selby (1974) presented the only available purpose-designed experiments to induce a state of immunological tolerance in mice to the nematode worm *T. muris*. Mice which were given cortisone acetate during the second week of *T. muris* infection were rendered tolerant for a period of up to 7 weeks. They concluded that this was not due to small doses of the antigen but rather to

reaction against the parasite (Stoner and Godwin, 1953, 1954; Weinstein, 1953, 1955; Coker, 1954, 1955, 1956a, b; Markel and Lewis, 1957; Markel, 1958; Campbell and Collette, 1962; Campbell, 1963; Ogilvie, 1965^b; Tewari, 1970; Wakelin, 1970; Hopkins et al., 1972; Moss, 1972). Also to introduce parasites into naturally resistant hosts (Roman, 1956; Bezubik, 1960; Cross, 1960, 1963; Harley and Gallicchio, 1970) to inhibit the action of the immune response of hosts immunized with antigens from worms (Avlaidou, 1970) and to induce a state of immunological tolerance (Wakelin and Selby, 1974; Behnke, 1975). Corticosteroids depress the immune response to parasites, allowing them to persist in the host beyond the normal period.

The persistence of adult worms in animals treated with cortisone has been reported in many different nematode (Weinstein, 1953, 1955; Ogilvie, 1965^b; Stoner and Goodwin, 1953, 1954; Coker, 1955, 1956; Markell and Lewis, 1957; Markell, 1958; Cross, 1960; Campbell and Collette, 1962; Campbell, 1963, 1968; Wakelin, 1970, 1974; Avlaidou, 1970), trematode (Tewari, 1970) and cestode infections (Hopkins et al., 1972; Moss, 1972), and this was usually accompanied by a rise in the egg or larval production.

Corticosteroid treatment suppressed acquired immunity following an initial infection with N. brasiliensis and a second infection was successfully established (Ogilvie, 1965^b). Mice treated with cortisone acetate during a primary infection with T. muris did not expel the developing larvae within the usual 3 weeks (Campbell, 1963; Wakelin, 1967) and the immune response against a challenge infection was suppressed (Wakelin, 1970). Prolonged cortisone treatment of rats infected with T. spiralis suppressed the acquisition of immunity to a primary infection and rendered immunised animals non-immune to a challenge infection (Coker, 1956a; Markell and Lewis, 1957).

Wakelin and Selby (1974) presented the only available purpose-designed experiments to induce a state of immunological tolerance in mice to the nematode worm T. muris. Mice which were given cortisone acetate during the second week of T. muris infection were rendered tolerant for a period of up to 7 weeks. They concluded that this was not due to small doses of the antigen but rather to

the immunosuppressive action of the drug during strategic points of infection when the immune system is about to recognise the antigenic materials present. This treatment stopped the usual process of worm elimination even when the drug was only used 4 times during the course of infection. However, Olivier (1962) working with Taenia taeniformis in mice obtained a long-lasting tolerance to this parasite following a few cortisone injections.

Human strongyloidiasis has always been thought of as innocuous, but incidences of overwhelming infections have been reported (Brown and Perna, 1958; Wilson and Thompson, 1964; Toh and Chow, 1969; Rivera et al., 1970; Behar, 1970, 1971; Huggins, 1971a, b; Adam et al., 1973). These infections produced enteritis (De Paolo et al., 1962; Louisy and Barton, 1971; Berkmen and Rabinowitz, 1972), necrotic jejunitis (Craven et al., 1971), peritonitis (Olurin, 1970), ulcerative colitis (de Campos, 1964), paralytic ileus (Nolasco and Africa, 1936; Frengley and Trewby, 1973) and occasional pulmonary (Poltra, 1972) and skin involvements (Thomas et al., 1971). Many reports have shown that corticosteroid therapy had grave repercussions on people infected with S. stercoralis. The most common cases of overwhelming human infection occur when patients with a non-patent or a very mild S. stercoralis infection are treated with corticosteroids. They develop severe and often fatal infection (Rogers and Nelson, 1966; Cruz et al., 1966; Civantos and Robinson, 1969; Yim et al., 1970; Fagundes et al., 1971; Cookson et al., 1972; Bitoun et al., 1972; Fowles, 1973; Neefe et al., 1973). Willis and Nwokolo (1966) recognised the dangers associated with steroid therapy unless patients were S. stercoralis-free and warned medical workers to be aware of this phenomenon. Such treatment, probably causes a shift in the equilibrium between the parasite and the host in favour of the parasite which leads to auto-infection and increased worm burden. They may penetrate and damage many organs of the body, and ultimately lead to the death of the host. This was also true in individuals inflicted with diseases characterised by depressed cell-mediated immunity, e.g. Leprosy and Tuberculosis (Purtillo et al., 1974).

The effects of corticosteroids on experimental S. ratti infections were, therefore, relevant to the clinical problem and to the

understanding of the role of the immune system in preventing auto-infection. The effects of Betamethasone on the course of initial and subsequent infections are reported in this chapter.

Administration of corticosteroids:

Betsolan (Betamethasone sodium phosphate, Glaxo) was administered intraperitoneally. Betsolan is a veterinary form of Betnesol used for the treatment of human conditions such as asthma, hay fever, rheumatoid arthritis, inflammatory skin disorders and other allergies (Mims, 1975). The rats were injected with an initial dose of 2 mg of Betamethasone (20 mg/kg body weight) which was followed by 1 mg (10 mg/kg) every two days for the whole period of the experiment, unless otherwise stated.

Administration of antibiotics:

In all experiments where corticosteroids were employed, 1000 units of penicillin (Triplopen, Glaxo) in 0.1 ml plus 2.5 mg of streptomycin in similar volume, both diluted in sterile distilled water, were administered intramuscularly on the same days as steroid treatment to avoid any opportunistic bacterial infections.

Experiment 8.1:

In this experiment, the effect of Betamethasone on the course of an initial infection was investigated. Eighty 100 gm rats were infected with 1000 (\pm 90) L3 each and then divided into 4 groups of 20. Group 1 received an initial dose of 2 mg of Betamethasone two days before infection, followed by 1 mg every two days for the duration of the experiment. Group 2 was first treated with Betamethasone 9 days PI followed by 1 mg every two days until the end of the experiment. Group 3 was first treated 25 days PI after

which time it received similar treatment to groups 1 and 2. Group 4 was not given steroid treatment. Faecal cultures were prepared on days 8, 11, 15, 18, 22, 25, 29, 32, 36, 39, 43, 46, and 50 PI. Four rats from each group were autopsied on days 10, 17, 24 and 31 PI and the number of worms counted and their lengths measured. The weight of the rats in all groups was monitored during the course of the experiment.

The results obtained are presented in Table 24. Larval production (Figure 30) in the groups which were treated with Betamethasone, from 2 days before and 9 after infection, was very low on day 8 PI. However, the number increased after day 22 PI and persisted with occasional fluctuations. The remaining rats in group 1 died on day 43 PI, with infection in the lungs possibly due to the effects of the drug. The larval production in group 2 increased to a very high peak on day 46 PI (15,800 L3/gm of faeces) and they died on days 51 and 52 PI.

Throughout the experiment, the number of adults recovered (Figure 31) from rats treated with Betamethasone on days -2 and 9 PI was lower than the untreated control. The mean number of adult worms was 98 in group 1 on day 10 PI and 147 in group 2, while the control rats harboured a mean of 430 on the same day. Group 3 before Betamethasone treatment also had a mean number of 404 worms on day 10. On day 24 PI, the control untreated group 4 had only 11 worms in the gut and group 3 had a mean of 12 worms while the corticosteroid-treated rats of groups 1 and 2 harboured 93 and 87 worms per rat, respectively.

When rats in group 3 were treated with Betamethasone on day 25 PI, they showed an immediate rise in the larval output on day 29 PI from 133 to 1970 L3s/gm of faeces. The larval production continued until day 43 PI, though on a lower level.

The number of adult worms in group 3 increased after the administration of Betamethasone on day 25 from a mean of 12 on day 24 to 77.7 on day 31.

The length of worms recovered (Figure 32) from groups 1 and 2 did not alter throughout the period of treatment while those of the untreated control group 4 followed the trend observed previously (see Chapter 3) and decreased in length after day 17 PI. The length of the worms recovered from group 3 after treatment increased from

Table 24 Exp. 8.1:

Effects of Betamethasone treatment on the course of initial infection
with S. ratti. A: Mean (SE) number of adult worms, B: Mean (SE) worm length (in mm.).

Days PI	<u>Group 1</u>		<u>Group 2</u>		<u>Group 3</u>		<u>Group 4</u>	
	A	B	A	B	A	B	A	B
10	98 (±10)	2.83 (±0.03)	147 (±23.7)	2.88 (±0.02)	404 (±11.6)	2.79 (±0.03)	430 (±19.3)	2.86 (±0.03)
17	77.7 (±9.55)	2.81 (±0.03)	78 (±9.6)	2.79 (±0.04)	254 (±20.8)	2.2 (±0.04)	311 (±15.1)	2.03 (±0.05)
24	93 (±19.5)	2.77 (±0.03)	87 (±14.8)	2.71 (±0.04)	12 (±1.8)	0.9 (±0.06)	11 (±1.03)	1.01 (±0.05)
31	93.5 (±7.3)	2.63 (±0.04)	57.7 (±15)	2.78 (±0.03)	77.7 (±13)	2.81 (±0.02)	1.5 (±0.64)	0.8 (±0.1)

Figure 30, Exp. 8.1:

Effect of Betamethasone on the level of larval production during an initial infection with *S. ratti*. ■—■ treated from day -2; □—□ treated from day 9 PI; ○—○ treated from day 25 PI; ●—● infected, not treated.






Figure 31, Exp. 8.1:

Number of adult *S. ratti* in four groups of rats:
 ■—■ treated with Betamethasone from day -2; □—□
 treated from day 9 PI; ○—○ treated from day 23 PI;
 ●—● infected, not treated.

Mean number of
adult worms




Figure 32, Exp. 8.1:

Changes in the length of adult worms recovered from four groups of rats. ■—■ treated from day -2; □—□ treated with Betamethasone from day 9 PI; ○—○ treated from day 23 PI; ●—● infected, not treated.





Figure 33, Exp. 8.1:

Changes in the body weight observed in 4 groups of rats. ■, □, ○ treated with Betamethasone;
● no treatment control; ↓ day Betamethasone treatment commenced.



0.9 mm on day 24 PI to 2.81 mm on day 31.

Corticosteroid treatment resulted in weight loss in all the treated groups (Figure 33). The spleens and mesenteric lymph nodes were also reduced in size. The walls of the small intestine in the treated animals became more translucent and less active and no Peyer's patches were seen. Most of the treated rats developed a diarrhoea and when they died they had patches of pus in their lungs.

Experiment 8.2:

This experiment was designed to repeat the previous experiment with special attention being paid to the distribution of the worms in the intestine in groups treated with corticosteroids before, early and late during infection, as well as the changes in the blood picture during such treatment.

Sixty four rats were infected on the same day with 1000 (\pm 78) L3/rat and were separated into 4 equal groups. Group 1 was treated with corticosteroids two days prior to infection and every two days until the end of experiment. Group 2 commenced similar treatment on day 9 and group 3 on day 23 PI. Faecal cultures were made from all groups on days 5, 7, 10, 17, 21, 25, 28, 31, 35 and 74. Autopsies of four rats from each group were performed on days 15, 23 and 33 PI. The intestine of these rats was divided into anterior and posterior halves to examine the distribution of these worms. In group 1, the larval production (Figure 34) was steady for the first 31 days but increased to over 4000 L3s/gm on day 35 PI and persisted up to 74 days when it reached a maximum of 5800 L3/gm. In group 2, larval output was also steady, although it stayed higher than group 1. The larval production also extended until day 74 as long as the treatment with the immunosuppressant was continued. The larval production in group 3 after being treated with Betamethasone on day 23 PI, increased gradually from 330 L3/gm on day 21 to 840 on day 25 PI and reached a maximum of 3200 on day 74 PI. The number of adult worms (Table 25 and Figure 35) in the three treated groups did not correspond to the increase in the larval

Figure 34, Exp. 8.2:

167

Effect of Dexamethasone on the level of larval production during an initial infection with *S. ratti*. ■—■ treated from day -2; □—□ treated from day 9 PI; ○—○ treated from day 23 PI; ●—● infected not treated.







Table 25 Exp. 8.2:

Effect of Betamethasone on the recovery and distribution of adult worms
in four groups of rats infected with S. ratti

	<u>Group 1</u>			<u>Group 2</u>			<u>Group 3</u>			<u>Group 4</u>		
	<u>Day 15</u>	<u>Day 23</u>	<u>Day 33</u>	<u>Day 15</u>	<u>Day 23</u>	<u>Day 33</u>	<u>Day 15</u>	<u>Day 23</u>	<u>Day 33</u>	<u>Day 15</u>	<u>Day 23</u>	<u>Day 33</u>
Mean (SE) total number of adult worms	127 (+32)	94.7 (+27)	141 (+30)	103 (+22)	105 (+11)	130 (+36)	322 (+13)	116 (+19)	275 (+79)	355 (+25)	100 (+6.6)	0.25 (+0.25)
Mean (SE) number of worms in anterior intestine	122 (+20)	91.7 (+26)	134 (+28)	99 (+20)	102 (+10)	122 (+32)	274 (+19)	1.75 (+0.8)	3.5 (+1.5)	342 (+26)	2.5 (+0.8)	0 -
Mean (SE) number of worms in posterior intestine	4.75 (+1.8)	3 (+1.1)	6.75 (+2.7)	4.25 (+2.3)	3.5 (+1.3)	7.25 (+4)	22.7 (+2.8)	114 (+18)	271 (+78)	13 (+2.1)	97.5 (+6.7)	0.25 (+0.25)
Mean (SE) length of adult worms (mm)	2.81 (+0.03)	2.77 (+0.05)	2.63 (+0.05)	2.75 (+0.04)	2.51 (+0.04)	2.53 (+0.06)	2.5 (+0.04)	1.28 (+0.05)	2.81 (+0.03)	2.48 (+0.05)	1.33 (+0.05)	- -

Figure 35, Exp. 8.2:

Recovery of adult *S. ratti* from the anterior and posterior intestines of rats treated with betamethasone: Group 1: commenced treatment on day -2; Group 2: commenced treatment on day 9 PI; Group 3: on day 23 PI; Group 4: not treated.



production. The number of adult worms in groups 1 and 2 were 127 on day 15, 94.7 on day 23 and 141 on day 33 PI. Rats in group 2 harboured a mean of 103, 105 and 130 worms on days 15, 23 and 33 PI, respectively. The distribution of the worms was constant in both groups 1 and 2, concentrating in the anterior half of the intestine throughout the period of corticosteroid treatment. The worms migrated from the anterior segment of the intestine (where they were mainly found on day 15 PI) in groups 3 and 4 to the posterior half as shown on day 23 PI. However, after group 3 was treated with Betamethasone on day 23 PI, the number of adult worms increased from 116 on day 23 to 275 on day 33 PI when the normal infection control group harboured a mean of 0.25 worms. The majority of the adult worms in group 3, on day 33, were found to occupy the last half of the intestine unlike the worm population in groups 1 and 2 on the same day. The distribution of the strings of eggs, which are laid by the females in the mucosa of the intestine (see Chapter 1) followed that of the adult worms particularly in the Betamethasone-treated groups. The worms in groups 1 and 2 did not shrink during the entire period of treatment, while those in groups 3 and 4 showed signs of such morphological damage on day 23 PI. However, the worms obtained on day 33 PI from group 3 had normal lengths (mean 2.81 mm). This again indicated that an auto-infection cycle might have been the reason for the rise in the number of worms and the change in their lengths following treatment with Betamethasone on day 23 PI.

The total white blood cell counts (Figure 36) were very low in groups 1 and 2 throughout the corticosteroid treatment. The WBC count in group 3 and 4 increased sharply on day 23 PI (prior to worm expulsion). However, the count dropped in group 3 after treatment with Betamethasone on day 23, and in group 4 on day 33 PI following the termination of the initial infection.

The differential white blood cell count (Table 26) on day 15 PI in all the groups, revealed that Betamethasone had a suppressive effect on lymphocytes and eosinophils, but there was no significant difference in the percentage of neutrophils and monocytes in all the four groups.

This experiment demonstrated that treatment with corticosteroids caused a delay in expulsion of S. ratti worms for as long as 49 days

Figure 36, Exp. 8.2:

Effect of Betamethasone treatment on the W.B.C. count in four groups of infected rats. ■—■ treatment commenced on day -2; □—□ treatment commenced on day 9 PI; ○—○ treatment commenced on day 23 PI; ●—● no Betamethasone treatment.




Table 26, Exp. 8.2:

Effect of Betamethasone, on day 15 of an initial S. ratti infection,
on the differential white blood cell (W.B.C.) count

Groups: (day treatment commenced)	% lymphocytes	% monocytes	% neutrophils	% eosinophils
1 (-2)	0	0	16	0
2 (9 PI)	0	0	18	0
3 (25 PI)	73	1	17	1
4 (No treatment)	75	1	16	1

when administered before or after infection. It also demonstrated that Betamethasone can cause a rise in the number of worms suggestive of auto-infection. The movement of worms from the anterior to the posterior half of the small intestine during the course of infection was inhibited by Betamethasone.

Experiment 8.3:

This experiment was designed to study the effect of ceasing immunosuppressive treatment on the restoration of the immune reaction in infected rats. It also investigated the effects of Betamethasone on the level of tissue eosinophils and on the degree of mesenteric mastocytosis. Forty eight rats were divided into three groups of 16 rats each, and infected with 1000 (\pm 77) L3 each. Another four rats of the same age were left as normal uninfected controls. Group 1 was treated with Betamethasone every two days commencing on day 0 until the end of the experiment while group 2 had similar treatment which was terminated on day 24 PI. Group 3 was left untreated. Four rats were killed from groups 1, 2 and 3 on days 10, 20, 25 and 32 PI and the number of adult worms counted. The number of tissue eosinophils and the level of mastocytosis was determined on day 25 PI when the four normal uninfected rats were killed. The mean number of adult worms (Table 27 and Figure 37) in both groups 1 and 2, during the period of Betamethasone treatment was 208 and 85 on day 10 and 134 and 138 on day 20 PI. The control group harboured 422 and 236 worms on the same days. One day after the withdrawal of the treatment from group 2, the worm burden was not different from that of group 1 (128 in group 2 and 101 in group 1), while the control group had a mean of 10 worms. However, the rats in group 2 harboured only 20 worms by day 32 PI, while group 1, which was receiving Betamethasone continuously, maintained the number of worms at 117. The normal infected, untreated, control group 3 had a mean of 0.25 worms on the same day. The number of tissue eosinophils (Figure 38) was quantified in a similar manner to that described in Chapter 8, i.e., dividing the intestinal villi into villus-crypt units and counting the number of eosinophils

Table 27 Exp. 8.3:

Effect of the withdrawal of Betamethasone on the course
of initial infection with S. ratti

Groups	Mean (SE) number of adult worms			
	<u>Day 10</u>	<u>Day 20</u>	<u>Day 25</u>	<u>Day 32</u>
1	208 (+38)	134 (+31)	101 (+11)	117 (+24)
2	85 (+17)	138 (+31)	128 (+19)	19.5 (+8)
3	422 (+21)	236 (+25)	10 (+1.9)	0.25 (+0.25)

Figure 37, Exp. 8.3:

Number of adult worms in three groups of infected rats.
 ■—■ treated with Betamethasone until day 32 PI; ○—○
 treated until day 24 PI; ●—● untreated control.

175





Figure 38, Exp. 8.3:

Effect of Betamethasone on the number of tissue eosinophils and percentage of mesenteric mast cell degranulation, on day 25 PI.

■ infected, treated with Betamethasone; ▨ infected non-treated; □ normal uninfected, untreated rats.



within 10 units. The rats of group 1 which were treated with Betamethasone and examined on day 25 PI, showed a mean of 0.88 tissue eosinophil per 10 VCU (Plate 54). The normal infection controls had a mean of 290 eosinophils/10 VCU and the normal uninfected rats had a mean of 92 eosinophils/10 VCU on the same day (see Plates 50 and 51, Chapter 8). No immune reaction was observed around sections of adults, larvae or embryonated eggs in the intestines of treated rats (Plate 55).

The rats of group 1 had a mean percentage of 26% degranulating mesenteric mast cells, while rats in group 3 showed a higher degree of degranulation on the same day. The normal rats of group 4 had a mean of 19% degranulating mast cells in the mesenteries on the same day (Figure 37). Macroscopically, the mesenteric lymph nodes of rats treated with Betamethasone (Plates 56 and 57) were capable of restoring their shape and size after the termination of treatment and sections of such nodes were similar to those of the untreated controls and contained numerous lymphocytes and precursor cells (Plates 58 and 59).

Experiment 8.4:

This experiment investigated the effect of corticosteroid treatment on a challenge infection into previously infected rats which had received Betamethasone after they had expelled their worm burden.

Thirty two rats were each infected with 1000 (\pm 85) L3 each and then divided into 4 groups of 8. Group 1 received 2 mg Betamethasone every two days commencing on the day of infection. Group 2 received 4 mg of Betamethasone on days 28, 29, and 30 PI. Group 3 received 4 mg of Betamethasone on days 29 and 30 PI and on day 3 post-challenge (see later). Group 4 received no Betamethasone. A fifth group of 4 rats was left uninfected and untreated to act as challenge control. Four rats from the first four groups were killed on day 13 PI. All five groups were treated with a single dose of 200 mg/kg of TBZ on day 26 PI and challenged with 1000 (\pm 97) L3 each on day 30. The rest of the rats in all




Plate 54 : Section through villi of rats treated with Betamethasone; note the absence of eosinophils (Lendrum's, x 250)




Plate 55 : Embryonated eggs embedded in the crypts of intestinal mucosa; note lack of reaction (Lendrum's, x 250)




Plate 54 : Section through villi of rats treated with Betamethasone; note the absence of eosinophils (Lendrum's, x 250)




Plate 55 : Embryonated eggs embedded in the crypts of intestinal mucosa; note lack of reaction (Lendrum's, x 250)




Plate 56 : Mesenteric lymph node treated with Betamethasone
(H. & E., x 80)




Plate 57 : Types of cells present in MLN treated with
Betamethasone (H. & E., x 250)




Plate 56 : Mesenteric lymph node treated with Betamethasone
(H. & E., x 80)




Plate 57 : Types of cells present in MN treated with
Betamethasone (H. & E., x 250)




Plate 58 : MLN from a non-treated control rat (H.& E., x 30)




Plate 59 : Types of cells present in normal MLN (H.& E., x 250)




Plate 58 : MLN from a non-treated control rat (H. & E., x 30)




Plate 59 : Types of cells present in normal MLN (H. & E., x 250)

groups were autopsied on day 13 post-challenge. In both autopsies, the number of adult worms in the anterior and posterior small intestine was counted and their lengths measured. Faecal cultures from all groups were made on day 13 post-challenge. The results are presented in Table 28).

On day 13 PI, the rats of group 1 (i.e. continuous Betamethasone treated) had the lowest adult count (mean 128) while groups 2, 3 and 4 had mean totals of 406, 410 and 433, respectively. Worms in all groups were concentrated in the anterior small intestine.

Following challenge, 145 worms were recovered from group 1 and these were distributed anteriorly. The mean length of the adult worms was 2.9 mm, and their fecundity was very high at 26.3 larvae/worm. The rats of group 2 had a mean total of 19.5 worms, the majority of which were occupying the posterior half of the gut. The worms were stunted with a mean length of 1.8 mm and showed a very low fecundity rate (2.6 larvae/worm). The rats in group 3, which received corticosteroids on days 29, 30 PI and day 3 post-challenge harboured a mean of 388 worms of normal (2.8 mm) length. Their larval production was high with a fecundity rate of 19.7 larvae/worm. No adults were recovered from group 4 on day 13 post-challenge while 373 worms were collected from the challenge control group 5, mostly in the anterior intestine.

This experiment demonstrated that continuous treatment of Betamethasone during initial infection completely suppressed the development of any acquired resistance to reinfection. Corticosteroid treatment when introduced on three consecutive days prior to the introduction of the challenge dose was unable to destroy the immune memory against an initial infection with the parasite. However, the drug was effective in suppressing the manifestation of acquired immunity when introduced on two consecutive days before the secondary infection followed by another dose after the challenge.

Experiment 8.5:

This experiment attempted to induce the delay of expulsion of S. ratti worms by treatment with corticosteroids at different times

Table 28, Exp. 8.4: Effect of Betamethasone treatment on the course of initial and second infections with *S. ratti*, on days 13 PI (A) and 13 post-challenge (B)

	<u>Group 1</u>		<u>Group 2</u>		<u>Group 3</u>		<u>Group 4</u>		<u>Group 5</u>	
	A	B	A	B	A	B	A	B	A	B
Mean (SE) total number of adults	128 (+27.6)	144 (+13.4)	406 (+26)	19.5 (+8.4)	410 (+18.6)	388 (+60.2)	433 (+19.5)	0	0	373 (+20)
Mean (SE) number in anterior half	122 (+26)	141 (+13)	398 (+26)	2.25 (+2)	401 (+18)	301 (+35.6)	429 (+20.8)	0	0	363 (+19)
Mean (SE) number in posterior half	5 (+1.8)	3 (+0.7)	8 (+1.5)	17.2 (+6.5)	9 (+2)	87.2 (+60.3)	9.5 (+2.1)	0	0	10 (+2.3)
Mean (SE) length of adults (mm)	N.D.	2.9 (+0.03)	N.D.	1.8 (+0.04)	N.D.	2.8 (+0.03)	N.D.	0	0	2.8 (+0.03)
Mean (SE) number of larvae/gm	N.D.	3821 (+231)	N.D.	51 (+8)	N.D.	4380 (+371)	N.D.	0	0	3906 (+216)
Fecundity (larvae/worm)	N.D.	26.3	N.D.	2.6	N.D.	19.7	N.D.	0	0	21

during the course of an initial infection. Thirty eight rats were infected on the same day with 1000 (\pm 53) L3/rat and were divided into four groups:

Group 1: Ten rats received 2 mg Betamethasone every two days from the day of infection until day 23 PI. Group 2: Ten rats received 4 mg of Betamethasone on days 4, 6, 8 and 10 PI. Group 3: Ten rats received 4 mg Betamethasone on days 6, 10, 14 and 18 PI. Group 4: Eight rats were left untreated as controls.

Half of each group was killed on days 30 and 35 PI and the number of adult worms was counted. The results are presented in Table 29.

Very few worms were recovered from the control group 4, on day 30 PI, and no worms were recovered from this group on day 35. Rats in group 1 had a mean worm burden of 50 on day 30 and this number had dropped to 7.8 worms on day 35 PI. The rats in group 2, which were treated with Betamethasone on days 4, 6, 8 and 10 harboured a mean of 597 on day 30 PI and 537 on day 35. The members of group 3 harboured a statistically significant mean of 225 worms on day 30 PI, but the number dropped sharply to 16 on day 35.

This experiment demonstrated that treatment with Betamethasone on days 4, 6, 8 and 10 delayed the action of expulsion of adult S. ratti worms up to day 35 PI. Betamethasone injected every two days for a period of 23 days did not induce similar delays in worm expulsion, and when introduced on days 6, 10, 14 and 18 only resulted in a partial delay of worm elimination.

Discussion:

The experiments in this chapter have demonstrated that corticosteroids have an immunodepressive effect on the course of S. ratti infection. If the drug was used early during infection there was no increase in the number of the adult females in the gut, in fact, worm recovery was usually very low. This might be attributed to different factors; The action of Betamethasone on the mucosa of the gut was severe, whether introduced IP or IM. The villi of

Table 29

Exp: 8.5 Effect of betamethasone treatment when administered at specific times during primary infection

	Group 1		Group 2		Group 3		Group 4	
	Day 30 PI	Day 35 PI	Day 30	Day 35	Day 30	Day 35	Day 30	Day 35
Number of adult worms in individual rats	91	3	470	551	169	14	4	0
	78	1	721	671	351	7	0	0
	17	15	660	451	200	23	31	0
	23	12	501	533	215	20	12	0
	41	8	636	481	191	16	-	-
Mean (SE)	50(+14.7)	7.8(+ 2.6)	597(+48)	537(+37.8)	225(+32.3)	16(+2.7)	11.7(+6.8)	0

treated rats were much smaller than those of normal controls. It is also possible that the immunodepressive action of Betamethasone allowed multiplication of bacteria in the gut and that they adversely affected S. ratti. Treated rats had continuous diarrhoea or loose faeces. It was hoped that infected rats treated with immunodepressive drugs, would allow the parasite to undergo auto-infection and, thus, harbour more adult worms than the original inoculum, as seen in human cases of infected immunodepressed individuals (Rogers and Nelson, 1966; Cruz et al., 1966; Willis and Mucolo, 1966; Civantos and Robinson, 1969; Yim et al., 1970; Fagundes et al., 1971; Bitoun et al., 1972; Cookson et al., 1972; Neefe et al., 1973; Purtillo et al., 1974). This did not occur when treatment was begun early during infection. However, there was an increase in the number of adult worms when rats were treated around the time of worm expulsion. These worms were fertile and not stunted. Since it was shown that the changes in the morphology of adult S. ratti during the final stages of an initial infection were irreversible (see Chapter 4), this increase in number and size of adult worms indicated that corticosteroid treatment caused auto-infection. It was not clear whether the same treatment in groups 1 and 2 had caused the same cycle, due to the low number of worms recovered.

Adult S. ratti persisted in rats treated with Betamethasone for as long as the treatment continued. Enhancement of worm survival in animals was obtained following cortisone treatment in other nematode (Weinstein, 1953, 1955; Ogilvie, 1965; Stoner and Goodwin, 1953, 1954; Coker, 1955, 1956; Markell and Lewis, 1957; Markell, 1958; Cross, 1960; Campbell and Collette, 1962; Campbell, 1963, 1968; Wakelin, 1970, 1974; Avlavidou, 1970; Behrke, 1975), trematode (Tewari, 1970) and cestode infections (Hopkins et al., 1972; Moss, 1972). Similar results were obtained using more specific methods of immunosuppression such as neonatal thymectomy, irradiation, anti-lymphocyte serum and thoracic duct drainage (Ogilvie and Jones, 1967; Dineen and Adams, 1970; Hopkins et al., 1972; Di Natta et al., 1972).

The precise mechanisms of such immune suppression are still undetermined, but it seems that both arms of the immune response are affected. It has been suggested that T. cells, which have been shown to be involved in worm expulsion (see Chapter 6), are resistant

to cortisone (Cohen *et al.*, 1970; Cohen and Claman, 1971). However, North (1972) suggested that a population of T. cells were sensitive to the suppressive effects of cortisone. The action of Betamethasone on *S. ratti* may be aimed at such a population and possibly at the extra factors e.g. eosinophils and pharmacologically active substances (see Chapter 7), which might play some role in worm expulsion.

Betamethasone treatment inhibited the posterior movement of the worms which occurs normally during the course of an initial infection (see Chapter 3) and most adult worms in treated rats were found in the anterior intestine. In group 3 of experiment 8.2, the recovery of a majority of normal, fertile, adult worms from the posterior small intestine following treatment on day 23 might be attributed to the fact that auto-infection took place within the intestine and not by migration to the lung as classically known (Faust and De Groat, 1940). It is possible that when the immune status of the host is low, intestinal rhabditiform larvae moult twice in the lumen and penetrate the gut mucosa, where they undergo two more moults and develop into adult females. It is known that in *S. ratti* no moulting occurs in the lung (Wartheim, 1965), which suggests that the pulmonary stage is not critical. This hypothesis does not reject the possibility of pulmonary migration during auto-infection but offers an explanation for the unexpected recovery of new, normal, adult worms in the pre-expulsion site normally occupied by damaged worms. These experiments have also shown that structural and morphological damage of worms was not genotypic but phenotypic since all the worms which developed following treatment with Betsolan at the terminal period of an infection, were fertile, and normal in size.

The recovery of rats after the withdrawal of corticosteroid treatment was fast. Within 8 days of the cessation of treatment, rats were able to expel the majority of the worms from their guts and restore the shape and size of their lymph nodes. Restoration of immunity and abrupt expulsion of worms was also observed in other helminthic infections, following the withdrawal of cortisone treatment (Coker, 1955; Ogilvie, 1965; Hopkins *et al.*, 1972).

Continuous treatment with Betamethasone during an initial

infection with S. ratti suppressed all acquired resistance to reinfection. However, treatment was necessary prior to and after the introduction of the challenging dose, in rats untreated previously, to eliminate the immunological memory and suppress the immune response against a second infection. Wakelin (1970) showed that immunological unresponsiveness to T. muris could be achieved if cortisone was given at the time where the immune system was set to recognize the antigenic stimulus produced by the worms. It was later possible to achieve immune tolerance by subjecting infected mice to four injections of cortisone acetate on days 8, 10, 12 and 14 PI, or by giving a single large dose on day 11 PI (Wakelin and Selby, 1974).

Behnke (1975b) obtained long-term survival of Aspicularis tetraptera in mice given a short term hydrocortisone treatment. He suggested that this treatment brought about a state of tolerance, due to the suppression of the action of recognition, by the host, of the antigens released during the cryptic-phase (Behnke, 1975a) of infection. Olivier (1962) working with Taenia taeniformis in mice obtained a long-lasting tolerance to this parasite following few cortisone injections.

In experiment 8.5, the number of adult worms on day 35 was still high in the group treated with Betamethasone on days 4, 6, 8 and 10 PI rather than days 6, 10, 14 and 18 PI. This is not a final indication of immune tolerance in this group but could give a provisional indication that corticosteroid treatment on days 4, 6, 8 and 10 only can paralyse the normal immune response against S. ratti and render the rats unresponsive for a period of 35 days, at least.

There is still a wide field of research to be explored concerning the effects of immune suppression in this host-parasite system, particularly as concerns the mechanisms involved in the cortisone-induced auto-infection. Apart from chemical suppression, the role of the specific suppression of T. and B. cells using thymectomy and whole body irradiation, as well as anti-lymphocyte or anti-thymocyte serum, requires investigation.

CHAPTER 9Summary and Conclusions

The experiments carried out during this project can be classified under four major headings:

1. Those which studied parasitological aspects of acquired immunity during primary, secondary or multiple infections.
2. Those which studied the effects of the immune response on the adult worm population.
3. Those which considered some of the mechanisms of worm expulsion.
4. Those which investigated the effects of immunosuppressive treatment on the course of primary and secondary infections.

Following a primary infection, rats developed strong resistance to S. ratti which manifested itself by a sharp drop in larval production after day 14 PI and a mass expulsion of the parthenogenetic adult females, although a few adult worms persisted for a period as a residual population. The worms also migrated towards the distal half of the small intestine, before expulsion.

Following a challenge infection, S. ratti was expelled from the gut within 10 days and immediately before expulsion, the worms occupied the posterior half of the intestine.

Resistance to reinfection with S. ratti was maintained in rats for a period of 3 months. The residual population of worms had no apparent role in this.

The host response during an initial infection was aimed at the adult worms in the gut. However, during a second infection, there was a severe cellular reaction around the migratory stages in the lungs, where larvae were trapped and killed. Only few larvae avoided the serious response in the lungs, and developed in the gut into stunted and almost sterile worms.

Following multiple exposures, the immune response killed the infective larvae in the skin and there was a generalized dermal

reaction with infiltration of eosinophils and lymphocytes.

A full infection with S. ratti gave complete protection to the rats against reinfection. 10 day-old adult females also conferred a substantial degree of protection. The migratory stages did not stimulate immunity.

Worms which were obtained from infected rats 8-16 days I.P. were 2.88 mm in length and their uteri were full of embryonated eggs. When these worms were transplanted into clean rats, they continued their egg production as if they were in the original host.

Worms which were harvested from infected rats, between days 18-25 PI, were much shorter in length averaging 0.9 mm and contained no eggs. When these worms were transplanted, into new rats, they did not recover and were expelled immediately, indicating that they had been irreversibly damaged.

These two types of worms were studied with light and electron microscopy. This revealed that the damaged worms had a disorganised endoplasmic reticulum with increased numbers of lysosomes and vacuoles. No fat droplets were seen but whorls of lipoprotein were spotted, invariably, within the lumen of damaged worms. These were suggested to be immune complexes, which had been either formed internally, or swallowed. Precipitates around the mouth of the worms fluoresced in a direct F.A.T., showing them to be immune complexes, but it was not possible to demonstrate specific fluorescence of the lumen contents of the guts in sections of damaged worms.

When short and stunted worms, obtained early during a secondary infection, were transferred to new rats, they were able to establish themselves. They renewed egg production, which had ceased in the original hosts. Worm adaptation, similar to that observed in N. brasiliensis (Ogilvie, 1972), was discussed as a possibility.

Experiments were carried out to investigate the mechanisms of worm expulsion. It was concluded that worm expulsion was an immunological event, since immune MINC were capable of inducing a reduction in the worm burden. MINC transferred on day 12 PI were more effective than those transferred on day 6. Immune serum had a blocking effect and, in 3 out of 4 experiments, resulted in

auto-infection in the recipient rats. It is suggested that antibodies may paralyse the cellular arm of the immune response by either masking the worms, and possibly larvae, or through producing specific suppressive factors, which inhibit the function of the specific population of lymphocytes. However, immune serum caused a reduction in the size of the adult worms given early during infection.

Auto-infection also occurred in rats which received immune serum and MNC on day 6 PI, but the best rate of worm expulsion was recorded when serum and MNC were transferred on day 12 PI.

Immune MNC were capable of conferring a substantial degree of protection when given 1 day before infection, but better protection was obtained when immune serum was added on day 0. However, immune serum alone given on day 0 did not confer any significant degree of protection. It was demonstrated that tissue eosinophilia increased during the period of worm expulsion. No success was achieved in repeated endeavours to stain and demonstrate intestinal mast cells during the period of expulsion, but studies on the level of degranulation of mesenteric mast cells revealed that there was a very significant rise in the percentage of degranulating mast cells in the adjoining tissues during worm expulsion in initial and second infection. There was a parallel rise in the total white blood cell count, but blood eosinophils did not follow this pattern. The number of peripheral lymphocytes was very high in infected rats.

Histamine injected directly into the stomach of infected rats caused a slight acceleration of both worm damage and expulsion. Aspirin, a potent inhibitor of prostaglandins, when introduced daily, per os, delayed worm expulsion by 5 days. These two results suggest a role for both histamine and prostaglandins in the elimination of S. ratti from the rats.

The final sets of experiments dealt with the effects of immunosuppressive corticosteroid treatment on initial and subsequent infections with S. ratti. Corticosteroid treatment caused a long persistence of the worm burden during an initial infection, for as long as the treatment was continued. The withdrawal of the drug caused the rats to recover and expel their infection in a

short period. No auto-infection was demonstrable when corticosteroids were introduced early during infection and continuously every two days. In fact the number of adult worms in these rats was very low compared to the untreated controls. However, when corticosteroids were introduced 2 days before worm expulsion (day 23 PI), auto-infection occurred and there was an increase in the number of adult worms, these being normal in their measurements and anatomy. This indicated that worm damage was phenotypic and due to an immune phenomenon rather than senescence. Larval production of worms in rats treated with corticosteroids was maintained for the duration of continuous treatment. A period of unresponsiveness was induced when rats were treated with corticosteroids at strategic points during infection. Adult worms persisted up to day 35 PI when rats were treated with 4 large doses of Betamethasone on days 4, 6, 8 and 10 PI.

S. ratti infections are like many nematode infections of rodents in that they are expelled from the intestine of the host within 2 or 3 weeks of arriving in the gut. The major complication of the host-parasite relationship is the possibility of auto-infection which does not occur in other nematode systems.

The parasite S. ratti has proved a very good laboratory model for the study of aspects of acquired immunity to nematode infections.

REFERENCES

- Abadie, S. H. (1963). The life cycle of Strongyloides ratti. J. Parasitol., 49, 241-248.
- Abe, Y., Kagei, N., Hori, E., Kawamoto, T., Miyagami, S., Tagawa, N., Mukai, T. and Arikawa, S. (1960). Studies on Strongyloides and Strongyloidiasis. Distribution of Strongyloides ratti Sandground, 1925 in house rats in Southern Kyushu. Kagoshima Med. J., 33, 446-461.
- Alfieri, F. (1908). Intorno alla genesi della "Anquillula intestinalis" de la rana. Glor. R. Accad. Med. Torino, 14, 265-267.
- Ali-Khan, Z. and Seemayer, T. A. (1975). Fatal bowel infraction and sepsis: an unusual complication of systemic strongyloidiasis. Trans. R. Soc. trop. Med. Hyg., 69, 473-476.
- Allison, F. (1965). Anti-inflammatory agents: In Zwerfach, B., Grant, L. and McCluskey (eds.) The inflammatory process. Academic Press, New York. pp. 559.
- Alphay, T. J. W. (1970). Studies on the distribution and site of location of Nippostrongylus brasiliensis within the small intestine of laboratory rats. Parasitology, 61, 449-460.
- Amano, R., Mizuno, H., Tange, H. and Tanaka (1959). The daily course of the infesting number of Strongyloides ratti after experimental infection in albino rats. Ochanomizu Igaku Zasshi, 7, 2928-2931.
- Anderson, C. V. and Leonard, A. B. (1940). Immunity produced in rats by the intestinal phase of Trichinella infections. J. Parasitol., 26, 42.
- Araujo, P. (1967). Distribucao de helmintos no intestino delgado de Rattus norvegicus naturalmente infestados. II. Strongyloides ratti, Sandground, 1925 e S. venezuelensis Brumpt, 1934. Revista de la Facultad de farmacia e bioquímica da Universidad de Sao Paulo, Brasil, 5, 179-190.
- Archer, R. K. (1956). The eosinophil response in the horse to intramedullary and intradermal injections of histamine, ACTH and cortisone. J. Path. Bact., 72, 87-94.

- Archer, R. K. (1963). The eosinophil leucocytes. Blackwell Scientific Publications, Oxford.
- Archer, R. K. (1965). On the functions of eosinophils in the antigen-antibody reaction. Br. J. Haematol., 11, 123-129.
- Archer, R. K. (1970). Regulatory mechanisms in eosinophil leukocyte production, release and distribution. In: Gordon, A. S. (ed), Regulation of haematopoiesis. Appleton-Century-Crofts, New York. pp. 925-941.
- Archer, Q. T. and Bosworth, N. (1961). Phagocytosis by eosinophils following antigen-antibody reactions in vitro. Aust. J. Exptl Biol., 39, 157-164.
- Ashton, N. and Cook, C. (1952). In vivo observations of the effect of cortisone upon the blood vessels in rabbit ear chambers. Bri. J. Exptl Pathol., 33, 445-450.
- Askanazy, M. (1900). Ueber Art und Zweck der Invasion der Anquillula intestinalis in die Darmwand. Centralbl. F. Bakt. Parasitenk., 27, 569-578.
- Avlavidou, T. (1970). Inhibiting action of hydrocortisone on the response of rabbits immunized with antigens from whole Ascaris suum. Scripta Sci. Med., 8, 171-174.
- Bagai, R. C. and Subrahmanyam, D. (1968). Studies on the host-parasite relation in albino rats infected with Litomosoides carinii. Amer. J. trop. Med. Hyg., 17, 833-839.
- Bailenger, J. and Caroenac, F. (1974). Repercussions of parasitism by Strongyloides ratti on the secretion of the glucocorticosteroids in the rat. Internat. J. Parasitol., 4, 307-310.
- Bailenger, J. and Faraggi, G. (1975a). Mechanism of the hypocorticosteronemic action of Strongyloides ratti in the rat. Ann. Parasit. hum. comp., 50, 187-197.
- Bailenger, J. and Faraggi, G. (1975b). Action of Reserpine on parasitism of the rat by S. ratti: An attempt to explain the mechanism of verminous hyper-corticosteronemia. Ann. Parasit. hum. comp., 50, 199-208.
- Barth, E. E. E., Jarrett, W. F. H. and Urquhart, G. M. (1966). Studies on the mechanism of the self-cure reaction in rats infected with Nippostrongylus brasiliensis. Immunology, 10, 459-464.

- Basir, M. A. (1950). The morphology and development of the sheep nematode Strongyloides papillosus (Wedl, 1856). Can. J. Res., 28, 173-196.
- Baylis, H. A. (1923). Report on a collection of parasitic nematodes, mainly from Egypt. Part III. Parasitology, 15, 24-38.
- Beach, T. D. (1935). Experimental propagation of Strongyloides in culture. Proc. Soc. exp. Biol. Med., 32, 1484-1486.
- Beach, T. D. (1936). Experimental studies on human and primate species of Strongyloides. V. The free living phase of the life cycle. Amer. J. Hyg., 23, 243-277.
- Behar, R. R. (1970). Complications of Strongyloidiasis. Revta Cuba Pediat., 42, 187-209.
- Behar, R. R. (1971). Complications of strongyloidiasis in infancy. Revta Cuba Pediat., 43, 147-165.
- Behnke, J. M. (1975). Suppression of expulsion of Aspicularis tetraptera in hydrocortisone and methotrexate treated mice. Parasitology, 71, 109-116.
- Bennett, A. (1972). Effects of prostaglandins on the gastro-intestinal tract; In: Karim (Ed). The prostaglandins, progress in research. Medical and Technical Publishing Co., London. pp. 205-221.
- Berglund, K. (1962). Inhibition of antibody formation by prednisolone: location of a short sensitive period. Acta path. microbiol. Scand., 55, 187-202.
- Berkmen, Y. M. and Rabinowitz, J. (1972). Gastrointestinal manifestations of the strongyloidiasis. Amer. J. Roentgenology, 115, 306-311.
- Bartram, D. S. (1966). Dynamics of parasitic equilibrium in cotton rat filariasis. Adv. Parasitol., 4, 255-319.
- Bezubik, B. (1960). Effect of cortisone on the susceptibility of hamsters and guinea-pigs to the sheep and rabbit strains of Strongyloides papillosus. J. Parasit., 46, 30-31.
- Bezjack, B. (1968). A clinical trial of thiabendazole in strongyloidiasis. Amer. J. trop. Med. Hyg., 17, 733-736.
- Bitoun, A., Foupon, R., Wilds, J. L. and Rault, P. (1972). Severe strongyloidiasis revealed by corticosteroid therapy. Report of one case. Nouvelle Presse Medicale, 1, 1935-1937.

- Bloom, G. and Kelly, J. W. (1960). The copper phthalocyanin dye "Astrablan" and its staining properties, especially the staining of mast cells. Histochemi, 2, 48-57.
- Bolla, R. I. and Roberts, L. S. (1968). Gametogenesis and chromosomal complement in Strongyloides ratti (Nematoda: Rhabdiasoidea). J. Parasit., 54, 849-855.
- Brackett, S. and Bliznick, A. (1949). An attempt to adapt Strongyloides ratti to the mouse. J. Parasit., 35, 41-44.
- Brambell, M. R. (1965). The distribution of a primary infestation of N. brasiliensis in the small intestine of laboratory rats. Parasitology, 55, 313.
- Brenner, K. C., Ogilvie, B. M., Keith, R. K. and Berrie, D. A. (1973). Acetylcholinesterase secretion by parasitic nematodes. III. Oesophagostomum spp. Internat. J. Parasit., 3, 609-618.
- Brown, H. W. and Perna, V. P. (1958). An overwhelming Strongyloides infection. J. Amer. Med. Ass., 168, 1648-1651.
- Brumpt, E. (1922). Precis de Parasitologie. 3rd ed., Paris.
- Butterworth, A. E., Sturrock, R. F., Houba, V. and Rees, P. H. (1974). Antibody dependent cell-mediated damage to schistosomulae in vitro. Nature, 252, 503-505.
- Butterworth, A. E., Sturrock, R. F., Houba, V., Mahmoud, A. A. F., Sher, A. and Rees, P. H. (1975). Eosinophils as mediators of antibody-dependent damage to schistosomulae. Nature, 256, 727-729.
- Cahill, K. M. (1967). Thiabendazole in massive strongyloidiasis. Amer. J. trop. Med. Hyg., 16, 451-453.
- Cameron, D. G. and Watson, G. M. (1949). The blood counts of the adult albino rat. Blood, J. Haemat., 4, 816-819.
- Campbell, W. C. (1963). Spontaneous cure in Trichuris muris infections in albino mice and its suppression by cortisone. J. Parasit., 49, 628-632.
- Campbell, W. C. (1965). Immunising effect of enteral and enteral-parenteral infections of Trichinella spiralis in mice. J. Parasit., 51, 185-194.
- Campbell, W. C. (1967). Distribution of Trichinella spiralis in the small intestine of young mice. J. Parasit., 53, 395-397.
- Campbell, W. C. (1968). Effect of anti-inflammatory agents on spontaneous cure of Trichinella and Trichuris in mice. J. Parasit., 54, 452-456.

- Campbell, W. C. and Collette, J. V. (1962). Effect of cortisone upon infection with Trichuris muris in albino mice. J. Parasit., 48, 933-934.
- Campbell, W. C., Hartman, R. K. and Cuckler, A. C. (1963). Effects of certain antihistamine and antiserotonin agents upon experimental trichinosis in mice. Expl Parasit., 14, 23-28.
- Casey, W. J. and McCall, C. E. (1971). Suppression of the cellular interactions of delayed hypersensitivity by corticosteroid. Immunology, 21, 225-231.
- Chaia, G. (1963). Pharmacological activity of Thiabendazole ~~2~~- (4-thiazolyl)-benzimidazole⁷ in rats experimentally infected with Strongyloides ratti. Rev. Inst. Med. trop. S. Paulo, 5, 154-159.
- Chaia, G. and Cunha, A. S. DA. (1966). Experimental therapy with Thiabendazole in strongyloidiasis. Rev. Inst. Med. trop. S. Paulo, 8, 17-21.
- Chaia, G. and Murta, C. C. (1967). Immunization of rats with irradiated Strongyloides larvae. Rev. Inst. Med. trop. S. Paulo, 9, 163-168.
- Chandler, A. C. (1925). The species of Strongyloides (Nematoda). Parasitology, 17, 426-433.
- Chandler, A. C. (1935). Studies on the nature of immunity to intestinal helminths. I. The local nature of the immunity of white rats to Nippostrongylus infection. Amer. J. Hyg., 22, 157-168.
- Chandler, A. C. (1936). Studies on the nature of immunity to intestinal helminths. III. Renewal of growth and egg production in Nippostrongylus muris after transfer from immune to non-immune rats. Amer. J. Hyg., 23, 46-54.
- Chandler, A. C. (1938a). Diploscaptor coronata as a facultative parasite of man, with a general review of vertebrate parasitism by rhabditoid worms. Parasitology, 30, 44-55.
- Chandler, A. C. (1938b). Further experiments on passive immunity of rats to Nippostrongylus muris infections. Amer. J. Hyg., 28, 51-62.
- Chandler, A. C. and Read, C. P. (1961). Introduction to Parasitology (10th edition). Toppan Company, Ltd., Tokyo. 822 p.
- Chang, P. C. H. and Graham, G. L. (1957). Parasitism, parthenogenesis and polyploidy: the life cycle of Strongyloides papillosus. J. Parasit., 13, 13.

- Chitwood, B. G. and Chitwood, M. B. (1950). An introduction to nematology. Section I. Anatomy. Monumental Printing Co., Baltimore. 213 p.
- Chitwood, B. G. and Graham, G. L. (1940). Absence of vitelline membranes on developing eggs in parasitic females of Strongyloides ratti. J. Parasit., 26, 183-190.
- Chitwood, B. G. and McIntosh, A. (1934). A new variety of Alloionema (Nematoda: Diplogasteridae), with a note on the genus. Proc. Helm. Soc. Wash., 1, 37-38.
- Civantos, F. and Robinson, M. J. (1969). Fatal strongyloidiasis following corticosteroid therapy. Amer. J. Dig. Dis., 14, 643.
- Cochrane, C. G. and Dixon, F. J. (1962). Antibody production by transferred cells. Adv. Immunol., 2, 205-239.
- Cohen, J. J. and Claman, H. N. (1971). Thymus marrow immunocompetence. V. Hydrocortisone resistant cells and processes in the haemolytic antibody response of mice. J. expl Med., 133, 1026-1034.
- Cohen, J. J., Fischbach, M. and Claman, H. N. (1970). Hydrocortisone resistance of graft vs. host activity in mouse thymus, spleen and bone marrow. Immunology, 106, 1146-1150.
- Cohen, S. and Ward, P. A. (1971). In vitro and in vivo activity of a lymphocyte and immune complex - dependent chemotactic factor of eosinophils. J. exp. Med., 133, 133-146.
- Coker, C. M. (1954). Effects of cortisone on established immunity to Trichinella spiralis in mice, with reference to the role of the cellular factor(s) in the immunity mechanism. Ph.D. Thesis, University of North Carolina, U.S.A.
- Coker, C. M. (1955). Effects of cortisone on Trichinella spiralis infections in non-immunised mice. J. Parasit., 41, 498-504.
- Coker, C. M. (1956a). Some effects of cortisone in mice with acquired immunity to Trichinella spiralis. J. Inf. Dis., 98, 39-44.
- Coker, C. M. (1956b). Cellular factors in acquired immunity to Trichinella spiralis, as indicated by cortisone treatment of mice. J. Inf. Dis., 98, 187-197.
- Cornan, R. M. (1972). Passive protection with homologous antiserum against Trichostrongylus colubriformis in the guinea-pig. Immunology, 23, 647-650.

- Connan, R. M. (1974). The distribution of Nippostrongylus brasiliensis in prolonged infections in lactating and neonatally infected rats. Parasitology, 68, 347-354.
- Cookson, J. B., Montgomery, R. D., Morgan, H. V. and Tudor, R. W. (1972). Fatal paralytic ileus due to strongyloidiasis. Brit. Med. J., 4, 771-772.
- Cornwell, R. L. (1962). Blood eosinophil response of calves to normal and irradiated larvae of Dictyocaulus viviparus. J. Comp. Path., 72, 170-180.
- Crandall, R. B. and Crandall, C. A. (1972). Trichinella spiralis, immunologic response to infection in mice. Exptl Parasit., 31, 378-398.
- Craven, J. C., Cantrell, E. G. and Lewis, M. G. (1971). Strongyloides stercoralis infection presenting as necrotizing jejunitis. (Corresp.) Trans. R. Soc. trop. Med. Hyg., 65, 532-533.
- Cross, J. H. (1960). The natural resistance of the white rat to Nematospiroides dubius and the effect of cortisone on this resistance. J. Parasit., 46, 175-185.
- Cross, J. H. (1963). Nematospiroides dubius in the abnormal host. Ann. New York Acad. Sci., 113, 88-99.
- Cruz, T., Rebougas, G. and Rocha, H. (1966). Fatal strongyloidiasis in patients receiving corticosteroids. New Eng. J. Med., 275, 1093-1096.
- Culberston, J. T. (1942). Active immunity against Trichinella spiralis. J. Parasit., 28, 197-202.
- Culberston, J. T. (1943). Natural transmission of immunity against Trichinella spiralis from mother rats to their offspring. J. Parasit., 29, 114-116.
- Culberston, J. T. and Kaplan, S. S. (1937). Passive immunity in experimental trichinosis. J. Parasit., 23, 560.
- Culberston, J. T. and Kaplan, S. S. (1938). A study upon passive immunity in experimental trichinosis. Parasitology, 30, 156-166.
- Darling, S. T. (1911a). The intestinal worms of 300 insane patients detected by special methods. Bull. Soc. Path. Expt., 4, 334-341.
- Darling, S. T. (1911b). Strongyloides infections in man and animals in the Isthmian Canal Zone. J. Exp. Med., 14, 1-24.

- Connan, R. M. (1974). The distribution of Nippostrongylus brasiliensis in prolonged infections in lactating and neonatally infected rats. Parasitology, 68, 347-354.
- Cookson, J. B., Montgomery, R. D., Morgan, H. V. and Tudor, R. W. (1972). Fatal paralytic ileus due to strongyloidiasis. Brit. Med. J., 4, 771-772.
- Cornwell, R. L. (1962). Blood eosinophil response of calves to normal and irradiated larvae of Dictyocaulus viviparus. J. Comp. Path., 72, 170-180.
- Crandall, R. B. and Crandall, C. A. (1972). Trichinella spiralis, immunologic response to infection in mice. Exptl Parasit., 31, 378-398.
- Craven, J. C., Cantrell, E. G. and Lewis, M. G. (1971). Strongyloides stercoralis infection presenting as necrotizing jejunitis. (Corresp.) Trans. R. Soc. trop. Med. Hyg., 65, 532-533.
- Cross, J. H. (1960). The natural resistance of the white rat to Nematospiroides dubius and the effect of cortisone on this resistance. J. Parasit., 46, 175-185.
- Cross, J. H. (1963). Nematospiroides dubius in the abnormal host. Ann. New York Acad. Sci., 113, 88-99.
- Cruz, T., Rebougas, G. and Rocha, H. (1966). Fatal strongyloidiasis in patients receiving corticosteroids. New Eng. J. Med., 275, 1093-1096.
- Culberston, J. T. (1942). Active immunity against Trichinella spiralis. J. Parasit., 28, 197-202.
- Culberston, J. T. (1943). Natural transmission of immunity against Trichinella spiralis from mother rats to their offspring. J. Parasit., 29, 114-116.
- Culberston, J. T. and Kaplan, S. S. (1937). Passive immunity in experimental trichinosis. J. Parasit., 23, 560.
- Culberston, J. T. and Kaplan, S. S. (1938). A study upon passive immunity in experimental trichinosis. Parasitology, 30, 156-166.
- Darling, S. T. (1911a). The intestinal worms of 300 insane patients detected by special methods. Bull. Soc. Path. Expt., 4, 334-341.
- Darling, S. T. (1911b). Strongyloides infections in man and animals in the Isthmian Canal Zone. J. Exp. Med., 14, 1-24.

- Darrach, M. (1959). Effect of steroids on antibody production in vivo. In: The mechanisms of hypersensitivity (Shaffer, J. H., LoGrippo, G. A. and Chase, M. W., eds). Henry Ford Symposium, London, Churchill. pp. 613-622.
- De Campos, E. P. (1964). Chronic ulcerative colitis due to Strongyloides stercoralis. Revta Inst. Adolfo Lutz, 24, 51-56.
- De Harven, E. (1967). Methods in electron microscope cytology. In: Methods in Cancer Research. Vol. I (Busch, H., ed.). Academic Press, New York. pp. 3-44.
- Delaney and Garraty (1975). Handbook of Haematology and blood transfusion technique, 2nd edit. Butterworths.
- Denham, D. A. (1966a). Immunity to Trichinella spiralis. I. The immunity produced by mice to the first four days of the intestinal phase of the infection. Parasitology, 56, 323-327.
- Denham, D. A. (1966b). Immunity to Trichinella spiralis. II. Immunity produced by the adult worm in mice. Parasitology, 56, 745-751.
- Denham, D. A. (1968). Immunity to Trichinella spiralis. III. The longevity of the infection in mice. J. Helminth., 42, 257-268.
- Denham, D. A. (1969). Immunity to Trichinella spiralis. IV. Passive immunity in mice. Folia Parasitol., Praha, 16, 183-187.
- Denham, D. A. and Martinez, A. R. (1970). Studies with Methyridine and Trichinella spiralis. II. The use of the drug to study the rate of larval production in mice. J. Helminth., 44, 357-363.
- Denham, D. A., Pennudurai, T., Nelson, G. S., Rogers, R. and Guy, F. (1972). Studies with Brugia pahangi. 2. The effect of repeated infection on parasite levels in cats. Internat. J. Parasit., 2, 401-408.
- De Paolo, Dias, L. B., Da Silva, J. R. (1962). Enteritis due to Strongyloides stercoralis - A report of 5 fatal cases. Amer. J. Dig. Dis., 7, 1086.
- Despommier, D. D. (1971). Immunogenicity of the newborn larva of Trichinella spiralis. J. Parasit., 57, 531-535.
- Despommier, D. D. and Muller, M. (1970). Functional antigens of Trichinella spiralis. J. Parasit., 56, 76.
- Desportes, C. (1945). Sur Strongyloides stercoralis (Bavay, 1876) et sur le Strongyloides de primates. Ann. Parasit., 20, 160-190.

- Dineen, J. K. and Adams, D. B. (1971). The role of the recirculating thymus-dependent lymphocytes in resistance to Trichostrongylus colubriformis in the guinea-pig. Immunology, 20, 109-113.
- Dineen, J. K. and Kelly, J. D. (1973a). Immunological unresponsiveness of neonatal rats infected with Nippostrongylus brasiliensis. The competence of neonatal lymphoid cells in worm expulsion. Immunology, 25, 141-150.
- Dineen, J. K. and Kelly, J. D. (1973b). Expulsion of Nippostrongylus brasiliensis from the intestine of rats: the role of a cellular component derived from the bone marrow. Internat. Arch. Aller. Appl. Immunol., 45, 759-766.
- Dineen, J. K., Kelly, J. D. and Love, R. J. (1973a). The competence of lymphocytes obtained from immune and non-immune donors to cause expulsion of Nippostrongylus brasiliensis in the rat (DA strain). Internat. Arch. Aller. Appl. Immunol., 45, 504-512.
- Dineen, J. K., Ogilvie, B. M. and Kelly, J. D. (1973b). Expulsion of Nippostrongylus brasiliensis from the intestine of rats. Collaboration between humoral and cellular components of the immune response. Immunology, 24, 467-476.
- Dineen, J. K., Kelly, J. D., Goodrich, B. S. and Smith, I. D. (1974). Expulsion of Nippostrongylus brasiliensis from the small intestine of the rat by prostaglandin-like factors from ram semen. Internat. Arch. Aller. Appl. Immunol., 46, 360-371.
- Dineen, J. K. and Wagland, B. M. (1966). The cellular transfer of immunity to Trichostrongylus colubriformis in an isogenic strain of guinea pig. II. The relative susceptibility of the larval and adult stages of the parasite to immunological attack. Immunology, 11, 47-57.
- Dinetta, J., Katz, F. and Campbell, W. C. (1972). Effect of heterologous anti-lymphocyte serum on the spontaneous cure of Trichinella spiralis infections in mice. J. Parasit., 58, 636-637.
- Dorin, R. P. (1946). The preparation and demonstration of an anti-serum fro Trichinella spiralis. J. Parasit., 32, 83-86.
- Duckett, M. G. (1970). An investigation of the importance of the host strain in the host-parasite relationship of Trichinella spiralis and the mouse. M.Sc. dissertation, University of London.

- Duckett, M. G., Denham, D. A. and Nelson, G. S. (1972). Immunity to Trichinella spiralis. V. Transfer of immunity against the intestinal phase from mother to baby mice. J. Parasit., 58, 550-554.
- Edwards, A. J., Burt, J. S., and Ogilvie, B. M. (1971). The effect of immunity upon some enzymes of the parasitic nematode, Nippostrongylus brasiliensis. Parasitology, 62, 339-347.
- Enerbäck, L. (1966). Mast cells in rat gastro-intestinal mucosa. 2. Dye-binding and metachromatic properties. Acta Pathol. Microbiol. Scand., 66, 303-312.
- Fagundes, L. A., Busato, O. and Brentano, L. (1971). Strongyloidiasis: Fatal complication of renal transplantation. (Corresp.) Lancet, August, 439-440.
- Faubert, C. E. (1976). Depression of the plaque-forming cells to sheep red blood cells by the new-born larvae of Trichinella spiralis. Immunology, 30, 485-489.
- Faubert, G. M. and Tanner, C. E. (1975). Leucoagglutination and cytotoxicity of the serum of infected mice and of extracts of Trichinella spiralis larvae and the capacity of infected mouse serum to prolong skin allografts. Immunology, 28, 1041-1050.
- Faust, E. C. (1933). Experimental studies on human and primate species of Strongyloides. II. The development of Strongyloides in the experimental host. Amer. J. Hyg., 18, 114-132.
- Faust, E. C. and De Groat (1940). Internal auto-infection in human strongyloidiasis. Amer. J. trop. Med., 20, 350-375.
- Franz, K. H. (1963). Clinical trials with thiabendazole (MK-360) against human strongyloidiasis. Amer. J. trop. Med. Hyg., 12, 211-214.
- Frenckley, J. D. and Trewby, P. N. (1973). Fatal paralytic ileus due to strongyloidiasis. Br. Med. J., 2, 308.
- Fülleborn, F. (1914). Untersuchungen über den infektionsweg bei Strongyloides und Ankylostomum und die biologie dieser Parasiten. Arch. f. Schiffs- und Tropen-Hyg., Beih, 5, 26-80.
- Goldgraber, M. B. and Lawert, M. L. (1965). Immunological injury of mast cells and connective tissue in mice infected with Strongyloides ratti. J. Parasit., 51, 169-174.

- Gould, S. E., Gomberg, H. J., Bethell, F. H., Vilella, J. B. and Hertz, C. S. (1955). Studies on Trichinella spiralis. IV. Effect of feeding irradiated Trichinella larvae on the production of immunity to reinfection. Amer. J. Path., 31, 949-953.
- Graham, G. L. (1935). Single larva infections of Strongyloides ratti Sandground, 1925, as an approach to certain problems of Strongyloides bionomics. J. Parasit., 21 (Abstract), 432.
- Graham, G. L. (1936). Studies on Strongyloides. I. S. ratti in parasitic series, each generation in the rat established with a single homogenic larva. Amer. J. Hyg., 24, 71-87.
- Graham, G. L. (1938a). Studies on Strongyloides. II. Homogenic and heterogenic progeny of the single, homogenically derived S. ratti parasite. Amer. J. Hyg., 27, 221-234.
- Graham, G. L. (1938b). Studies on Strongyloides. III. The fecundity of single S. ratti of homogenic origin. J. Parasit., 24, 233-243.
- Graham, G. L. (1939a). Studies on Strongyloides. IV. Seasonal variation in the production of heterogenic progeny by singly established S. ratti from homogenically derived line. Amer. J. Hyg., 30, Sec. D, 15-27.
- Graham, G. L. (1939b). Studies on Strongyloides. V. Constitutional differences between a homogenic and a heterogenic line of S. ratti. J. Parasit., 25, 365-375.
- Graham, G. L. (1940). Studies on Strongyloides. VI. Comparison of two homogenic lines of singly established S. ratti. J. Parasit., 26, 207-218.
- Grassi, G. B. (1878). L'Anquillula intestinalis. Nota preventiva. Gazz. Med. Ital. Lomb., 5, 471-474.
- Grassi, G. B. (1879). Soura l'Anquillula intestinale. Revd. Inst. Lomb. di Sc. e Lett., Rendic, Milano, 12, 228-233.
- Grassi, G. B. and Segre, R. (1887). Nuova osservazione sull'eterogenia del Rhabdonema (Anquillula) intestinale. Reviewed by Braun. Centr. Bakt. u. Parasit., 2, 413-415.
- Griffin, L. S. P. (1972). The effect of immunosuppressants on Strongyloides ratti infection. M.Sc. project, University of London.
- Gudson, J. P. Jr., Moore, V. L., Myrvik, O. H. and Holyfield, P. A. (1972). Promethazine hydrochloride as an immunosuppressant. J. Immunol., 108, 1340-1344.

- Gyllenstein, L. (1962). The cellular composition of lymphatic tissue during involution after administration of corticosteroid. Acta Path. Microbiol. Scand., 56, 35-45.
- Hall, M. C. (1916). Nematode parasites of mammals of the order Rodentia, Lagomorpha and Hyracoidea. Proc. U.S. Nat. Mus., 50, 1-284.
- Harada, Y. and Mori, O. (1952). Diagnosis of Ancylostomiasis with cultivation method and skin test. Rinshyo Igaku, 39, 219-223.
- Harley, J. P. and Gallicchio, V. (1970). Effect of cortisone on the establishment of Nippostrongylus brasiliensis in the rabbit. J. Parasit., 56, 271-276.
- Harness, E., Sellwood, S. A. and Young, E. R. (1971). Experimental Haemonchus placei infection in calves. Influence of anaemia and numbers of larvae on worm development. J. Comp. Path., 81, 129-136.
- Hartung, W. and Becht, H. (1960). Über die Wirkung des Betastrahlen des P³² auf Trichinella spiralis. Strahlentherapie, 112, 613-620.
- Hellström, K. E. and Hellström, I. (1970). Immunological enhancement as studied by cell culture technique. Annual R. Microbiol., 24, 373.
- Hellström, I., Evans, C. A. and Hellström, K. E. (1969). Cellular immunity and its serum-mediated inhibition in Shope-virus-induced rabbit papillomas. Internat. J. Cancer, 4, 601-607.
- Hellström, I. and Hellström, K. E. (1969). Studies on cellular immunity and its serum-mediated inhibition in Moloney-virus-induced mouse sarcomas. Internat. J. Cancer, 4, 587-600.
- Hendricks, J. R. (1950). The relationship between precipitin titre and number of Trichinella spiralis in the intestinal tract of mice following test infections. J. Immunol., 64, 173-177.
- Hendricks, J. R. (1952). Studies in mice on the dual antibody basis of acquired resistance to Trichinella spiralis. J. Elisha Mitchell Sci. Soc., 68, 12-36.
- Hendricks, J. R. (1953). Passive transfer of resistance to Trichinella spiralis in mice. J. Parasit., 39, 40-41.
- Henney, C. S., Bourne, H. R. and Lichtenstein, L. M. (1972). The role of cyclic 3'5' adenosine monophosphate in the specific cytolytic activity of lymphocytes. J. Immunol., 108, 1526-1534.

- Henney, R. W., Maclean, J. M. and Mulligan, W. (1971). The effect of host immunity on the metabolism of Nippostrongylus brasiliensis. Immunology, 21, 711-718.
- Henry, A. C. L. (1931). Les parasites et les maladies parasitaires du ragondin. Bull. Soc. Nat. Acclimat. France, 78, 421-447.
- Herlich, H. (1963). Resistance to Trichostrongylus colubriformis in the guinea-pig. J. Parasit., 49 (Suppl.), 33.
- Herlich, H., Douvres, F. W. and Isenstein, R. S. (1956). Experimental infections of guinea pigs with Trichostrongylus colubriformis, a parasite of ruminants. Proc. Helminth. Soc. Wash., 23, 104-105.
- Hogarth-Scott, R. S. and Bingley, J. B. (1971). Prolongation and enhancement of Nippostrongylus infection in the laboratory rat by a heterologous antiserum to rat peritoneal cells - A possible role for pharmacologically-active cells in immunity. Immunology, 21, 87-99.
- Hopkins, C. A., Subramanian, G. and Stallard, H. (1972). The effect of immunosuppressants on the development of Hymenolepis diminuta in mice. Parasitology, 65, 111-120.
- Huggins, D. (1971a). Severe strongyloidiasis: A case report. Revta Soc. Bras. Med. Trop., 5, 159-174.
- Huggins, D. (1971b). Severe infection in strongyloidiasis, report of a case. Anais. Escola Nac. Saude Publ. Med. Trop., 5, 271-282.
- Humphrey, J. H. and White, R. G. (1970). Immunology for students of medicine. Blackwell Scientific Publications, Oxford. (2nd ed.)
- Hunter, G. C. and Leigh, L. C. (1961). Studies on the resistance of rats to the nematode Nippostrongylus muris (Yokogawa, 1922). III. Attempt to transfer immunity adoptively. Parasitology, 51, 357-366.
- James, E. R. and Danham, D. A. (1975). Immunity to Trichinella spiralis. VI. The specificity of the immune response stimulated by the intestinal stage. J. Helminth., 49, 43-47.
- Jannar, M. P. (1966). Cytochemical and electron microscope studies on the pharynx and intestinal epithelium of Nippostrongylus brasiliensis. J. Parasit., 52, 1116.
- Jarrett, W. F. H., Jennings, F. W., McIntyre, W. I. M., Mulligan, W. and Urquhart, G. M. (1960). Immunological studies on Dictyocaulus viviparus infection. Immunity produced by the administration of irradiated larvae. Immunology, 3, 145-151.

- Henney, R. W., Maclean, J. M. and Mulligan, W. (1971). The effect of host immunity on the metabolism of Nippostrongylus brasiliensis. Immunology, 21, 711-718.
- Henry, A. C. L. (1931). Les parasites et les maladies parasitaires du ragondin. Bull. Soc. Nat. Acclimat. France, 78, 421-447.
- Herlich, H. (1963). Resistance to Trichostrongylus colubriformis in the guinea-pig. J. Parasit., 49 (Suppl.), 33.
- Herlich, H., Douvres, F. W. and Isenstein, R. S. (1956). Experimental infections of guinea pigs with Trichostrongylus colubriformis, a parasite of ruminants. Proc. Helminth. Soc. Wash., 23, 104-105.
- Hogarth-Scott, R. S. and Bingley, J. B. (1971). Prolongation and enhancement of Nippostrongylus infection in the laboratory rat by a heterologous antiserum to rat peritoneal cells - A possible role for pharmacologically-active cells in immunity. Immunology, 21, 87-99.
- Hopkins, C. A., Subramanian, G. and Stallard, H. (1972). The effect of immunosuppressants on the development of Hymenolepis diminuta in mice. Parasitology, 65, 111-120.
- Huggins, D. (1971a). Severe strongyloidiasis: A case report. Revta Soc. Bras. Med. Trop., 5, 159-174.
- Huggins, D. (1971b). Severe infection in strongyloidiasis, report of a case. Anais. Escola Nac. Saude Publ. Med. Trop., 5, 271-282.
- Humphrey, J. H. and White, R. G. (1970). Immunology for students of medicine. Blackwell Scientific Publications, Oxford. (2nd ed.)
- Hunter, G. C. and Leigh, L. C. (1961). Studies on the resistance of rats to the nematode Nippostrongylus muris (Yokogawa, 1922). III. Attempt to transfer immunity adoptively. Parasitology, 51, 357-366.
- James, E. R. and Denham, D. A. (1975). Immunity to Trichinella spiralis. VI. The specificity of the immune response stimulated by the intestinal stage. J. Helminth., 49, 43-47.
- Jenaur, M. P. (1966). Cytochemical and electron microscope studies on the pharynx and intestinal epithelium of Nippostrongylus brasiliensis. J. Parasit., 52, 1116.
- Jarrett, W. F. H., Jennings, F. W., McIntyre, W. I. M., Mulligan, W. and Urquhart, G. M. (1960). Immunological studies on Dictyocaulus viviparus infection. Immunity produced by the administration of irradiated larvae. Immunology, 3, 145-151.

- Jarrett, W. F. H., Jarrett, E. E. E., Miller, H. R. P. and Urquhart, G. M. (1968). Quantitative studies on the mechanism of self-cure in Nippostrongylus brasiliensis infections. Proc. 3rd. Internat. Congr. Wld Ass. Adv. Vet. Parasit. In the: Reaction of the host to parasitism (Ed. Soulsby, Elwert, Marburg/L). pp. 191-198.
- Jarrett, E. E. E. and Urquhart, G. M. (1971). The immune response to nematode infections. Internat. Rev. Trop. Med., 4, 53-96.
- Jennings, F. W., Mulligan, W. and Urquhart, G. M. (1963). Variables in X-ray "inactivation" of Nippostrongylus brasiliensis larvae. Expl Parasit., 13, 367-373.
- Jones, V. E., Edwards, A. J. and Ogilvie, B. M. (1970). The circulating immunoglobulins involved in protective immunity to the intestinal stage of N. brasiliensis in the rat. Immunology, 18, 621-633.
- Jones, V. E. and Ogilvie, B. M. (1971). Protective immunity to Nippostrongylus brasiliensis: the sequence of events which expels worms from the rats intestine. Immunology, 20, 549-561.
- Jones, V. E. and Ogilvie, B. M. (1972). Protective immunity to Nippostrongylus brasiliensis in the rat. III. Modulation of worm acetylcholinesterase by antibodies. Immunology, 22, 119-129.
- Jones, W. O., Rothwell, T. L. W., Dineen, J. K. and Griffiths, D. A. (1974). Studies on the role of histamine and 5-hydroxytryptamine in immunity against the nematode Trichostrongylus colubriformis. II. Amine levels in the intestine of infected guinea pigs. Internat. Arch. Allerg. Appl. Immunol., 46, 14-27.
- Jorge, J. and Barbosa, W. (1964). Treatment of strongyloidiasis with Thiabendazole. Rev. Goiana Med., 10, 135-140.
- Katz, F. F. (1961). Distribution of the nematode Strongyloides ratti in the small intestine of gonadectomized and sham-operated male and female rats. Amer. Zool., 1, 363-364.
- Katz, F. F. (1962). Differences in Strongyloides ratti worm burdens in male and female rats. J. Parasit., 48 (Suppl.), 51.
- Katz, F. F. (1963). Testosterone and estrogen treatments of gonadectomized rats in experiments on the host sex difference in S. ratti intestinal burdens. J. Parasit., 49 (Suppl.), 52.
- Katz, F. F. (1967). On a host sex difference in Strongyloides ratti intestinal worm burdens in rats. Proc. Penn. Acad. Sci., 41, 30-37.

- Kay, A. B. (1974). The eosinophil in infectious diseases. J. Inf. Dis., 129, 606-613.
- Kazacos, K. R. (1976). Increased resistance in the rat to Strongyloides ratti following immunization against Trichinella spiralis. J. Parasit., 62, 493-494.
- Kazacos, K. R. and Thomson, R. E. (1975). Cross-resistance between Nippostrongylus brasiliensis and Strongyloides ratti in rats. J. Parasit., 61, 525-529.
- Keller, R. (1971). Nippostrongylus brasiliensis in the rat: failure to relate intestinal histamine and mast cell levels with worm expulsion. Parasitology, 63, 473-481.
- Keller, R. and Keist, R. (1972). Protective immunity to Nippostrongylus brasiliensis in the rat: central role of the lymphocyte in worm expulsion. Immunology, 22, 767-773.
- Keller, R. and Ogilvie, B. M. (1972). The effects of drugs on worm expulsion in Nippostrongylus brasiliensis infected rats: a discussion of the interpretation of drug action. Parasitology, 64, 217-227.
- Kelly, J. D. (1972). Immunological suppression of parasite rejection by removal of selected lymphocyte populations, the effect of neonatal thymectomy and anti-lymphocyte serum administration on the resistance of rats to infection with the nematode Nippostrongylus brasiliensis. Aust. J. exp. Biol. Med. Sci., 50, 477-489.
- Kelly, J. D. and Dineen, J. K. (1972). The suppression of rejection of Nippostrongylus brasiliensis in the rat by promethazine hydrochloride. The site of immunological impairment. Immunology, 22, 361-370.
- Kelly, J. D. and Ogilvie, B. M. (1972). Intestinal mast cells and eosinophil numbers during worm expulsion in nulliparous and lactating rats infected with Nippostrongylus brasiliensis. Internat. Arch. Aller. Appl. Immunol., 43, 497-509.
- Kelly, J. D., Dineen, J. K. and Love, R. J. (1973). Expulsion of Nippostrongylus brasiliensis from the intestine of rats: evidence for a third component in the rejection mechanism. Internat. Arch. Aller. Appl. Immunol., 45, 767-779.
- Kelly, J. D., Dineen, J. K., Goodrich, B. S. and Smith, I. D. (1974). Expulsion of Nippostrongylus brasiliensis from the intestine of rats. Role of prostaglandins and pharmacologically active amines (histamine and 5-hydroxytryptamine) in worm expulsion. Internat. Arch. Aller. Appl. Immunol., 47, 458-465.

- Kim, C. W. (1957). Immunity to Trichinella spiralis in mice infected with irradiated larvae. J. Elisha Mitchell Sci. Soc., 73, 308-317.
- Kreis, H. (1932). Studies on the genus Strongyloides (Nematodes). Amer. J. Hyg., 16, 450-491.
- Larsh, J. E. Jr. (1967). The present understanding of the mechanism of immunity to Trichinella spiralis. Amer. J. trop. Med. Hyg., 16, 123-132.
- Larsh, J. E. Jr. and Hendricks, J. R. (1949). The probable explanation for the difference in the localisation of adult Trichinella spiralis in young and old mice. J. Parasit., 35, 101-106.
- Larsh, J. E. Jr., Gilchrist, H. B. and Greenberg, B. G. (1952). A study of the distribution and longevity of adult Trichinella spiralis in immunised and non-immunised mice. J. Elisha Mitchell Sci. Soc., 68, 1-11.
- Larsh, J. E. Jr., Goulson, H. T. and Weatherly, N. F. (1964a). Studies on delayed (cellular) hypersensitivity in mice infected with Trichinella spiralis. I. Transfer of lymph node cells. J. Elisha Mitchell Sci. Soc., 80, 133-135.
- Larsh, J. E. Jr., Goulson, H. T. and Weatherly, N. F. (1964b). Studies on delayed (cellular) hypersensitivity in mice infected with Trichinella spiralis. II. Transfer of peritoneal exudate cells. J. Parasit., 50, 496-498.
- Larsh, J. E. Jr., Goulson, H. T., Weatherly, N. F. and Chaffee, E. F. (1970). Studies on delayed (cellular) hypersensitivity in mice infected with Trichinella spiralis. VI. Results in recipients injected with antiserum or "freeze-thaw" spleen cells. J. Parasit., 56, 1206-1209.
- Larsh, J. E. Jr. and Race, G. J. (1954). A histopathologic study of the anterior small intestine of immunized and non-immunized mice infected with Trichinella spiralis. J. Inf. Dis., 94, 262-272.
- Larsh, J. E. Jr. and Race, G. J. (1975). Allergic inflammation as a hypothesis for the expulsion of worms from tissues. A review. Exptl Parasit., 37, 251-266.
- Larsh, J. E. Jr., Race, G. J. and Jeffries, W. B. (1956). The association in young mice of intestinal inflammation and the loss of adult worms following an initial infection with Trichinella spiralis. J. Inf. Dis., 99, 63-71.

- Lawler, H. J. (1940). Passive transfer of immunity to the nematode Strongyloides ratti. Amer. J. Hyg., 31, 28-31.
- Lee, C. L. and Lewert, R. M. (1957). Studies on the presence of muco-polysaccharidase in penetrating helminth larvae. J. Inf. Dis., 101, 287-294.
- Lee, D. L. (1965). The cuticle of adult Nippostrongylus brasiliensis. Parasitology, 55, 173.
- Lee, D. L. (1969a). Changes in adult Nippostrongylus brasiliensis during the development of immunity to this nematode in rats. 1. Changes in ultrastructure. Parasitology, 59, 29-39.
- Lee, D. L. (1969b). Nippostrongylus brasiliensis: some aspects of the fine structure and biology of the infective larva and the adult. Symp. Bri. Soc. Parasitol., 59, 29-39.
- Lee, D. L. (1970). The fine structure of the excretory system in adult Nippostrongylus brasiliensis (Nematoda) and a suggested function for the "excretory glands". Tissue and Cell, 2, 225-231.
- Leichtenstem, O. (1899). Zur lebensgeschichte der Anguillula intestinalis. Zbl. Bakt., 2, 226-231.
- Lendrum, A. C. (1944). The staining of eosinophil polymorphs and enterochromaffin cells in histological sections. J. Path. Bact., 56, 441.
- Leuckart, R. (1882). Ueber die Lebensgeschichte der sogenannten A. stercoralis und deren Beziehungen zu der Sogenannten A. intestinalis. Bericht. Verhandl. sachs. Gesellsch. Wissensch. Math. - Phys. Klasse, Leipzig., 34, 85-107.
- Levin, A. J. and Evans, T. C. (1942). The use of roentgen radiation in locating an origin of host resistance to Trichinella spiralis infections. J. Parasit., 28, 477-483.
- Litt, M. (1964). Studies in experimental eosinophilia. VI. Uptake of immune complexes by eosinophils. J. Cell Biol., 23, 355-361.
- Little, M. D. (1961). Concurrent infections of Strongyloides venezuelensis and S. ratti in wild rats in New Orleans. J. Parasit., 44 (Suppl.), 17.
- Little, M. D. (1966a). Comparative morphology of six species of Strongyloides (Nematoda) and redefinition of the genus. J. Parasit., 52, 69-84.
- Little, M. D. (1966b). Seven new species of Strongyloides (Nematoda) from Louisiana. J. Parasit., 52, 85-97.

- Ljungström, I. and Ruitenberg, E. J. (1976). A comparative study of the immunohistological and serological response of intact and T. cell-deprived mice to Trichinella spiralis. Clin. Exp. Immunol., 24, 146-156.
- Looss, A. (1904). Die Wanderung der Ancylostomum und Strongyloides larven von der Haut nach dem Darm. C.r. 6me. Congres. Internat. de Zoologie, Berne, 225-233.
- Looss, A. (1911). The anatomy and life history of Ancylostoma duodenale Dub. A monograph. Part II. The development in the free state. Rec. Egypt Gov. Schl. Med., 4, 159-613.
- Louch, C. D. (1962). Increased resistance to Trichinella spiralis in the laboratory rat following infection with Nippostrongylus muris. J. Parasit., 48, 24-26.
- Louisy, C. L. and Barton, C. J. (1971). The radiological diagnosis of Strongyloides stercoralis enteritis. Radiology, 98, 535-541.
- Love, R. J. (1975). Nippostrongylus brasiliensis infections in rats. Both antibodies and sensitized cells are necessary for the immunological control of developing larvae. Internat. Arch. Aller. Appl. Immunol., 48, 211-219.
- Love, R. J., Kelly, J. D. and Dineen, J. K. (1974). Nippostrongylus brasiliensis: effects of immunity on the pre-intestinal and intestinal larval stages of the parasite. Internat. J. Parasit., 4, 183-191.
- Love, R. J., Ogilvie, B. M. and McLaren, D. J. (1976). The immune mechanism which expels the intestinal stages of Trichinella spiralis from rats. Immunology, 30, 7-15.
- Lukashenko, N. P. (1960). Microprecipitation on larvae of Trichinella spiralis in vitro as a method of vital diagnostics of trichinellosis in swine. Helminthologia, 2, 105-110.
- Markell, E. K. (1958). The effect of cortisone treatment upon the longevity and productivity of Trichinella spiralis in the rat. J. Inf. Dis., 102, 158-161.
- Markell, E. K. and Lewis, W. P. (1957). Effect of cortisone treatment on immunity to subsequent reinfections with Trichinella in the rat. Amer. J. trop. Med. Hyg., 6, 553-561.
- Marks, C. F., Thomson, I. J. and Castro, C. E. (1968). Dynamics of the permeation of nematodes by water, nematocides and other substances. Exptl Parasit., 22, 321-337.

- Medawar, P. (1969). Immunosuppressive agents with special reference to antilymphocytic serum. Proc. Roy. Soc. B., 174, 155-172.
- McCoy, O. R. (1931). Immunity of rats to reinfection with Trichinella spiralis. Amer. J. Hyg., 14, 484-494.
- McCoy, O. R. (1940). Rapid loss of Trichinella fed to immune rats and its bearing on the mechanism of immunity. Amer. J. Hyg., 32, 105-116.
- McCoy, O. R. and Bond, F. F. (1941). Attempted passive transfer to rats and mice of immunity to Trichinella spiralis. J. Parasit., 27, 24-25.
- McLaren, D. J., Burt, J. S. and Ogilvie, B. M. (1974). The anterior glands of adult Necator americanus (Nematoda : Strongyloidea) II. Cytochemical and functional studies. Internat. J. Parasit., 5, 39-46.
- Michel, J. F., Lancaster, H. B. and Hong, C. (1972). Host induced effects on the vulval flap of Ostertagia ostertagi. Internat. J. Parasit., 2, 305-318.
- Miller, T. A. (1964). Effect of X-irradiation upon the infective larvae of Ancylostoma caninum and the immunogenic effect in dogs of a single infection with 40 Kr irradiated larvae. J. Parasit., 50, 735-742.
- Miller, T. A. (1966). Comparison of the immunogenic efficiencies of normal and X-irradiated Ancylostoma caninum larvae in dogs. J. Parasit., 52, 512-519.
- Millonig, G. (1961). Advantages of a phosphate buffer for osmium tetroxide solutions in fixation. J. Appl. Physiol., 32, 1637.
- Mills, C. K. and Kent, N. H. (1965). Excretions and secretions of Trichinella spiralis and their role in immunity. Expl Parasit., 16, 300-310.
- Moss, G. D. (1972). The effect of cortisone acetate treatment on the growth of Hymenolepis microstoma in mice. Parasitology, 64, 311-320.
- Mougeot, G. and Lancaster, F. (1973). Trichinose experimentale (3^e note) - Transplantations intestinales des Trichinella spiralis adultes. Ann. Parasit. Hum. Comp., 48, 55-59.
- Mulligan, W., Urquhart, G. M., Jennings, F. W. and Neilson, J. T. M. (1965). Immunological studies on Nippostrongylus brasiliensis infection in the rat. The "self-cure" phenomenon. Expl Parasit., 16, 341-347.

- Murray, M., Jarrett, W. F. H. and Jennings, F. W. (1971a). Mast cells and macromolecular leak in intestinal immunological reaction. The influence of sex of rats infected with Nippostrongylus brasiliensis. Immunology, 21, 17-31.
- Murray, M., Miller, H. R. P., Sanford, J. and Jarrett, W. F. H. (1971b). 5-hydroxytryptamine in intestinal immunological reaction. Its relationship to mast cell activity and worm expulsion. Internat. Arch. Aller. Appl. Immunol., 40, 236-247.
- Murray, M., Smith, W. D., Waddell, A. H. and Jarrett, W. F. H. (1971c). Nippostrongylus brasiliensis. Histamine and 5-hydroxytryptamine inhibition and worm expulsion. Exptl Parasit., 30, 58-63.
- Nashimura, H. (1943). Helminths of alimentary tract of house rats in Fukuoka city. Fukuoka Ika Daigaku Zasshi, 38, 726-741.
- Neefe, L. I., Pinilla, O., Garagusi, V. F. and Bauer, H. (1973). Disseminated strongyloidiasis with cerebral involvement: A complication of corticosteroid therapy. Amer. J. Med., 55, 832-838.
- Nolasco, J. O. and Africa, C. M. (1936). A fatal case of paralytic ileus associated with severe Strongyloides infection suggesting internal auto-infection. J. Philipp. Isl. Med. Ass., 16, 275-283.
- Nordstrom, C., Dahlquist, A. and Josefsson, L. (1968). Quantitative determination of enzymes in different parts of the villi and crypts of rat small intestine. Comparison of alkaline phosphatase, disaccharidases and dipeptidases. J. Histochem. Cytochem., 15, 713.
- North, R. J. (1972). The action of cortisone on cell-mediated immunity to infection: histogenesis of the lymphoid cell response and selective elimination of committed lymphocytes. Cell Immunol., 3, 501-505.
- Noto-Soediro, R. (1933). Een Strongyloides - soort bij een haan. Nederl. - Ind. Blad. Diergeneesk., 45, 547-550.
- Ogilvie, B. M. (1964). Nippostrongylus brasiliensis: a study of the life cycle and the immunological response of the host. Ph.D. Thesis, University of Cambridge, England.
- Ogilvie, B. M. (1965a). Role of adult worms in immunity of rats to Nippostrongylus brasiliensis. Parasitology, 55, 325-335.

- Ogilvie, B. M. (1965b). Use of cortisone derivatives to inhibit resistance to Nippostrongylus brasiliensis and to study the fate of parasites in resistant hosts. Parasitology, 55, 723-730.
- Ogilvie, B. M. (1967). Reagin-like antibodies in rats infected with the nematode parasite Nippostrongylus brasiliensis. Immunology, 12, 113-131.
- Ogilvie, B. M. (1969). Immunity to Nippostrongylus brasiliensis. Symp. Bri. Soc. Parasit., 7, 31-40.
- Ogilvie, B. M. (1972). Protective immunity to Nippostrongylus brasiliensis in the rat. II. Adaptation by worms. Immunology, 22, 111-118.
- Ogilvie, B. M. and Hockley, D. J. (1968). Effect of immunity on Nippostrongylus brasiliensis adult worms: reversible and irreversible changes in infectivity, reproduction and morphology. J. Parasit., 54, 1073.
- Ogilvie, B. M., and Jones, V. E. (1967). Reaginic antibodies and immunity to Nippostrongylus brasiliensis in the rat. I. The effect of thymectomy, neonatal infections and splenectomy. Parasitology, 57, 335-349.
- Ogilvie, B. M. and Jones, V. E. (1968). Passive protection with cells and antiserum against Nippostrongylus brasiliensis in the rat. Parasitology, 58, 939-949.
- Ogilvie, B. M. and Jones, V. E. (1969). Reaginic antibodies and helminth infections, In: Movat's "Cellular and humoral mechanisms in anaphylaxis and allergy". Karger, Basel. pp. 13-22.
- Ogilvie, B. M. and Jones, V. E. (1971). Nippostrongylus brasiliensis: A review of immunity and the host/parasite relationship in the rat. Exptl Parasit., 29, 138-177.
- Ogilvie, B. M. and Jones, V. E. (1973). Immunity in the parasitic relationship between helminths and hosts. Progr. Allergy, 17, 93-144.
- Ogilvie, B. M. and Love, R. J. (1974). Cooperation between antibodies and cells in immunity to a nematode parasite. Transplant. Rev., 19, 147-168.
- Ogilvie, B. M. and Wilson, R. J. M. (1976). Evasion of the immune response by parasites. Bri. Med. Bull., 32, 177-181.

- Ogilvie, B. M. and Worms, M. J. (1976). Immunity to Nematode Parasite of man with special reference to Ascaris, hookworms and filariae. In: Immunology of parasitic infections (Cohen, S. and Sadun, E. H., eds.). Blackwell Scientific Publications. pp. 380-407.
- Ogilvie, B. M., Rothwell, T. L. W., Bremner, K. C., Schnitzerling, H. J., Nolan, J. and Keith, R. K. (1973). Acetylcholinesterase secretion by parasitic nematodes. I. Evidence for secretion of the enzyme by a number of species. Internat. J. Parasit., 3, 589-597.
- Ohira, T. (1918). Problem whether human Strongyloides is infective on the other animals, in addition to auto-infection. Tokyo Iji Shinshi, 2096, 2003-2009.
- Olivier, L. (1962). Studies on natural resistance to Taenia taeniaeformis in mice. II. The effect of cortisone. J. Parasit., 48, 758-762.
- Oliver-Gonzalez, J. (1940). The dual antibody basis of acquired immunity in trichinosis. J. Inf. Dis., 69, 254-270.
- Oliver-Gonzalez, J. (1940). The in vitro action of immune serum on the larvae and adults of Trichinella spiralis. J. Inf. Dis., 67, 292-300.
- Olurin, E. O. (1970). Strongyloidiasis causing fatal peritonitis. W. African Med. J., 19, 102-104.
- Opie, E. (1904). An experimental study of the relation of cells with eosinophile granulation to infection with an animal parasite (Trichinella spiralis). Amer. J. Med. Sci., 127, 477-493.
- Pampiglione, S. and Ricciardi, M. L. (1972a). Experimental infestation with human strain S. fülleborni in man. Lancet, March 25, 663-665.
- Pampiglione, S. and Ricciardi, M. L. (1972b). Geographic distribution of Strongyloides fülleborni in humans in tropical Africa. Parassitologia, 14, 329-338.
- Pampiglione, S. and Ricciardi, M. L. (1971a). The presence of Strongyloides fülleborni (Von Linstow, 1905) in man in central and east Africa. Parassitologia, 13, 257-269.
- Pampiglione, S. and Ricciardi, M. L. (1971b). Experimental human infection with S. fülleborni of human origin. Parassitologia, 13, 505-511.

- Perroncito, E. (1880). Osservazione elmintologiche relative alla malattia sull'upatasi endemica negli operai del Gottardo. Atti R. Accad. Lincei, Roma, 1, 381-433.
- Perry, R. H. (1974). Transfer of immunity to Trichinella spiralis from mother to offspring. J. Parasit., 60, 460-465.
- Persaud, C. R. and Grant, L. S. (1972). In vitro comparison of the activity of Mintzol (Thiabendazole) and Ketrax (Levamisole) against S. stercoralis. (Corresp.) Trans. R. Soc. trop. Med. Hyg., 66, 368-369.
- Piper, P. J. and Vane, J. R. (1969). The release of prostaglandins during anaphylaxis in guinea-pig isolated lung; in Mantegazza and Horton (eds.) Prostaglandins, peptides and amines. Academic Press, New York. pp. 15-19.
- Podhajecy, K. (1962). Localisation of intestinal trichinellae in the small intestine of mice in their intestinal phase. Wiadomosci Parazyto'ogiczne, 8, 633-636.
- Poltra, A. A. (1972). Pulmonary strongyloidiasis. (Corresp.) Trans. R. Soc. trop. Med. Hyg., 66, 520.
- Purtilo, D. T., Meyers, W. M. and Connor, D. H. (1974). Fatal strongyloidiasis in immunosuppressed patients. Amer. J. Med., 56, 488-493.
- Raff, M. C. (1970). Two distinct populations of peripheral lymphocytes in mice distinguishable by immunofluorescence. Immunology, 19, 637-650.
- Ransom, B. H. (1907). Observations on the life history of Strongyloides longur. Bureau Ani. Ind., Circ. 116, 4-6, Washington.
- Ransom, B. H. (1911). The nematodes parasitic in the alimentary tract of cattle, sheep and other ruminants. Bureau Ani. Ind., Bull. 127, 132, Washington.
- Rappaport, I. and Wells, H. S. (1951). Studies in trichinosis. I. Immunity to reinfection in mice following a single light infection. J. Inf. Dis., 88, 248-253.
- Rago, A. A. and Kahtalian, A. (1972). Attempts to immunize rats using irradiated larvae of Strongyloides ratti. Rev. Brasileira de Biol., 32, 97-99.
- Reiff, O. F. and De Moraes, C. R. (1972). Strongyloidiasis: report of two fatal cases presenting an acute abdomen. Revista Paulista de Medicina, 80, 151-154.

- Reynolds, D. G., Brim, J. and Sheehy, T. W. (1967). The vascular architecture of the small intestinal mucosa of the monkey (Macaca mulatta). Anat. Rec., 159, 211.
- Rickard, M. D. (1974). Hypothesis for the long term survival of Taenia pisiformis cysticerci in rabbits. Z. Parasitenk., 44, 203-209.
- Rickard, M. D. and Outteridge, P. M. (1974). Antibody and cell-mediated immunity in rabbits infected with the larval stages of Taenia pisiformis. Z. Parasitenk., 44, 187-201.
- Riley, J. F. (1959). The mast cells. E. & S. Livingstone, Edinburgh.
- Rivera, E., Madonado, N., Velez-Garcia, Grillo, A. J. and Malaret, G. (1970). Hyperinfection syndrome with Strongyloides stercoralis. Ann. Int. Med., 72, 199-204.
- Roberts, F. H. S. and Keith, R. K. (1959a). The efficiency of Phenothiazine against the immature stages of the bovine stomach worm, Haemonchus placei (Place, 1893) Ransom 1911. Aust. Vet. J., 35, 38-39.
- Roberts, F. H. S. and Keith, R. K. (1959b). Observations on the effect of treatment with Phenothiazine on the development of resistance by calves to infestation with the stomach worm Haemonchus placei (Place, 1893) Ransom 1911. Aust. Vet. J., 35, 409-414.
- Roberts, F. H. S., Riek, R. F. and Keith, R. K. (1963). Studies on resistance in calves to experimental infection with the nodular worm Oesophagostomum radiatum (Rudolphi, 1803). II. The role of the respective stages of the parasitic life cycle in the stimulation of resistance. Aust. J. Agric. Res., 14, 704-715.
- Rogers, W. A. Jr. and Nelson, B. (1966). Strongyloidiasis and malignant lymphoma, "opportunistic infection" by a nematode. J. Amer. Med. Ass., 195, 685-687.
- Roman, E. (1951). Etude ecologique et morphologique sur les acanthocephales et la nematodes parasites des rats de la region Lyonnaise. Mém. Mus. Nat. d'Hist. Nat., Ser. A.2, 49-270.
- Roman, E. (1956). Specificite parasitaire de Strongyloides ratti du surmulot. Effets de la cortisone sur l'infestation d'autres rongeurs par le nematode. Ann. Parasit. Hum. Comp., 31, 552-571.

- Roth, H. (1939). Experimental studies on the course of trichina infection in guinea-pigs. III. Immunity of guinea-pigs to reinfection with Trichinella spiralis. Amer. J. Hyg., 30, 35-64.
- Rothwell, T. L. W. and Dineen, J. K. (1973). The response of the regional lymph node of guinea-pigs to primary and challenge infection with the nematode Trichostrongylus colubriformis. Internat. J. Parasit., 3, 201-208.
- Rothwell, T. L. W., Dineen, J. K. and Love, R. J. (1971). The role of pharmacologically active amines in resistance to Trichostrongylus colubriformis in the guinea-pig. Immunology, 21, 925-938.
- Rothwell, T. L. W., Ogilvie, B. M. and Love, R. J. (1973). Acetylcholinesterase secretion by parasitic nematodes. II. Trichostrongylus spp. Internat. J. Parasit., 3, 599-608.
- Rothwell, T. L. W., Prichard, R. K. and Love, R. J. (1974a). Studies on the role of histamine and 5-hydroxytryptamine in immunity against the nematode Trichostrongylus colubriformis. I. In vivo and in vitro effects of the amines. Internat. Arch. Aller. Appl. Immunol., 46, 1-13.
- Rothwell, T. L. W., Jones, W. O. and Love, R. J. (1974b). Studies on the role of histamine and 5-hydroxytryptamine in immunity against the nematode Trichostrongylus colubriformis. III. Inhibition of worm expulsion from guinea pigs by treatment with Reserpine. Internat. Arch. Aller. Appl. Immunol., 47, 875-886.
- Rovelli, C. (1888). Ricerche sugli organi degli Strongyloides Como. Reviewed by Brau, M. Centralbl. f. Bakt. und Parasitenk., 4, 660-661.
- Ruitenberg, E. J. and Steerenberg, P. A. (1974). Intestinal phase of Trichinella spiralis in congenitally athymic (nude) mice. J. Parasit., 60, 1056-1057.
- Sabesin, S. M. (1963). A function of the eosinophil: phagocytosis of antigen-antibody complexes. Proc. Soc. Exptl. Biol. Med., 112, 667-670.
- Sanderson, B. E. (1969). Acetylcholinesterase activity in Nippostrongylus brasiliensis (Nematoda). Comp. Biochem. Physiol., 29, 1207-1213.

- Sandground, J. H. (1925). Speciation and specificity in the nematode genus Strongyloides. J. Parasit., 12, 59-80.
- Sandground, J. H. (1926). Biological studies on the life cycle of the genus Strongyloides Grassi, 1879. Amer. J. Hyg., 6, 337-388.
- Sandground, J. H. (1928). Some studies on susceptibility, resistance and acquired immunity to infection with Strongyloides stercoralis (Nematoda) in dogs and cats. Amer. J. Hyg., 8, 507-538.
- Sarles, M. P. (1938). The in vitro action of immune serum on the nematode Nippostrongylus muris. J. Inf. Dis., 62, 337-348.
- Sarles, M. P. and Taliaferro, W. H. (1936). The local points of defense and the passive transfer of acquired immunity to Nippostrongylus muris in rats. J. Inf. Dis., 59, 207-220.
- Sasa, M., Hayashi, S., Tanaka, H., Sato, K., Miura, A., Wakasugi, M., Shirasaka, R., Takata, A., Tokuriki, K., Nakamura, E. and Nagano, M. (1957). Comparative studies on parasitic infections among coal miners in Kyushu and Hokkaido. Koshu Eisei, 21, 1-9.
- Schacher, J. F. and Chee-Hock, C. (1960). Malaysian parasites. XLVII. Nematode parasites of three common house rat species in Malaya, with notes on Rictularia tani Hoeppli, 1929. Stud. Inst. Med. Res., Malaya, 29, 209-216.
- Selby, G. R. and Wakelin, D. (1973). Transfer of immunity against Trichuris muris in the mouse by serum and cells. Internat. J. Parasit., 3, 717-722.
- Semrad, J. E. and Coors, M. J. (1951). The rise and fall of immunity to Trichinella spiralis in the albino rat and its effects on growth and reproduction of the parasite. Trans. Illin. Acad. Sci., 44, 253-258.
- Sheldon, A. J. (1937a). Some experimental studies on Strongyloides ratti. Amer. J. Hyg., 25, 39-52.
- Sheldon, A. J. (1937b). Studies on active acquired resistance, natural and artificial, in the rats to infection with Strongyloides ratti. Amer. J. Hyg., 25, 53-65.
- Sheldon, A. J. (1937c). Studies on routes of infection of rats with S. ratti. Amer. J. Hyg., 26, 358-373.
- Sheldon, A. J. (1937d). Successful infection of mice with S. ratti. J. Parasit., 23, 98.

- Sheldon, A. J. (1939). Specificity of artificial acquired immunity to *S. ratti*. *Amer. J. Hyg.*, 29, 47-50.
- Sinclair, I. J. (1970). The relationship between circulating antibodies and immunity to helminthic infections. *Adv. Parasitol.*, 8, 97-138.
- Sinski, E. and Krzysztofowicz, M. (1971). The blood picture and immunological reactions in rabbits exposed to repeated doses of the sheep and rabbit strains of *Strongyloides papillosus*. *Acta Parasitol. Polonica*, 19, 427-444.
- Sjögren, H. O., Hellström, I., Bansal, S. C. and Hellström, K. E. (1971). Suggestive evidence that the "blocking antibodies" of tumor-bearing individuals may be antigen-antibody complexes. *Proc. Nat. Acad. Sci.*, 68, 1372-1375.
- Smith, I. D., Goodrich, B. S., Kelly, J. D. and Dineen, J. K. (1974). Prostaglandin-like activity of various fractions of ram semen: their role in the immune rejection of the nematode *Nippostrongylus brasiliensis* from the small intestine of rats. *Prostaglandins*, 5, 87-96.
- Smith, E. W. and Atkinson, W. B. (1956). Simple procedure for identification and rapid counting of mast cells in tissues. *Science N.Y.*, 123, 941.
- Smithers, S. R. (1962). Stimulation of acquired resistance to *Schistosoma mansoni* in monkeys: role of eggs and worms. *Exptl Parasit.*, 12, 263-273.
- Smithers, S. R. and Terry, R. J. (1965). The infection of laboratory hosts with cercariae of *Schistosoma mansoni* and the recovery of the adult worms. *Parasitology*, 55, 695-700.
- Smyth, J. D. (1962). Introduction to animal parasitology. The English Universities Press Ltd., London. 470 p.
- Søndergaard, J. and Greaves, M. W. (1970). Recovery of a pharmacologically active fatty acid during the inflammatory reaction invoked by patch testing in allergic contact dermatitis. *Internat. Arch. Allerg. Appl. Immunol.*, 39, 56-61.
- Sotolongo Guarra, F. (1972). Parasitic males of *S. stercoralis*. *Parassitologia*, 14, 29-33.
- Soulaby, E. J. L. (1958). Studies on the heterophil antibodies associated with helminth infections. III. Heterophil antibody in *Oesophagostomum dentatum* infections in pigs. *J. Comp. Path. Therap.*, 68, 380-387.

- Soulsby, E. J. L. (1962). Antigen-antibody reactions in helminth infections. Adv. Immunol., 2, 265-308.
- Spindler, L. A. (1958). The occurrence of the intestinal thread-worms Strongyloides ratti in the tissues of rats, following experimental percutaneous infection. Proc. Helm. Soc. Wash., 25, 106-111.
- Stankiewicz, M., Bezubik, B. and Sinaki, E. (1971). The effect of UV-rays on the infective larvae of Strongyloides papillosus in the rabbit. III. Production of immunity by irradiated larvae. Acta Parasitol. Polonica, 19, 327-336.
- Stoner, R. D. and Godwin, J. T. (1953). The effects of ACTH and cortisone upon susceptibility to trichinosis in mice. Amer. J. Path., 29, 943-950.
- Stoner, R. D. and Godwin, J. T. (1954). The effects of adrenocorticotrophic hormone and cortisone upon acquired immunity to trichinosis in mice. Amer. J. Path., 30, 913-918.
- Tanabe, H. and Takeishi, H. (1936). A survey of helminths in the digestive system of the rat. Keio, Igaku, 16, 1767-1785.
- Taliaferro, W. H. (1940). The mechanism of acquired immunity to infections with parasitic worms. Physiol. Rev., 20, 469.
- Taliaferro, W. H. and Sarles, M. P. (1939). The cellular reactions in the skin, lung and intestine of normal and immune rats after infection with Nippostrongylus muris. J. Inf. Dis., 64, 157-192.
- Tashiro, I. (1912). Studies on Strongyloides stercoralis in Japan, with special reference to its infection in dogs. Saikingaku Zasshi, 204, 805-811.
- Tewari, H. C. (1970). Studies on the resistance of guinea-pigs to Fasciola hepatica infection. Ind. J. Anim. Hlth, 9, 157-161.
- Thomas, H. (1970). Quantitative investigation of host-parasite relationship in Strongyloides ratti infected rats. (Abstract) J. Parasit., 56.
- Thompson, J. and van Furth, R. (1970). The effect of glucocorticosteroid on the kinetics of mononuclear phagocytes. J. expl Med., 131, 429-442.
- Toh, C. C. S. and Chow, K. W. (1969). Malabsorption syndrome in a patient infected with Strongyloides stercoralis. Ann. trop. Med. Parasit., 63, 493-497.

- Urquhart, G. M., Mulligan, W., Eadie, R. M. and Jennings (1965). Immunological studies on Nippostrongylus brasiliensis infection in the rat. The role of local anaphylaxis. Expl Parasit., 17, 210-217.
- Uvnas, B. and Wold, J. K. (1967). Isolation of a mast cell degranulating polypeptide from Ascaris suis. Acta Physiol. Scand., 70, 269-276.
- Van Drum, P. (1902). Quelques notes sur les embryons de Strongyloides intestinalis et leur penetration par la peau. Thompson Yates Lab. Rept., Liverpool, 4, 471-474.
- Venable, J. H. and Coggeshall, R. E. (1965). A simplified lead citrate stain for use in electron microscopy. J. Cell Biol., 25, 407-408.
- Wagland, B. M. and Dineen, J. K. (1965). The cellular transfer of immunity to Trichostrongylus colubriformis in an isogenic strain of guinea-pigs. Aust. J. expl. Biol. Med. Sci., 43, 429-438.
- Wakelin, D. (1967). Acquired immunity to Trichuris muris in the albino laboratory mouse. Parasitology, 57, 515-524.
- Wakelin, D. (1969). Studies on the immunity of albino mice to Trichuris muris. The stimulation of immunity by chemically abbreviated infections. Parasitology, 59, 549-555.
- Wakelin, D. (1970). Studies on the immunity of albino mice to Trichuris muris; suppression of immunity by cortisone acetate. Parasitology, 60, 229-237.
- Wakelin, D. (1975). Immune expulsion of Trichuris muris from mice during a primary infection: analysis of the components involved. Parasitology, 70, 397-405.
- Wakelin, D. and Lloyd, M. (1976a). Immunity to primary and challenge infections of Trichinella spiralis in mice: a re-examination of conventional parameters. Parasitology, 72, 173-182.
- Wakelin, D. and Lloyd, M. (1976b). Accelerated expulsion of adult Trichinella spiralis in mice given lymphoid cells and serum from infected donors. Parasitology, 72, 307-315.
- Wakelin, D. and Selby, G. R. (1974a). The induction of immunological tolerance to the parasitic nematode T. muris in cortisone-treated mice. Immunology, 26, 1-10.

- Walker-Smith, J. A., McMillan, B., Middleton, A. W., Robertson, S. and Hopcroft, A. (1969). Strongyloidiasis causing small-bowel obstruction in an aboriginal infant. Med. J. Aust., 20, 1263-1265.
- Walls, R. S., Basten, A., Leuchars, E. and Davies, A. J. S. (1971). Mechanism for eosinophilic and neutrophilic leucocytoses. Bri. Med. J., 111, 157-159.
- Walls, R. S., Carter, R. L., Leuchars, E. and Davies, A. J. S. (1973). The immunopathology of trichiniasis in T. cell-deficient mice. Clin. Exp. Immunol., 13, 231-242.
- Weinstein, P. P. (1953). The effect of cortisone on the development of the immune response in the white rat to Nippostrongylus muris. J. Parasit., 39, 35.
- Weinstein, P. P. (1955). The effect of cortisone on the immune response of the white rat to Nippostrongylus muris. Amer. J. trop. Med. Hyg., 41, 61-74.
- Wells, P. D. (1962). Mast cell, eosinophil and histamine levels in Nippostrongylus brasiliensis infected rats. Exptl Parasit., 12, 82-101.
- Wertheim, G. (1963). Helminth parasites of the rat Rattus norvegicus from Haifa and Tel Aviv. Bull. Res. Council Israel, Sec. E, 10E, 125-129.
- Wertheim, G. (1970a). Growth and development of Strongyloides venezuelensis Brumpt, 1934 in the albino rat. Parasitology, 61, 381-388.
- Wertheim, G. (1970b). Experimental concurrent infections with Strongyloides ratti and S. venezuelensis in laboratory rats. Parasitology, 61, 389-395.
- Wertheim, G. and Lengy, J. (1964). The seasonal occurrence of Strongyloides ratti Sandground, 1925 and S. venezuelensis Brumpt, 1934, in a population of Rattus norvegicus. J. Helminth., 38, 393-398.
- Wertheim, G. and Lengy, J. (1965). Growth and development of Strongyloides ratti Sandground, 1925, in the albino rat. J. Parasit., 51, 636-639.
- Whur, P. (1966). Relationship of globule leucocytes to gastro-intestinal nematodes in sheep, and Nippostrongylus brasiliensis and Hymenolpis nana infections in rats. J. Comp. Path., 76, 57-67.

- Whur, P. (1967). Globule leucocyte response in hyper-immune rats infected with Nippostrongylus brasiliensis. J. Comp. Path., 77, 271-277.
- Willis, A. J. P. and Nwokolo, C. (1966). Steroid therapy and strongyloidiasis. Lancet, 1, 1396-1398.
- Wilson, R. J. M., Jones, V. E. and Eskowitz, S. (1967). Thymectomy and anaphylactic antibody in rats infected with Nippostrongylus brasiliensis. Nature (Lond.), 213, 398-399.
- Wilson, S. and Thompson, A. (1964). A fatal case of strongyloidiasis. J. Path. Bact., 87, 169-176.
- World Health Organization (1963) African Conference on Ancylostomiasis. Technical Report Series, 225, CCTA/WHO.
- Yamaguti, S. (1961). Systema helminthum. Vol. III. The nematodes of vertebrates, Parts I and II. Interscience Publishers, Inc., New York.
- Yim, Y., Kikkawa, Y., Tanowitz, H. and Wittner, M. (1970). Fatal strongyloidiasis in Hodgkin's disease after immunosuppressive therapy. J. trop. Med. Hyg., 73, 245-249.
- Yoeli, M., Most, H., Berman, H. H., and Scheinsson, G. P. (1963). The clinical picture and pathology of a massive Strongyloides infection in a child. Trans. R. Soc. trop. Med. Hyg., 57, 346.
- Zaiman, H. (1970). Drug treatment of trichinosis. Chapter 12 in "Trichinosis in man and animals" (Gould, S. E., ed.). Charles C. Thomas, Illinois.
- Zaiman, H., Scardino, V., Bersen, P. and Stern, R. C. (1963). Further studies on the eosinophil response of rats infected with irradiated Trichinella. Expl Parasit., 14, 1-8.
- Zaiman, H., Omi, G. and Gapinski, L. (1953). Studies on the nature of immunity to Trichinella spiralis in parabiotic "uninfected" rats as determined by observations of muscular invasion. Amer. J. Hyg., 57, 306-310.
- Zinn, W. (1899). Ueber A. intestinalis. Centralbl. f. Bakt. Parasitenk., 26, 396-398.

ADDENDUM

- Adam, M., Morgan, O., Persaud, C. and Gibbs, W. N. (1973). Hyperinfection syndrome with Strongyloides stercoralis in malignant lymphoma. Br. Med. J., 1, 264-266.
- Aiken, J. W. (1972). Aspirin and indomethacin prolong parturition in rats. Evidence that prostaglandins contribute to expulsion of foetus. Nature, 240, 21-25.
- Boggs, D. R., Athens, J. W., Cartwright, G. E. and Wintrobe, (1964). The effect of adrenal gluco-corticosteroids upon the cellular composition of inflammatory exudates. Amer. J. Path., 44, 763-773.
- Cavallero, C. and Braccini, C. (1950). Cortisone e mastociti. Boll. Soc. Ital. Pat., 1, 3.
- Cavallero, C. and Braccini, C. (1951). Effect of cortisone on the mast cells of the rat. Proc. Soc. Exp. Biol., 78, 141.
- Ebert, R. H. (1952). In vivo observations on the effect of cortisone on experimental tuberculosis using the rabbit ear chamber technique. Amer. Rev. Tuberculosis, 65, 64-74.
- Fernex, M. (1968). The mast-cell system: Its relationship to Athero-Sclerosis, Fibrosis and Eosinophils. S. Karger, Basle.
- Kay, A. B. (1976). Function of eosinophil leukocytes. (Annotation). Br. J. Haemat., 33, 313-318.
- Smith, I. D. and Temple, D. (1973). The influence of analgesic drugs on the action of prostaglandin F₂ on the human uterus in vivo and rabbit myometrial strips in vitro. Prostaglandins, 4, 469-477.
- Sotolongo Guerra, F., Kouri, J. E. and Perez, S. Q. (1971). Finding of a male parasitic Strongyloides stercoralis. Revta Cub. Med. Trop., 23, 3-11.
- Thomas, J., Renambot, J., Barbotin, M., Riquaud, J. and Saint-Anore, P. (1971). Urticaire lineaire recidivante mobile au cours de la strongyloïdose guerison par le Thiabendazole. Bull. Soc. Path. Exot., 64, 486-497.
- Vane, J. R. (1971). Inhibition of prostaglandin synthesis as a mechanism of action of aspirin-like drugs. Nature, 231, 232-235.