

## RESEARCH ARTICLE

# Inter-pregnancy interval and associated adverse maternal outcomes among women who delivered at Kilimanjaro Christian Medical Centre in Tanzania, 2000-2015

Leah Anku Sanga<sup>1\*</sup>, Tara Mtuy<sup>1,2</sup>, Rune Nathaniel Philemon<sup>1,3</sup>, Michael Johnson Mahande<sup>1</sup>

**1** Department of Epidemiology & Biostatistics, Institute of Public Health, Kilimanjaro Christian Medical University College, Kilimanjaro, Tanzania, **2** Department of Global Health & Development, London School of Hygiene & Tropical Medicine, London, United Kingdom, **3** Department of Community Health, Institute of Public Health, Kilimanjaro Christian Medical University College, Kilimanjaro, Tanzania

☞ These authors contributed equally to this work.

\* [leah.ameye@gmail.com](mailto:leah.ameye@gmail.com)



## OPEN ACCESS

**Citation:** Sanga LA, Mtuy T, Philemon RN, Mahande MJ (2020) Inter-pregnancy interval and associated adverse maternal outcomes among women who delivered at Kilimanjaro Christian Medical Centre in Tanzania, 2000-2015. PLoS ONE 15(2): e0228330. <https://doi.org/10.1371/journal.pone.0228330>

**Editor:** Joshua Amo-Adjei, University of Cape Coast, GHANA

**Received:** May 3, 2019

**Accepted:** January 13, 2020

**Published:** February 6, 2020

**Copyright:** © 2020 Sanga et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The data from this study is can not be shared publicly because it contains personal identifiers and sensitive data on women's reproduction. This data was obtained from Kilimanjaro Christian Medical Centre who is the sole owner of the Medical Birth Registry and has imposed data protection policy on the database. It therefore has legal and ethical restrictions on sharing. However, data are available upon request from the Kilimanjaro Christian Medical Centre Institutional Data access committee

## Abstract

Inter-pregnancy interval is an important determinant of both maternal and child health. Sub-optimal inter-pregnancy interval has been associated with adverse maternal outcomes -including postpartum hemorrhage and hypertensive disorders, direct causes of maternal mortality. Both overall maternal mortality and the contribution of postpartum hemorrhage on maternal mortality have increased in Tanzania. If we are to achieve sustainable development goal (SDG) number 3.1 to reduce the global maternal mortality ration to less than 70 per 100,000 live births by 2030, it is highly important that such contributors are dealt with. This study aimed to determine the distribution and trends of inter-pregnancy interval and to deduce its association with adverse maternal outcomes among women who delivered at Kilimanjaro Christian Medical Centre (2000–2015). A retrospective cohort study was designed using Kilimanjaro Christian Medical Centre medical birth registry data for women who delivered from 2000 to 2015. Women with at least two births recorded in the registry were included. A total of 7,995 births from 6,612 mothers were analyzed. Anemia during pregnancy, post-partum hemorrhage and pre-eclampsia were adverse maternal outcomes of interest. Data analysis was performed using multivariable logistic regression models allowing for robust standard errors. Crude and adjusted odds ratio with their respective 95% confidence intervals were estimated. More than half (51.7%) of non-first births were born within sub-optimal IPI. The median IPI was 34 months (IQR: 33.5 months). The median IPI increased from 11 months in 2002 to 35 months in 2006 and plateaued until 2014, but it rose to 41.6 months in 2015. Median IPI was shorter in young women <20 years and in birth order seven and above (16 vs. 27 months, respectively). Short IPI was associated with lower risk of pre-eclampsia [aOR: 0.71, 95%CI: 0.52, 0.97] while long IPI was associated with lower risk of post-partum hemorrhage [aOR: 0.70, 95%CI: 0.52, 0.94]. This study found an association between long and short IPI with adverse maternal outcomes. Even though these results should be interpreted with caution based on the fact that the data was sampled from a referral hospital and hence there could be overrepresentation of women with maternal complications,

which can be contacted through e-mail: [kcmcadmin@kcmc.ac.tz](mailto:kcmcadmin@kcmc.ac.tz) or through postal address: Executive Director, KCMC, P. O. Box 3010, Moshi, Tanzania

**Funding:** The author(s) received no specific funding for this work

**Competing interests:** The authors have declared that no competing interests exist

our findings still warrant the importance of supporting modern family planning methods as a measure to improve IPI and thereby improve maternal outcomes as well.

## Introduction

Inter-pregnancy interval (IPI) is the time lapse between termination of one pregnancy and conception of a subsequent one [1]. Optimal IPI can ensure optimal health for both mother and infant [2] whereas sub-optimal IPI has been associated with several maternal morbidities and mortality [3–10]. World Health Organization (WHO) recommends an interval of at least 24 months between a live birth and the next pregnancy [1] while others have argued that IPI of 3–5 years further reduces the risk for adverse maternal outcomes [11]. Different theories—particularly maternal depletion theory and physiological regression theory—link inter-pregnancy interval with adverse maternal outcomes [3,12].

Hemorrhage and hypertensive disorders are among maternal morbidities associated with IPI (Fig 1). They are among direct causes of maternal deaths and account for approximately half of the maternal deaths globally [13–15]. Approximately 800 women die every day globally due to pregnancy and childbirth related complications [14] with the global maternal mortality ratio (MMR) being 211 per 100,000 live births [16]. Most (99%) of these deaths occur in low income countries, with majority (66%) from Sub-Saharan Africa, including Tanzania [14,17]. It is estimated that 20–35% of maternal deaths can be prevented by adhering to recommended IPI [18].

Maternal mortality and morbidity can be averted by dealing with preventable factors that cause them. Despite several interventions [19–22], maternal mortality has risen from 432 in 2012 to 524 per 100,000 live births in 2017 in Tanzania [16]. The country failed to reach the Millennium Development Goal (MDG) number 5 target for improved maternal health and the trend is not promising to enhance achievement of Sustainable Development Goal number 3 to reduce maternal mortality ratio to less than 70 per 100,000 live births by 2030 [20,23,24]. Furthermore, the proportion of maternal deaths attributed to hemorrhage has increased from 15% in 2010 to 18% in 2015 and there has been no improvement in proportion of deaths caused by hypertensive disorders from 2006 to 2015 [19,25].


Several factors have been reported to influence IPI [26–30], but of ultimate importance is modern contraceptive use [30]. However, utilization of modern contraceptives in Tanzania is still low and the proportion of births born within short IPI has increased by 3% in the past decade [19,25].

Non-adherence to optimal IPI could be one of the drivers increasing MMR in Tanzania since it is associated with direct causes of maternal mortality. However, in our settings, little is known about the role of IPI on adverse maternal outcomes. Understanding the association between IPI and adverse maternal outcomes may help to provide important information needed to help design evidence-based interventions to reduce its impact, and hence accelerate the achievement of SDG3 by 2030. This study aimed to determine the distribution, trend and patterns of IPI among women who delivered at Kilimanjaro Christian Medical Centre in northern Tanzania between 2005 and 2015. The study also assessed the association between IPI and adverse maternal outcomes.

## Materials and methods

### Study setting

This study was conducted at the Department of Obstetrics and Gynecology of Kilimanjaro Christian Medical Centre (KCMC). KCMC is a consultant and university teaching hospital



**Fig 1. Directed acyclic diagram showing relationship between inter-pregnancy interval and adverse maternal outcomes.**

<https://doi.org/10.1371/journal.pone.0228330.g001>

located in Moshi urban district, Kilimanjaro region in northern Tanzania. It is one of the four zonal hospitals in Tanzania which serves over 15 million people, including the local community and referred cases from six regions in Tanzania. Pregnant high-risk women are referred to KCMC for observation and delivery while women from the local community may come to deliver on their own accord. The main catchment area of KCMC hospital is Kilimanjaro region.

### Design, sample and sampling

This study used secondary data from a cohort of women who delivered at KCMC from 2000 to 2015. Women were followed retrospectively through their medical records. We restricted our analysis to two or more deliveries from the same mother, delivered at KCMC during the study period. Women who did not have consecutive deliveries during the study time were excluded. A total of 19,000 deliveries were recorded during the study period. Of these, 7,995 births from 6,612 mothers were eligible for the study.

### Data collection

Data were extracted from the medical birth registry at KCMC. The medical birth registry records information for all mothers who deliver in the Obstetrics and Gynecology Department at KCMC within 24 hours after delivery or as soon as mothers have recovered in case of complicated deliveries. Information was collected on a daily basis by a trained nurse/midwife including, but not limited to, maternal and paternal socio-demographics, mother's reproductive history and conditions and complications of the mother before, during and after pregnancy/delivery and puerperium. In the interview, women also provided information about births

delivered outside KCMC during the study period. However, information on miscarriages and abortions, whether they happened at KCMC or elsewhere, were not captured in the birth registry. Each woman and her pregnancy outcome were given a unique identifier which was used to link deliveries to a specific mother. Data were entered in a computerized system of the birth registry.

### Data analysis

Data analysis was performed using STATA version 13.1 statistical software (StatCorp, College Station, TX).

The study looked at three adverse maternal outcomes; post-partum hemorrhage (PPH), pre-eclampsia and anemia. PPH was considered if a woman lost  $\geq 500$ mls of blood after vaginal delivery or  $\geq 1000$ mls of blood after caesarean delivery [1]. Blood loss was visually estimated by a nurse midwife after vaginal delivery and by the doctor performing the surgery in caesarean delivery. A woman was considered to have anemia if at any point in her pregnancy she had measured (by any validated method) Hemoglobin level below 8.5 grams per deciliter (8.5g/dl). Pre-eclampsia was defined as a disorder characterized by development of hypertension 140/90 mm Hg or more with proteinuria after the 20<sup>th</sup> week of gestation in a previously normotensive and non-proteinuric woman[1].

Inter-pregnancy interval (IPI) was the main exposure for these outcomes and it was computed as follows:

$$\text{IPI (days)} = [\text{D.O.B of index child} - \text{D.O.B of preceding child}] - \text{Gestation age at birth of index child}$$

IPI was categorized into <24 months (short IPI), 24–59 (reference group/optimal IPI) and 60+ months (long IPI). Mother's age, education level, occupation, religion, marital status, tribe, current residence, pregnancy type, family planning use, alcohol use, anemia in previous pregnancy, PPH in previous pregnancy, pre-eclampsia in previous pregnancy, referral status, delivery mode, parity, any ANC visit, number of ANC visits and death of a preceding child were considered as confounders.

Since children are clustered within a mother, hierarchy of data was considered but there was no effect of clustering on the outcomes. In the univariate and multivariable analysis, logistic regression with robust standard errors was used to assess the association between IPI and each of the selected adverse maternal outcomes. Confounding was established if a variable changed the Odds Ratios for the effect of IPI on any adverse maternal outcome by 10% or more. To determine if there was a dose-response relationship between IPI and adverse maternal outcomes, Generalized Additive Models were used. However, all three maternal outcomes were deemed to have a linear association with IPI. While analyzing trends of IPI over the years, the years 2000 and 2001 were excluded from the analysis due to few numbers of subjects. Significance was considered at 5% level.

### Ethical considerations

For practical reasons, since the interview was administered just after the woman had given birth, consent was given orally. The midwife nurse gave every woman oral information about the birth registry, the data needed to be collected from them and the use of the data for research purposes. Then informed consent was sought verbally from every woman. Following the consent, the woman could still opt not to reply to individual questions. Inclusion of the orally conducted interview in the database was conditional on informed consent which was followed as far as the orally conducted interview was concerned. All consent procedures were approved by the Kilimanjaro Christian Medical Centre ethical committee.

Ethical clearance for the establishment of Medical Birth registry was granted from Ethics Committee at Kilimanjaro Christian Medical Centre, approved by Tanzania's Ministry of Health and National Ethics Committee in Norway. Approval to carry out this study was obtained from Tumaini University Makumira through Kilimanjaro Christian Medical University College Research and Ethical Committee.

## Results

### Socio-demographic and pregnancy characteristics of study participants at baseline delivery

The mean mother's age of all women in the sample was 30.5 (SD: 5.3 years) with most women aged between 20–34 years. The majority of women were employed, married and living in urban areas (Table 1). Most of them used modern family planning and attended four or more ANC visits. A lower proportion (2.8%) of women with long IPI had PPH in preceding pregnancy while a higher proportion of women with short IPI had pre-eclampsia (4.9%) and had experienced perinatal death in their preceding birth (11.2%) (Table 2).

### Characteristics of women who had adverse maternal outcomes

Of all pregnancies studied, a total of 309 (3.9%) experienced PPH, 80 (1.0%) experienced anaemia and 270 (3.4%) experienced Pre-eclampsia. Distribution of proportion of these outcomes by inter-pregnancy interval is explained in Table 3. A total of 692 (10.5%) women experienced one of the adverse maternal outcomes in at least one of their subsequent deliveries. PPH was the most frequent adverse outcome (53%) among women who had adverse maternal outcomes. A few (12%) of these women had anemia and 39% had pre-eclampsia.

More than half of all women in the sample had IPI of 24–59 months (51.1%), were aged 20–34 years (65.9%) and were married (97%) (Table 4).

Across all adverse outcomes, most women delivered by caesarean section. Higher proportion (14.1%) of women who had pre-eclampsia had experienced perinatal death in the preceding child compared to those who had anemia (3.8%) and PPH (8.4%) (Table 5).

### Distribution and trends of IPI across birth orders

About half of the women had sub-optimal IPI (short = 32.9%; long = 18.8%) with the median IPI 34 months (IQR: 33.4 months). The median IPI improved from 11 months in 2002 to 34.8 months in 2006 and remained stable up to 2014. In 2015, it raised to 41.6 months (Fig 2).

There was no difference in median IPI in the 2<sup>nd</sup> to 3<sup>rd</sup> birth order to that in 4<sup>th</sup> to 6<sup>th</sup> birth order (median IPI = 33.6 and 35 months, respectively). In higher birth orders (7 and above), median IPI was shorter (26.6 months) compared to lower birth orders (Fig 3).

The median IPI was longer in older women aged 35 and above (median IPI = 32.3 months) and shorter in younger women <20 years (median IPI = 15.7 months) (Fig 4).

### Association between IPI and adverse maternal outcomes

In the univariate analysis, long IPI was significantly associated with anemia at 5% level while there was no significant association between long IPI and pre-eclampsia or PPH. Short IPI was not significantly associated with any of the adverse maternal outcomes at 5% level. Parity, delivery mode and pregnancy type were associated with all of the three adverse maternal outcomes in univariate analysis (Table 6). After adjusting for other variables, short IPI was significantly associated with lower risk of pre-eclampsia while long IPI was significantly associated with lower risk of PPH at 5% level. Women with short IPI had 29% lower odds of having pre-

**Table 1. Socio-demographic characteristics of women by inter-pregnancy interval at baseline (N = 6,612).**

| Variable                          | Inter-pregnancy interval (months) |      |                 |       |               |      |
|-----------------------------------|-----------------------------------|------|-----------------|-------|---------------|------|
|                                   | <24 N = 2,174                     |      | 24–59 N = 3,193 |       | 60+ N = 1,245 |      |
|                                   | n                                 | %    | n               | %     | n             | %    |
| <b>Age categories</b>             |                                   |      |                 |       |               |      |
| ≤19                               | 51                                | 2.3  | 11              | 0.3   | 3             | 0.2  |
| 20–34                             | 1,853                             | 85.2 | 2,581           | 80.8  | 801           | 64.3 |
| ≥35                               | 268                               | 12.3 | 596             | 18.7  | 441           | 35.4 |
| Missing                           | 2                                 | 0.1  | 5               | 0.2   | -             | -    |
| <b>Mother’s education</b>         |                                   |      |                 |       |               |      |
| No formal education               | 28                                | 1.3  | 29              | 0.9   | 13            | 1.0  |
| Primary                           | 1,075                             | 49.4 | 1,750           | 54.8  | 720           | 57.8 |
| Secondary and above               | 1,068                             | 49.1 | 1,409           | 44.1  | 512           | 41.1 |
| Missing                           | 3                                 | 0.1  | 5               | 0.2   | -             | -    |
| <b>Mother’s occupation</b>        |                                   |      |                 |       |               |      |
| Employed                          | 1,528                             | 70.3 | 2,370           | 74.2  | 1,010         | 81.1 |
| Unemployed                        | 636                               | 29.3 | 814             | 25.5  | 235           | 18.9 |
| Missing                           | 10                                | 0.5  | 9               | 0.3   | -             | -    |
| <b>Religion</b>                   |                                   |      |                 |       |               |      |
| Catholic                          | 911                               | 41.9 | 1,243           | 38.9  | 521           | 41.8 |
| Protestant                        | 852                               | 39.2 | 1,314           | 41.2  | 491           | 39.4 |
| Muslim                            | 398                               | 18.3 | 621             | 19.4  | 229           | 18.4 |
| Others                            | 7                                 | 0.3  | 11              | 0.3   | 2             | 0.2  |
| Missing                           | 6                                 | 0.3  | 4               | 0.1   | 2             | 0.2  |
| <b>Marital status</b>             |                                   |      |                 |       |               |      |
| Married                           | 2,094                             | 96.3 | 3,098           | 97.0  | 1,182         | 94.9 |
| Single                            | 72                                | 3.3  | 83              | 2.6   | 63            | 5.1  |
| Missing                           | 8                                 | 0.4  | 12              | 0.4   | -             | -    |
| <b>Mother’s current residence</b> |                                   |      |                 |       |               |      |
| Rural                             | 764                               | 35.1 | 1,055           | 33    | 366           | 29.4 |
| Urban                             | 1,301                             | 59.8 | 2,018           | 63.2  | 827           | 66.4 |
| Semi urban                        | 107                               | 4.9  | 119             | 3.7   | 52            | 4.2  |
| Missing                           | 2                                 | 0.1  | 1               | <0.01 | -             | -    |

Modern family planning include: pills, injections, implants and intra-uterine devices

<https://doi.org/10.1371/journal.pone.0228330.t001>

eclampsia as compared to those with optimal IPI [aOR: 0.71, 95%CI: 0.52, 0.97]. On the other hand, women with long IPI had 30% lower odds for post-partum hemorrhage [aOR: 0.70, 95% CI: 0.52, 0.94] compared to those with optimal IPI (Table 7).

### Discussion

This study found that about half of the women at KCMC do not adhere to WHO recommendation on IPI. The median IPI increased yearly from 2002 to 2006, and then it plateaued and rose again in 2015.

We also noted that, IPI was shorter in younger women (<20 years) compared to older women and shorter in higher birth orders (seven and above) compared to lower birth orders. Our results suggest that women with short IPI are at lower risk of pre-eclampsia and those with long IPI are at lower risk of PPH. This implies sub-optimal IPI is protective for these adverse maternal outcomes. Lower risk of pre-eclampsia in short IPI was however incongruent

Table 2. Health-related characteristics of women by inter-pregnancy interval at baseline birth (N = 6,612).

| VARIABLES                                  | Inter-pregnancy interval (Months) |      |                 |      |               |      |
|--|-----------------------------------|------|-----------------|------|---------------|------|
|  | <24 N = 2,174                     |      | 24–59 N = 3,193 |      | 60+ N = 1,245 |      |
|  | n                                 | %    | n               | %    | n             | %    |
| <b>Type of family planning used</b>        |                                   |      |                 |      |               |      |
| None                                       | 375                               | 17.2 | 179             | 5.6  | 61            | 4.9  |
| Modern                                     | 1,073                             | 49.4 | 2,335           | 73.1 | 1,034         | 83.1 |
| Traditional                                | 708                               | 32.6 | 660             | 20.7 | 147           | 11.8 |
| Missing                                    | 18                                | 0.8  | 19              | 0.6  | 3             | 0.2  |
| <b>Alcohol use before/during pregnancy</b> | 587                               | 27.0 | 984             | 30.8 | 430           | 34.5 |
| <b>PPH in preceding birth</b>              | 74                                | 3.4  | 112             | 3.5  | 35            | 2.8  |
| <b>Anemia in preceding birth</b>           | 36                                | 1.7  | 53              | 1.7  | 15            | 1.2  |
| <b>Preeclampsia in preceding birth</b>     | 106                               | 4.9  | 110             | 3.4  | 43            | 3.5  |
| <b>Referred from</b>                       |                                   |      |                 |      |               |      |
| Home                                       | 1,945                             | 89.5 | 2,964           | 92.8 | 1,183         | 95.0 |
| Regional hospital                          | 44                                | 2.0  | 73              | 2.3  | 19            | 1.5  |
| District hospital                          | 34                                | 1.6  | 42              | 1.3  | 19            | 1.5  |
| Other                                      | 44                                | 2.0  | 71              | 2.2  | 23            | 1.8  |
| Missing                                    | 107                               | 4.9  | 43              | 1.3  | 1             | 0.1  |
| <b>Delivery mode</b>                       |                                   |      |                 |      |               |      |
| Spontaneous                                | 1,442                             | 66.3 | 1,943           | 60.9 | 726           | 58.3 |
| Assisted                                   | 19                                | 0.9  | 36              | 1.1  | 16            | 1.3  |
| CS   | 710                               | 32.7 | 1,208           | 37.8 | 502           | 40.3 |
| Missing                                    | 3                                 | 0.1  | 6               | 0.2  | 1             | 0.1  |
| <b>Death of preceding birth</b>            | 243                               | 11.2 | 77              | 2.4  | 17            | 1.4  |
| Missing                                    | 4                                 | 0.2  | 10              | 0.3  | 2             | 0.2  |
| <b>Parity</b>                              |                                   |      |                 |      |               |      |
| 2  | 1,301                             | 59.8 | 1,765           | 55.3 | 550           | 44.2 |
| 3  | 490                               | 22.5 | 834             | 26.1 | 412           | 33.1 |
| >3   | 383                               | 17.6 | 594             | 18.6 | 283           | 22.7 |
| <b>Antenatal care attendance</b>           | 2,162                             | 99.6 | 3,178           | 99.8 | 1,241         | 99.7 |
| <b>Number of ANC visits</b>                |                                   |      |                 |      |               |      |
| <4   | 864                               | 39.7 | 1,082           | 33.9 | 442           | 35.5 |
| ≥4   | 1,279                             | 58.8 | 2,083           | 65.2 | 794           | 63.8 |
| Missing                                    | 31                                | 1.4  | 28              | 0.9  | 9             | 0.7  |

<https://doi.org/10.1371/journal.pone.0228330.t002>

with a study done in Latin America and The Caribbean[6], a difference which could probably be due to methodological differences. Conde-Agudelo’s [6] methodology ignored dependency which could have caused underestimation of standard errors, leading to smaller p-values and wrong inferences. In this study, data dependency was taken into consideration using robust

Table 3. Proportion of women with adverse outcomes by inter-pregnancy interval (N = 7,995).

| IPI categories (months) | Total<br>N | PPH (N = 369) |     | ANAEMIA (N = 80) |     | PRE-ECLAMPSIA (N = 270) |     |
|-------------------------|------------|---------------|-----|------------------|-----|-------------------------|-----|
|                         |            | n             | %   | n                | %   | n                       | %   |
| 24–59                   | 3,890      | 205           | 5.3 | 47               | 1.2 | 138                     | 3.5 |
| <24                     | 2,625      | 100           | 3.8 | 26               | 1.0 | 77                      | 2.9 |
| >59                     | 1,480      | 64            | 4.3 | 7                | 0.5 | 55                      | 3.7 |

<https://doi.org/10.1371/journal.pone.0228330.t003>

Table 4. Socio-demographic characteristics of women by adverse maternal outcomes (N = 692).

| VARIABLES                | PPH (n = 369) |      | Anemia (n = 80) |      | Pre-eclampsia(n = 270) |      |
|--------------------------|---------------|------|-----------------|------|------------------------|------|
|                          | n             | %    | n               | %    | n                      | %    |
| <b>IPI categories</b>    |               |      |                 |      |                        |      |
| 24–59                    | 205           | 55.6 | 47              | 58.8 | 138                    | 51.1 |
| <24                      | 100           | 27.1 | 26              | 32.5 | 77                     | 28.5 |
| >59                      | 64            | 17.3 | 7               | 8.8  | 55                     | 20.4 |
| <b>Age categories</b>    |               |      |                 |      |                        |      |
| ≤19                      | 1             | 0.3  | -               | -    | -                      | -    |
| 20–34                    | 270           | 73.2 | 63              | 78.8 | 178                    | 65.9 |
| ≥35                      | 98            | 26.6 | 17              | 21.3 | 92                     | 34.1 |
| <b>Education</b>         |               |      |                 |      |                        |      |
| None                     | 4             | 1.1  | -               | -    | 1                      | 0.4  |
| Primary                  | 206           | 56.1 | 40              | 50.0 | 112                    | 41.5 |
| Secondary and above      | 157           | 42.8 | 40              | 50.0 | 157                    | 58.1 |
| <b>Occupation</b>        |               |      |                 |      |                        |      |
| Employed                 | 273           | 74.2 | 49              | 61.3 | 217                    | 80.4 |
| Unemployed               | 95            | 25.8 | 31              | 38.8 | 53                     | 19.6 |
| <b>Religion</b>          |               |      |                 |      |                        |      |
| Catholic                 | 144           | 39.1 | 29              | 36.3 | 82                     | 30.4 |
| Protestant               | 151           | 41   | 36              | 45   | 124                    | 45.9 |
| Muslim                   | 72            | 19.6 | 15              | 18.8 | 64                     | 23.7 |
| Others                   | 1             | 0.3  | -               | -    | -                      | -    |
| <b>Marital status</b>    |               |      |                 |      |                        |      |
| Married                  | 352           | 96.4 | 75              | 94.9 | 261                    | 97   |
| Single                   | 13            | 3.6  | 4               | 5.1  | 8                      | 3    |
| <b>Current residence</b> |               |      |                 |      |                        |      |
| Rural                    | 142           | 38.5 | 29              | 36.3 | 69                     | 25.6 |
| Urban                    | 209           | 56.6 | 45              | 56.3 | 188                    | 69.6 |
| Semi urban               | 18            | 4.9  | 6               | 7.5  | 13                     | 4.8  |

<https://doi.org/10.1371/journal.pone.0228330.t004>

standard errors, which gives our study greater strength. Although Conde-Agudelo [6] found no statistical significance in association of IPI on PPH in their study, the direction of effect of IPI was in agreement with this study.

Although the results in this study are inconsistent with other studies that precede it, we have reason to believe that the characteristics of women in this study could have contributed to and could explain the uncommon results reported in this study. KCMC is a zonal referral hospital and women delivering here are in the most part, high risk-pregnant women. Being a referral hospital, the health care workers are hyper vigilant and more equipped to address complications that could arise during pregnancy and child birth. This means that the type of care offered to high risk-pregnant women could be different from the type of care that these women could have received in lower level facilities and hence prevent these women from complications such as PPH and Pre-eclampsia early on. This difference in care could be the reason why such associations are found to be protective.

The yearly increase in median IPI suggests an improvement during the early years of the study. This initial increase is likely to be influenced by increase in proportion of women using modern contraceptive, an almost constant total fertility rate from 1999 to 2005 and an increase in wanted fertility from 36% in 1999 to 42% in 2005 in Tanzania [31] The plateau from 2006 to 2014 implies changes in contraceptive use patterns. Although there was an increase in the

Table 5. Health-related characteristics of women by adverse maternal outcomes (N = 692).

| Variable                                   | PPH (n = 369) |      | Anemia (n = 80) |       | Pre-eclampsia (n = 270) |      |
|--|---------------|------|-----------------|-------|-------------------------|------|
|  | n             | %    | n               | %     | n                       | %    |
| <b>Type of family planning used</b>        |               |      |                 |       |                         |      |
| None                                       | 33            | 9.1  | 10              | 12.5  | 25                      | 9.4  |
| Modern                                     | 259           | 71.2 | 45              | 56.3  | 165                     | 61.8 |
| Traditional                                | 72            | 19.8 | 25              | 31.3  | 77                      | 28.8 |
| <b>Alcohol use before/during pregnancy</b> | 108           | 29.3 | 24              | 30    | 67                      | 24.8 |
| <b>PPH in preceding birth</b>              | 22            | 6    | 1               | 1.3   | 13                      | 4.8  |
| <b>Anemia in preceding birth</b>           | 9             | 2.4  | 6               | 7.5   | 5                       | 1.9  |
| <b>Preeclampsia in preceding birth</b>     | 16            | 4.3  | 5               | 6.3   | 70                      | 25.9 |
| <b>Referred from</b>                       |               |      |                 |       |                         |      |
| Home                                       | 338           | 91.6 | 75              | 94.9  | 237                     | 90.1 |
| Regional hospital                          | 9             | 2.4  | -               | -     | 9                       | 3.4  |
| District hospital                          | 9             | 2.4  | 2               | 2.5   | 7                       | 2.7  |
| Other                                      | 13            | 3.5  | 2               | 2.5   | 10                      | 3.8  |
| <b>Delivery mode</b>                       |               |      |                 |       |                         |      |
| Spontaneous                                | 50            | 13.6 | 39              | 48.8  | 115                     | 42.8 |
| Assisted                                   | 2             | 0.5  | 1               | 1.3   | 7                       | 2.6  |
| CS   | 317           | 85.9 | 40              | 50.0  | 147                     | 54.6 |
| <b>Death of preceding birth</b>            | 31            | 8.4  | 3               | 3.8   | 38                      | 14.1 |
| <b>Parity</b>                              |               |      |                 |       |                         |      |
| 2  | 135           | 36.6 | 43              | 53.8  | 86                      | 31.9 |
| 3  | 120           | 32.5 | 14              | 17.5  | 90                      | 33.3 |
| >3   | 114           | 30.9 | 23              | 28.7  | 94                      | 34.8 |
| <b>Antenatal care attendance</b>           | 367           | 99.5 | 80              | 100.0 | 269                     | 99.6 |
| <b>Number of ANC visits</b>                |               |      |                 |       |                         |      |
| <4   | 133           | 36.0 | 27              | 33.8  | 97                      | 35.9 |
| ≥4   | 231           | 62.6 | 51              | 63.8  | 169                     | 62.6 |
| Missing                                    | 5             | 1.4  | 2               | 2.5   | 4                       | 1.5  |

<https://doi.org/10.1371/journal.pone.0228330.t005>






Fig 2. Trend of inter-pregnancy interval over time. The red dotted line shows a demarcation for IPI at 24 and 59 months.

<https://doi.org/10.1371/journal.pone.0228330.g002>



**Fig 3. Distribution of inter-pregnancy interval by birth order.** The red dotted line shows a demarcation for IPI at 24 and 59 months.

<https://doi.org/10.1371/journal.pone.0228330.g003>



**Fig 4. Distribution of inter-pregnancy by age-group.** The red dotted line shows a demarcation for IPI at 24 and 59 months.

<https://doi.org/10.1371/journal.pone.0228330.g004>

Table 6. Crude association between inter-pregnancy interval and other confounders with adverse maternal outcomes.

| VARIABLE                                   | PPH (N = 7,360) |                  | ANAEMIA (N = 7,995) |              | PRE-ECLAMPSIA (N = 7,995) |                   |
|--|-----------------|------------------|---------------------|--------------|---------------------------|-------------------|
|  | cOR             | P-value          | cOR                 | P-value      | cOR                       | P-value           |
| <b>IPI Categories</b>                      |                 |                  |                     |              |                           |                   |
| 24–59                                      | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| <24  | 0.82            | 0.12             | 0.82                | 0.407        | 0.82                      | 0.174             |
| >59  | 0.77            | 0.081            | 0.39                | <b>0.02</b>  | 1.05                      | 0.767             |
| <b>Age (yrs)</b>                           |                 |                  |                     |              |                           |                   |
| <20  | 1.00            |                  |                     |              |                           |                   |
| 20–34                                      | 2.49            | 0.368            |                     |              |                           |                   |
| >34  | 2.97            | 0.284            |                     |              |                           |                   |
| <b>Mother's education</b>                  |                 |                  |                     |              |                           |                   |
| No formal education                        | 1.00            |                  |                     |              | 1.00                      |                   |
| Primary                                    | 0.91            | 0.851            |                     |              | 2.11                      | 0.461             |
| Secondary and above                        | 0.76            | 0.601            |                     |              | 3.51                      | 0.214             |
| <b>Mother's occupation</b>                 |                 |                  |                     |              |                           |                   |
| Employed                                   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Unemployed                                 | 1.13            | 0.302            | 1.93                | <b>0.005</b> | 0.73                      | <b>0.05</b>       |
| <b>Marital status</b>                      |                 |                  |                     |              |                           |                   |
| Married                                    | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Single                                     | 1.26            | 0.421            | 1.78                | 0.265        | 1.01                      | 0.969             |
| <b>Mother's current residence</b>          |                 |                  |                     |              |                           |                   |
| Rural                                      | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Urban                                      | 0.71            | <b>0.003</b>     | 0.80                | 0.36         | 1.42                      | <b>0.019</b>      |
| Semi urban                                 | 1.02            | 0.942            | 1.64                | 0.275        | 1.51                      | 0.236             |
| <b>Type of pregnancy</b>                   |                 |                  |                     |              |                           |                   |
| Multiple                                   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Singleton                                  | 0.35            | <b>&lt;0.001</b> | 0.29                | <b>0.01</b>  | 0.46                      | <b>0.014</b>      |
| <b>Type of family planning used</b>        |                 |                  |                     |              |                           |                   |
| None                                       | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Modern                                     | 0.91            | 0.603            | 0.55                | 0.088        | 0.80                      | 0.323             |
| Traditional                                | 0.78            | 0.259            | 0.90                | 0.785        | 1.11                      | 0.632             |
| <b>Alcohol use before/during pregnancy</b> |                 |                  |                     |              |                           |                   |
| No   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Yes  | 1.05            | 0.673            | 1.01                | 0.96         | 0.77                      | <b>0.08</b>       |
| <b>PPH in preceding birth</b>              |                 |                  |                     |              |                           |                   |
| No   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Yes  | 1.62            | <b>0.034</b>     | 0.34                | 0.277        | 1.38                      | 0.269             |
| <b>Anemia in preceding birth</b>           |                 |                  |                     |              |                           |                   |
| No   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Yes  | 1.64            | 0.156            | 5.27                | <b>0.001</b> | 1.19                      | 0.712             |
| <b>Preeclampsia in preceding birth</b>     |                 |                  |                     |              |                           |                   |
| No   | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Yes  | 1.24            | 0.409            | 1.73                | 0.238        | 11.50                     | <b>&lt;0.0001</b> |
| <b>Referred from</b>                       |                 |                  |                     |              |                           |                   |
| Home                                       | 1.00            |                  |                     |              | 1.00                      |                   |
| Regional hospital                          | 1.59            | 0.185            |                     |              | 1.98                      | 0.052             |
| District hospital                          | 2.03            | <b>0.045</b>     |                     |              | 2.14                      | 0.055             |
| Other                                      | 1.89            | <b>0.031</b>     |                     |              | 2.09                      | 0.027             |
| <b>Delivery mode</b>                       |                 |                  |                     |              |                           |                   |
| Spontaneous                                | 1.00            |                  | 1.00                |              | 1.00                      |                   |
| Assisted                                   | 2.47            | 0.215            | 1.45                | 0.7          | 3.72                      | <b>0.001</b>      |
| CS   | 11.98           | <b>&lt;0.001</b> | 1.73                | <b>0.017</b> | 2.20                      | <b>&lt;0.0001</b> |
| <b>Death of preceding birth</b>            |                 |                  |                     |              |                           |                   |
| No   | 1.00            |                  | 1.00                |              | 1.00                      |                   |

(Continued)

Table 6. (Continued)

| VARIABLE                         | PPH (N = 7,360) |         | ANAEMIA (N = 7,995) |         | PRE-ECLAMPSIA (N = 7,995) |         |
|----------------------------------|-----------------|---------|---------------------|---------|---------------------------|---------|
|                                  | cOR             | P-value | cOR                 | P-value | cOR                       | P-value |
| Yes                              | 2.16            | <0.001  | 0.78                | 0.679   | 3.50                      | <0.0001 |
| <b>Parity</b>                    |                 |         |                     |         |                           |         |
| 2                                | 1.00            |         | 1.00                |         | 1.00                      |         |
| 3                                | 1.29            | 0.047   | 0.47                | 0.013   | 1.54                      | 0.003   |
| >3                               | 1.67            | <0.001  | 1.02                | 0.923   | 2.15                      | <0.0001 |
| <b>Antenatal care attendance</b> |                 |         |                     |         |                           |         |
| No                               |                 |         |                     |         | 1.00                      |         |
| Yes                              |                 |         |                     |         | 1.14                      | 0.896   |
| <b>Number of ANC visits</b>      |                 |         |                     |         |                           |         |
| <4                               | 1.00            |         | 1.00                |         | 1.00                      |         |
| ≥4                               | 1.08            | 0.505   | 1.13                | 0.603   | 1.04                      | 0.784   |

cOR = Crude Odds Ratio

<https://doi.org/10.1371/journal.pone.0228330.t006>

proportion of women using modern contraceptives from 2005 to 2015, the greatest increase was noted in use of short term contraceptives [21] which are not as effective as long term contraceptives in affecting IPI[32]. This could explain the plateau. The rise seen in 2015 could suggest that, trends could be moving towards longer IPI. This could be due to re-launching of Green Star Campaign, a programmatic change made at the end of 2013 to promote the use of modern family planning methods, which could have influenced contraceptive use patterns [22]. The reported trend of median IPI from 2002 to 2015 in this study differs from that reported in the country [21] and this difference could be due to the fact that a survey reports a summary measure for all regions, therefore the regional differences are masked.

### Conclusion

There are still many women delivering within sub-optimal IPI (51.7%) in KCMC. Most women who have short IPI are young, less than 20years. This study has found a protective association between sub-optimal IPI with adverse maternal outcomes. However, these results

Table 7. Crude and adjusted association between inter-pregnancy interval and adverse maternal outcomes.

| Outcome                               | Inter-pregnancy interval categories (months) |                          |              |                          |                          |
|---------------------------------------|--|--------------------------|--------------|--------------------------|--------------------------|
|                                       | <24  | P-value                  | 24–59 (Ref.) | 60+                      | P-value                  |
| Pre-eclampsia (N = 7995)              | 0.82 (0.62, 1.09)                            | 0.174                    | 1.00         | 1.05 (0.76, 1.44)        | 0.767                    |
| Pre-eclampsia <sup>a</sup> (N = 7749) | <b>0.71 (0.52, 0.97)</b>                     | <b>0.032<sup>†</sup></b> | 1.00         | 0.92 (0.65, 1.27)        | 0.603                    |
| PPH (N = 7360)                        | 0.82 (0.65, 1.05)                            | 0.120                    | 1.00         | 0.77 (0.58, 1.03)        | 0.081                    |
| PPH <sup>a</sup> (N = 7304)           | 0.88 (0.68, 1.13)                            | 0.311                    | 1.00         | <b>0.70 (0.52, 0.94)</b> | <b>0.019<sup>†</sup></b> |
| Anemia (N = 7995)                     | 0.82 (0.51, 1.31)                            | 0.407                    | 1.00         | 0.39 (0.18, 0.86)        | <b>0.020</b>             |
| Anemia <sup>a</sup> (N = 7906)        | 0.72 (0.45, 1.17)                            | 0.184                    | 1.00         | 0.45 (0.20, 1.03)        | 0.059                    |

<sup>a</sup>adjusted for: age, mother’s occupation, religion, current residence, alcohol use, pre-eclampsia in previous pregnancy, PPH in previous pregnancy, anemia in previous pregnancy, delivery mode, perinatal death of previous child at birth, parity, number of ANC visits, ANC attendance, type of family planning used, pregnancy type and referral status.

<sup>†</sup>Significant p-value

<https://doi.org/10.1371/journal.pone.0228330.t007>

do not nullify the previous literature as they could be biased by the demographics of these women as well as the tertiary care they are receiving

## Acknowledgments

We would like to extend our sincere gratitude to Medical Birth Registry staff and staff at the Department of Obstetrics and Gynecology at Kilimanjaro Christian Medical Centre for working tirelessly in collecting and updating the data for the registry and the KCMC management for allowing the use of Birth Registry data for this manuscript. We also thank the Norwegian government for supporting the establishment of the KCMC Medical Birth Registry. Also to the entire Institute of Public Health at Kilimanjaro Christian Medical University College for their valuable inputs and encouragement, and our colleagues Fredrick Sigalla, Umami Abdul, Emmanuel Mtui, Nathaniel Kalengo and Ephrasia Hugho for their advice and moral support.

## Author Contributions

**Conceptualization:** Leah Anku Sanga, Michael Johnson Mahande.

**Data curation:** Leah Anku Sanga.

**Formal analysis:** Leah Anku Sanga.

**Methodology:** Leah Anku Sanga, Michael Johnson Mahande.

**Project administration:** Michael Johnson Mahande.

**Supervision:** Tara Mtuy, Rune Nathaniel Philemon, Michael Johnson Mahande.

**Visualization:** Leah Anku Sanga, Tara Mtuy, Rune Nathaniel Philemon, Michael Johnson Mahande.

**Writing – original draft:** Leah Anku Sanga.

**Writing – review & editing:** Leah Anku Sanga, Tara Mtuy, Rune Nathaniel Philemon, Michael Johnson Mahande.

## References

1. WHO. Report of a WHO Technical Consultation on Birth Spacing [Internet]. Geneva, Switzerland; 2007. Available from: [http://apps.who.int/iris/bitstream/10665/69855/1/WHO\\_RHR\\_07.1\\_eng.pdf%0Ahttp://www.who.int/maternal\\_child\\_adolescent/documents/birth\\_spacing.pdf](http://apps.who.int/iris/bitstream/10665/69855/1/WHO_RHR_07.1_eng.pdf%0Ahttp://www.who.int/maternal_child_adolescent/documents/birth_spacing.pdf)
2. Norton M. New evidence on birth spacing: promising findings for improving newborn, infant, child, and maternal health. *Int J Gynecol Obstet* [Internet]. 2005 Apr 1 [cited 2018 May 18]; 89:S1–6. Available from: <http://doi.wiley.com/10.1016/j.ijgo.2004.12.012>
3. Wendt A, Gibbs CM, Peters S, Hogue CJ. Impact of increasing inter-pregnancy interval on maternal and infant health. *Paediatr Perinat Epidemiol*. 2012; 26(SUPPL. 1):239–58.
4. Defranco EA, Seske LM, Greenberg JM, Muglia LJ. Influence of interpregnancy interval on neonatal morbidity. *Am J Obstet Gynecol*. 2015; 212(3):386.e1–386.e9.
5. Rutstein SO. TRENDS IN BIRTH SPACING [Internet]. DHS Comparative Reports. Calverton, Maryland, USA; 2011 [cited 2018 Jul 4]. Available from: <https://dhsprogram.com/pubs/pdf/CR28/CR28.pdf>
6. Conde-Agudelo A. Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study. *BMJ* [Internet]. 2000; 321(7271):1255–9. Available from: <http://www.bmj.com/cgi/doi/10.1136/bmj.321.7271.1255> PMID: 11082085
7. Lilungulu A, Matovelo D, Kihunrwa A, Gumodoka B. Spectrum of maternal and perinatal outcomes among parturient women with preceding short inter-pregnancy interval at Bugando Medical Centre, Tanzania. *Matern Heal Neonatol Perinatol* [Internet]. 2015; 1(1):1. Available from: <http://mhnjournal.biomedcentral.com/articles/10.1186/s40748-014-0002-1>

8. Seham FA., Ali Abd El Hamid A, Fathy Heiba M, Mohamed Elalem O. Effect of Interpregnancy Interval on Pregnancy Outcome Among Pregnant Women Attending Delivery At Belqas hospital. *IOSR J Nurs Heal Sci* [Internet]. 2015; 4(4):2320–1940. Available from: [www.iosrjournals.org](http://www.iosrjournals.org)
9. Habimana-Kabano I, Broekhuis A, Hooimeijer P. Inter-Pregnancy Intervals and Maternal Morbidity: New Evidence from Rwanda. *Afr J Reprod Health*. 2015; 19(3):77–86. PMID: 26897916
10. Hanley GE, Hutcheon JA, Kinniburgh BA, Lee L. Interpregnancy interval and adverse pregnancy outcomes: an analysis of successive pregnancies. *Obstet Gynecol*. 2017; 129(3):408–15. <https://doi.org/10.1097/AOG.0000000000001891> PMID: 28178044
11. USAID. STRENGTHENING FAMILY PLANNING POLICIES AND PROGRAMS IN DEVELOPING COUNTRIES: AN ADVOCACY TOOLKIT USAID [Internet]. Washington, DC; 2005 [cited 2018 May 15]. Available from: <http://www.policyproject.com/pubs/manuals/FamilyPlanningToolkitfinal.pdf>
12. Conde-Agudelo A, Rosas-Bermudez A, Castaño F, Norton MH. Effects of Birth Spacing on Maternal, Perinatal, Infant, and Child Health: A Systematic Review of Causal Mechanisms. 2012 [cited 2018 Mar 7]; 43(2). Available from: [https://www.k4health.org/sites/default/files/conde-agudelo\\_2012.pdf](https://www.k4health.org/sites/default/files/conde-agudelo_2012.pdf)
13. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: A WHO systematic analysis. *Lancet Glob Heal*. 2014; 2(6):323–33.
14. WHO. Maternal mortality [Internet]. World Health Organization; 2016 [cited 2018 Mar 17]. Available from: <http://www.who.int/news-room/fact-sheets/detail/maternal-mortality>
15. UN. THE GLOBAL STRATEGY FOR WOMEN'S, CHILDREN'S AND ADOLESCENTS' HEALTH (2016–2030) [Internet]. New York, USA; 2015 [cited 2018 Mar 7]. Available from: [http://globalstrategy.everywomaneverychild.org/pdf/EWEC\\_globalstrategyreport\\_200915\\_FINAL\\_WEB.pdf](http://globalstrategy.everywomaneverychild.org/pdf/EWEC_globalstrategyreport_200915_FINAL_WEB.pdf)
16. WHO, UNICEF, UNFPA, World Bank, United Nations. TRENDS IN MATERNAL MORTALITY: 2000 TO 2017. 2019.
17. WHO. Trends in Maternal Mortality: 1990 to 2008. 2008; Available from: [http://whqlibdoc.who.int/publications/2010/9789241500265\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241500265_eng.pdf)
18. WHO. Maternal mortality Fact Sheet. 2016;
19. MoHSW. The National Road Map Strategic Plan To Accelerate Reduction of Maternal, Newborn and Child Deaths in Tanzania. 2008;(April):1–102.
20. MoHSW. The National Road Map Strategic Plan to Improve Reproductive, Maternal, Newborn, Child & Adolescent Health. Dar es Salaam, Tanzania; 2015.
21. MoHCDGEC. Tanzania Demographic and Health Survey and Malaria Indicator Survey (TDHS-MIS) 2015–16 [Internet]. Vol. MoHCDGEC, Dar es Salaam, Tanzania, and Rockville, Maryland USA. Dar es Salaam, Tanzania and Rockville, Maryland, USA; 2016. Available from: <https://www.dhsprogram.com/pubs/pdf/FR321/FR321.pdf>
22. MoHSW. National Family Planning Costed Implementation Program [Internet]. Dar es Salaam, Tanzania; 2010 [cited 2018 Jul 20]. Available from: <https://www.fhi360.org/sites/default/files/media/documents/national-fp-costed-implementation-plan-tanzania-main-text.pdf>
23. Pile JM, Simbakalia C. Repositioning Family Planning-Tanzania Case Study: A Successful Program Loses Momentum [Internet]. New York; 2006 [cited 2018 Jul 20]. Available from: [www.acquireproject.org](http://www.acquireproject.org)
24. Afnan-Holmes H, Magoma M, John T, Levira F, Msemu G, Armstrong CE, et al. Tanzania's Countdown to 2015: an analysis of two decades of progress and gaps for reproductive, maternal, newborn, and child health, to inform priorities for post-2015. *Lancet Glob Heal* [Internet]. 2015 [cited 2018 Mar 7]; 3: e396–409. Available from: [http://www.thelancet.com/pdfs/journals/langlo/PIIS2214-109X\(15\)00059-5.pdf](http://www.thelancet.com/pdfs/journals/langlo/PIIS2214-109X(15)00059-5.pdf)
25. de Jonge HCC, Azad K, Seward N, Kuddus A, Shaha S, Beard J, et al. Determinants and consequences of short birth interval in rural Bangladesh: a cross-sectional study. *BMC Pregnancy Childbirth* [Internet]. 2014 Dec 24 [cited 2018 Mar 10]; 14:427. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25539669> <https://doi.org/10.1186/s12884-014-0427-6> PMID: 25539669
26. WHO. WHO | SDG 3: Ensure healthy lives and promote wellbeing for all at all ages [Internet]. World Health Organization; 2017 [cited 2018 Jun 29]. Available from: <http://www.who.int/sdg/targets/en/>
27. NBS, ICF Macro. Tanzania Demographic and Health Survey 2010 [Internet]. Dar es Salaam, Tanzania; 2011. Available from: <https://dhsprogram.com/pubs/pdf/FR243/FR243%5B24June2011%5D.pdf>
28. Exavery A, Mrema S, Shamte A, Bietsch K, Mosha D, Mbaruku G, et al. Levels and correlates of non-adherence to WHO recommended inter-birth intervals in Rufiji, Tanzania. *BMC Pregnancy Childbirth* [Internet]. 2012; 12:1–8. Available from: <http://www.biomedcentral.com/1471-2393/12/152> <https://doi.org/10.1186/1471-2393-12-1>
29. Begna Z, Assegid S, Kassahun W, Gerbaba M. Determinants of inter birth interval among married women living in rural pastoral communities of southern Ethiopia: a case control study. *BMC Pregnancy*

Childbirth [Internet]. 2013 Dec 20 [cited 2018 Mar 11]; 13(116). Available from: <http://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/1471-2393-13-116>

30. Illah E, Mbaruku G, Masanja H, Kahn K. Causes and risk factors for maternal mortality in rural Tanzania—case of Rufiji Health and Demographic Surveillance Site (HDSS). *Afr J Reprod Health* [Internet]. 2013; 17(3):119–30. Available from: [https://www.researchgate.net/publication/257133413\\_Causes\\_and\\_risk\\_factors\\_for\\_maternal\\_mortality\\_in\\_rural\\_Tanzania—case\\_of\\_Rufiji\\_Health\\_and\\_Demographic\\_Surveillance\\_Site\\_HDSS](https://www.researchgate.net/publication/257133413_Causes_and_risk_factors_for_maternal_mortality_in_rural_Tanzania—case_of_Rufiji_Health_and_Demographic_Surveillance_Site_HDSS) PMID: 24069774
31. NBS, ORC Macro. Tanzania Demographic and Health Survey 2004–05. Dar es Salaam, Tanzania; 2005.
32. Dutta DC. Textbook of Obstetrics including Perinatology and Contraception. Eighth. Konar H, editor. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd; 2015.