

Development, implementation and early results of a 12-week web-based intervention targeting 51 children age 5–13 years and their families

Annelie Thorén^{1,2}  | Annika Janson^{3,4} | Erling Englund⁵ | Sven-Arne Silfverdal¹ 

¹Department of Clinical Sciences, Pediatrics, Umeå University, Umeå, Sweden

²Department of Pediatrics, Sollefteå Hospital, Sollefteå, Sweden

³National Childhood Obesity Centre, Karolinska University Hospital, Stockholm, Sweden

⁴Division of Pediatric Endocrinology, Department of Women's and Children's Health Karolinska Institutet, Stockholm, Sweden

⁵Department of Research and Development, Västernorrland County Council, Sundsvall, Sweden

Correspondence

Annelie Thorén, Department of Pediatrics, Sollefteå Hospital, Child Health Care Unit, Västernorrland County Council, Sollefteå, SE 881 33, Sweden.

Email: annelie.thoren@rvn.se

Funding information

VINNOVA: 2018-01405

Summary

Background: Internet-based treatments have proven effective for various health issues. There is a need to scale up interventions targeting children with obesity, also in less densely populated areas where the prevalence in many countries is higher than in urban areas. The aim of this study was to design and implement an internet-based program as an add-on to standard treatment for childhood obesity.

Methods: Web-Childhood Obesity Prevention (Web-COP) was a prospective feasibility study with a pre- post- design. The intervention consisted of four group-based education sessions at the clinic, physical activity on prescription, and a new 12-week internet-based program. Web-COP was offered to children with obesity (International Obesity Task Force Body Mass Index (IOTF-BMI) ≥ 30 kg/m²) and their parents in two counties in Northern Sweden from August 2018 to June 2019. The primary outcome was change in BMI standard deviation score (BMI-SDS).

Results: The study included 55 children 5–13 years of age. The internet-based component was well received, and retention rate was 51/55 (92.7%). Data was analysed for 51 children. Mean BMI-SDS was 3.3 at start and decreased by 0.2, 0.3, and 0.4 at two, four, and six months from baseline. Using a continuous algorithm, 42/51 (81%), children lowered their BMI-SDS and 33/51 (65%) lowered their BMI.

Conclusion: Adding group sessions and an internet-based program to standard care was feasible and two thirds of included children with obesity reduced their BMI.

KEYWORDS

children, intervention, obesity, web-based treatment

1 | INTRODUCTION

Childhood obesity is a growing global health concern.¹ Obesity in childhood is likely to track into adulthood.^{2–4} Longitudinal data from

children born 1997–1999 in Sweden show that children with overweight or obesity at five years of age tend to remain living with overweight or obesity later in childhood.⁵ However, there is also considerable new development of obesity.^{2,4,6} Early interventions

Abbreviations: BMI, Body Mass Index; IOTF, International Obesity Task Force; Web-COP, Web-Childhood Obesity Prevention.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. Obesity Science & Practice published by World Obesity and The Obesity Society and John Wiley & Sons Ltd

targeting children with obesity 5–13 years of age serve as attractive opportunities for promoting a healthy lifestyle aiming at reducing future risk of obesity and associated health risks.

In data from Sweden's three largest cities, 9–16% of four-year-old children have overweight, and another 2–4% have obesity.^{7–9} The previous rapid increase in the prevalence of childhood obesity in Sweden may have halted.¹⁰ The prevalence of childhood and adult obesity is higher in the less densely populated northern parts of Sweden than in other parts of the country.^{11,12} A decline in childhood overweight and obesity has been noticed also in rural areas. In a study on four-year-old children living in Northern Sweden the prevalence of overweight among boys fell in five years from 17.2% to 14.2% and among girls from 22.3% to 19.0%.¹³

Treatments of childhood obesity emphasize lifestyle modification, that is, restrict calorie intake and increase energy expenditure by increased physical activity. Evidence suggests that multi-component interventions that aim at stimulating changes in behaviour may achieve small reductions in body weight status in children.^{14–16} Also, previous studies have shown that involving parents is beneficial in child obesity treatment.^{17–21} The benefit of parental involvement is likely multi-factorial, from being a role model and provide psychological support to the practical parental support that is pivotal for children to change dietary habits by providing healthy meals and limit exposure to unhealthy foods, promote physical exercise, and be able to travel and join treatment programs.^{13,22} Many previous trials on childhood obesity treatment report high levels of drop-out or loss to follow-up, with drop-out rates ranging from 40% to 79%.^{13,16,22} In less densely populated areas, the distance to treatment centers may further limit the ability for children to participate in obesity treatment programs.

Web-based treatments in various health-related areas have demonstrated to be effective with low risks. Using the internet for communication with patients has promoted patient satisfaction, coverage of treatment, and cost-effectiveness.²³ Weight loss programs delivered over the internet could be especially appealing to children and adolescents.²⁴ Inherent attributes of web-based interventions, such as interactivity, flexibility in time and location, and anonymity, which may limit obstacles due to stigma related to obesity, may be beneficial.^{24,25} Still, there is a need to verify that web-based interventions help to sustain long-term behavioural change. In a review of twelve studies describing eight randomized web-based interventions, four of the studies showed a reduction of obesity in the different measurements of obesity used, whereas two studies showed no results. In two studies, the BMI increased at follow-up.²⁶ More than a decade ago, a randomized controlled study of a comprehensive one-year intervention promoting lifestyle changes in overweight and obese children age 8–12 years in Northern Sweden showed modest effects,¹⁶ but, interestingly, the study had an internet-based follow-up during the second year. The authors concluded that families were not comfortable with the internet-based part of the program and the platform was not much used.²⁷

In the area of this study, standard care treatment for children with obesity comprises a doctor's appointment and, in some cases, contact with a dietician. This study, the Web-Childhood Obesity Prevention

(Web-COP) study, offered an add-on to existing standard care. The study aimed at designing a web-based module and investigating whether the addition of a web-based module was feasible from a technical and practical point before planning for a randomized controlled study. More specifically, the study wanted to assess if the various means of recruitment of patients would work, if there were technical issues on the use of the internet-based modules, and if internet based weekly modules could assist by supporting other parts of the program and increase the participation of the families, limit drop-outs and increase the frequency of treatment contacts, without the family needing to attend the clinic. The hypothesis was that school nurses would be able to recruit patients that had not previously been in treatment for obesity, that the parents and families would use and find the new internet-based program attractive, that children would remain in treatment. This paper aims to present the design and feasibility aspects of patient recruitment, internet-based activities, and retention rate, as well as, changes in BMI-SDS at two, four, and six months after inclusion.

2 | MATERIALS AND METHODS

2.1 | Study area, recruitment, and study population

The study took place in Västernorrland Region in the north of Sweden. In this area, 98% of the population has access to high-speed internet.²⁸ Families from both urban and rural areas, including parents living in separate homes, were invited to participate in the study if interested and eligible. Participants were recruited in cycles 2018/2019, with two groups starting the program in August 2018 and one group starting in January 2019. A variety of recruitment strategies were utilized, including newspapers, study pamphlets, and direct contact (AT) with school nurses. Children and parents were invited to the paediatric health care clinics at the hospitals in Sollefteå and Sundsvall for a baseline visit and measures of weight, height, and waist circumference prior to the first group session.

Inclusion criteria were obesity as defined by the International Obesity Task Force (IOTF) cutoff IOTF-BMI ≥ 30 ,^{29,30} 5–13 years of age, families able to speak and write Swedish, and having a high-speed internet connection in their homes. Previous obesity treatment was not an exclusion criterion.²⁵ Syndromic obesity, or a major neurological or endocrine disease, were exclusion criteria. A modification of the protocol extending the upper age limit to 13 years was made to enable five children age 13 years to participate.

2.2 | Study design

This was a two-center, single-group, non-randomized, prospective feasibility study with a pre-test and post-test design aimed at evaluating the feasibility of the new intervention, including educational group sessions, and a web-based treatment program. The Web-COP study consisted of three components: 1) a series of four educational group




FIGURE 1 The timeline of the Web-COP program

sessions, 2) a 12-week web-based program, including coaching, and 3) physical activity on prescription³¹ (Figure 1).

2.3 | Group sessions

After the initial inclusion visit to the paediatrician, group sessions were given weekly over a four-week-period to parents and children. The group sessions took place at an exercise center in each community. Each session lasted 120 minutes and was attended by the child and at least one parent. Parents and children met in separate groups for physical activity and nutrition lessons. At all sessions, the children engaged in physical exercise for 45–60 min. Children were given the opportunity to try various sports like outdoor training, swimming, Tabata, and circuit training. Meanwhile, parents were given education in nutrition and physical activity.

Two experienced members of staff, a dietician and a physical therapist, led the classes and encouraged discussions on different

topics. Lessons on nutrition focused on the importance of increasing fruit and vegetables, avoiding food high in sugar, and introducing *my plate* (Tallriksmodellerna) to focus both on meal size and meal composition. These lessons were provided to children through interactive activities. Parents and children also had the opportunity to prepare one simple healthy meal together, and parents were offered a tour with a dietician in a grocery store to learn more about healthy and unhealthy food choices. Between the meetings, children and parents had home assignments related to the theme of the meeting. In one group session, the parents were provided instructions on how to use the internet-based program and were given technical support. The outline of the internet-based program content was delivered verbally to the group supplemented with a slide presentation. Printed material was provided to the parents during two sessions.

2.4 | Internet-based program

The mobile-phone friendly internet-based program was designed for parents to support their children's obesity treatment. The internet-based program was developed by a multidisciplinary team of researchers and experts in nutrition, paediatric medicine, engineering, child psychology, web-design, and physical activity. Gamification, application of game-design elements, was used to support families in achieving positive health behaviour goals. The parents received a personal login to the internet-based program. For 12 weeks, the parents participated in a program consisting of online lessons with different topics. Parents could complete the program at their own pace. Each week ended with a test for the participating parent or family. After finishing each week, the parents received a short message to praise the parent for a completed week and to encourage them to start the next week in the program.

The twelve sessions aimed at encouraging healthy eating habits, such as choice of breakfast and reducing sugar intake, and gave examples on how to increase the amount of fruit and vegetables when planning a meal. In addition, the sessions aimed at motivating the family to be more physically active. The program also contained quizzes on healthy food choices and information about how to promote healthy sleeping and screen-time behaviour. If parents had not used the internet-based program for a couple of days, they received a reminder message via mail. Both parents were asked to participate in the web-based program.

2.5 | Physical activity on prescription

At the inclusion visit, the paediatrician discussed the importance of getting 60 minutes or more of physical activity per day with the children and parents and prescribed "physical activity on prescription". Physical activity on prescription is a Swedish structured national model that enables health care providers from different cadres to prescribe physical activity.³² The prescription enabled the children to get free access to activities from a local provider of the family's choice.

The younger children could choose between different physical activities, including family aerobics and swimming, while the older could choose spinning, gym weight training, aerobic circuit training, or swimming. Parents were encouraged to exercise with their children and to promote more daily activities together and time outdoors.

3 | ETHICS

Children and families were invited to participate and received information, both verbal and written. Verbal informed consent was received for the baseline measurements and written informed consent from both child and parents was received at the baseline doctor's appointment. The study was approved by the Ethics Board at Umeå University (Dnr 2018–113-31 M, Dnr 2018–461-32 M).

4 | DATA COLLECTION

Patient files at the clinic were used to determine if patients were previously treated for obesity. Patients who had no previous contact recorded for obesity were defined as “new” patients. Attendance was taken at all group sessions. The registration to use the internet-based program was recorded for each family, and weekly use was reported by the parents.

4.1 | Measurements

BMI was calculated by weight (kg) divided by height squared (m^2) and using growth references of the Swedish growth charts for numerical value of BMI-SDS.³³ Children's heights and weights were measured at the start of the intervention (baseline) and after two, four, and six months. Height was measured to the nearest 0.1 cm. Weight was measured with the children barefoot in light clothing the nearest 0.1 kg using a digital scale (Tanita Digital Scale). Data was collected in Sollefteå and Sundsvall separately, and the same paediatric nurse made all measurements of children in each clinic. Data were entered in the Swedish Childhood Obesity Treatment Register (BORIS), a national quality register supervised by the National Board of Health and Welfare in Sweden.

4.2 | Statistical analysis

As data on BMI-SDS were skewed and not normally distributed, data were log transformed. Standard descriptive statistics (geometric mean, as the analysis was based on log-transformed values and 95% confidence interval) were computed using the continuous algorithm for BMI-SDS using the Swedish growth references and growth charts.³³ A paired two-tailed t-test was used for significance testing with the geometric mean of the ratio of paired values as measures of

difference with a 95% confidence interval. The significance level was set at $p < 0.05$ (IBM SPSS Statistics, Version 25.0, IBM Corp, Armonk, NY, USA) was used for the analyses. Seven children have a missing value at one of the measure points. In these cases, imputations with *Last observation carried forward* (and similar if an observation was carried backward) or means of nearby values, was used. Of these, one child has a missing BMI-SDS at baseline, and it was replaced by the value at two months. Two children had a missing BMI-SDS at four months and it was replaced by the mean of two and six months. Four children had a missing BMI-SDS at six months, and it was replaced by the value at four months.

5 | RESULTS

A total of 59 children (31 girls) and their parents were recruited in two communities. Recruitment was primarily done by school nurses, but also children already in treatment at the paediatric clinic were asked if they would be interested in participating. Of included children, 28 (16 girls) had previously not been attending obesity treatment at the paediatric clinic or elsewhere. Since many persons were involved in recruiting, we were unable to determine how many families were informed about the project.

Three children were excluded because they passed 13.9 years of age at the start of the intervention, and one participant was excluded because of taking medical treatment for ADHD. Four left the study before four months. In total, 51/55 (92.7%) of included children (27 girls) aged 5–13 completed the intervention (Figure 2).

In 28 (54%) of 51 included children, at least one parent and child participated in all the four educational group sessions. At least one parent to each child registered in the web-based program and most parents were also active in the internet-based program. The four families that withdrew did so in the period of the group sessions, and their rationale for not continuing was that they had a long way to drive to the group sessions and did not see obesity in the child as a big problem. The remaining 51 children and their parents completed the group sessions and proceeded to support through the web-based program. All families completed the web-based part of the program.

Of the 51 children, 42 lowered their BMI-SDS during the six-month intervention. Four children had a higher BMI-SDS after the intervention, and data at six months are missing for four children. The mean BMI-SDS was reduced from 3.3 (95% CI: 3.0–3.6) at baseline to 2.9 (95% CI: 2.7–3.1) (geometric) measured at six months ($P < 0.001$), the ratio of six months and baseline: 0.88 (95% CI: 0.85–0.91), meaning that BMI SDS has decreased 12% since baseline. A decrease in BMI-SDS was shown already after two months (0.94 (95% CI: 0.92–0.96), $P < 0.001$). The children in Sollefteå had higher BMI-SDS at baseline compared with the children in Sundsvall, 3.5 (95% CI: 2.9–4.3) 3.2 (3.0–3.5), respectively. The reduction in BMI-SDS was greater for the children in Sollefteå. The reduction in both Sundsvall and Sollefteå were significant ($p < 0.001$, Sundsvall: 0.91 (95% CI: 0.88–0.94) and Sollefteå: 0.82 (95% CI: 0.77–0.88)) (Figure 3).




FIGURE 2 Web-COP CONSORT




FIGURE 3 Web-COP BMI-SDS

6 | DISCUSSION

This study, the Web-COP program, was developed as an add-on to standard care and aimed at increasing parents' awareness, motivation, and ability to modify habits affecting their children's weight and health. The three components of the program were meant to

reinforce each other: group sessions, a web-based program, and organized physical activity free of charge. This was a feasibility study³⁴ with a focus on the study process, and we conclude that the recruitment process worked and that school nurses were effectively informing families about the study. The sex distribution of the included patients was balanced. The drop-outs occurred during the period of

group sessions only, and all remaining families were participating in the internet-based program. There were no major technical issues, and we observed no negative side-effects of the program. Data on changes in BMI-SDS were quite promising in relation to other treatments of childhood obesity.^{14,15,35–37}

With the high prevalence of childhood obesity there is a need to scale up interventions that are easily accessible and appreciated as helpful by the parents. This study included 24 (47%) children that were not previously in treatment. Also, the attrition rate (92.7%) was higher than in many programs targeting childhood obesity,¹ likely beneficial for the outcome in terms of BMI-SDS. With the aid of the child's parents, interventions should aim at diminishing the risks of persisting obesity and associated risks. Previous studies support the programs that target parents to children with obesity,^{19–21} but results on web-based treatments for treating childhood obesity were inconclusive. The authors of the review of twelve studies on using internet in obesity treatment concluded that controlled studies on stand-alone web-based treatments might elucidate their usefulness.²⁶ In contrast, our design was based on a presumption that web-based components would be useful as one of several components in an intervention. In the previously mentioned comprehensive one-year randomized controlled study aimed at promoting lifestyle changes in overweight and obese children age 8–12 years in Northern Sweden 2006–2009, the authors reported that the web-based support in the second year was not much used by the families and the drop-out rate of the total study was 45%.^{16,27,38} In our view, the increased usage of all kinds of internet applications over the last decade has largely contributed to the higher acceptability of the web-based component in this study, and likely also to the attrition rate.²⁶

The prevalence of overweight in children and their parents has shown to be higher in rural compared with urban areas.^{12,39} Moraes et al. have suggested that targeting rural areas might be an effective approach to reach children with the greatest needs.³⁹

Two-thirds of the children in this study reduced their degree of obesity. The difference in results between the two sites was most likely due to higher BMI-SDS at baseline in some children in Sollefteå compared with the children in Sundsvall. Previous studies have shown that even moderate changes in BMI among children can reduce risk indicators for cardiovascular disease.⁴⁰

Our findings suggest that future childhood obesity interventions may benefit from having components like Web-COP with group sessions supported by a web-based program. The inclusion of a mobile-friendly web-based program is consistent with recommendations of multi-component interventions to address the complexity of childhood obesity. Based on this feasibility study, further development of the Web-COP project will include more interactive game-like activities, less text in the web-based information, additional group sessions, and the children will be offered a smart-watch to track daily activity. To further study the effect of Web-COP, a randomized controlled trial with cross-over design on a larger number of children for a longer period of time was started in August 2019.

The strength of this study is that about 90% of the included patients completed the study, which is a high attrition rate for interventions on obesity. A limitation of the design was that families left the study during the period of group sessions, whereas they could have benefitted from the web-based intervention that came after the group-sessions. This will be considered for future studies. The major limitation of this study was its lack of randomization and control group.⁴¹ Also, we cannot determine which components of Web-COP were useful, just that the whole intervention achieved these results. Another limitation is the short-term follow-up. Also, the study could be subject to confounding factors due to selection bias as families were recruited for their willingness to participate. Since many persons were involved in the recruitment and the study was presented in schools and media, it was unfortunately impossible to determine how many families were introduced to the study and asked whether they wanted to participate. Recruiting patients based on their willingness to participate is a built-in limitation to the design of this study aimed at encouraging reflecting on habits, and sustainable behavioural change, which can only be achieved with the active involvement of the patients. More studies are necessary to confirm the evidence of combined group sessions and web-based family interventions.

7 | CONCLUSION

The Web-COP program with group sessions, an internet-based program, and physical activity free of charge was feasible in terms of patient recruitment and attrition, and the web-based component was well accepted and used by the families. The results on changes in BMI-SDS were promising and 42 of the 51 children lowered their BMI-SDS, and the average reduction was 0.4 SD during the six-month intervention. Even a small reduction in mean BMI-SDS is beneficial to the child's health. To confirm the findings of this feasibility study, a randomized controlled trial with a longer follow up is now underway.

ACKNOWLEDGEMENTS

The development of the web-based program was in part financed by Vinnova 2018-01405.

ORCID

Annelie Thorén  <https://orcid.org/0000-0003-4193-3141>

Sven-Arne Silfverdal  <https://orcid.org/0000-0002-3606-3797>

REFERENCES

1. Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet (London, England)*. 2014;384:766–781.
2. Simmonds M, Llewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev Off J Int Assoc Study Obes*. 2016;17:95–107.

3. Simmonds M, Burch J, Llewellyn A, et al. The use of measures of obesity in childhood for predicting obesity and the development of obesity-related diseases in adulthood: a systematic review and meta-analysis. *Health Technol Assess (Winch Eng)*. 2015;19:1-336.
4. Freedman DS, Goodman AB, King RJ, Blanck HM. Tracking of obesity among 2- to 9-year-olds in an electronic health record database from 2006 to 2018. *Obes Sci Pract*. 2020;6:300-306.
5. Duchon K, Jones M, Faresjö ÅO, Faresjö T, Ludvigsson J. Predicting the development of overweight and obesity in children between 2.5 and 8 years of age: the prospective ABIS study. *Obes Sci Pract*. 2020;1-8.
6. Hakkanen P, But A, Ketola E, Laatikainen T. Distinct age-related patterns of overweight development to guide school healthcare interventions. *Acta Paediatrica (Oslo, Norway: 1992)*. 2020;109:807-816.
7. Stockholm R. Årsrapport Barnhälsovård i Stockholms län 2018 [Web page]. 2018 [cited 2017]. Available from: <https://www.sll.se/globalassets/1.-halsa-och-varld/bilagor---nyhet/bilagor-nyheter-2019/arsrapport-barnhalsovard-2018-1.pdf>
8. Thorn J, Waller M, Johansson M, Mårild S. Overweight among Four-Year-Old Children in Relation to Early Growth Characteristics and Socioeconomic Factors. *J Obes*. 2010;2010:1-6.
9. Mangrio E, Lindström M, Rosvall M. Early life factors and being overweight at 4 years of age among children in Malmö, Sweden. *BMC Public Health*. 2010;10:764.<https://doi.org/10.1186/1471-2458-10-764>.
10. Lissner L, Sohlstrom A, Sundblom E, Sjöberg A. Trends in overweight and obesity in Swedish schoolchildren 1999-2005: has the epidemic reached a plateau? *sObes Rev Off J Int Assoc Study Obes*. 2010;11:553-559.
11. Neovius M, Janson A, Rössner S. Prevalence of Obesity in Sweden. *Obes Rev*. 2006;7:1-3.
12. Hemmingsson E, Ekblom O, Kallings LV, et al. Prevalence and time trends of overweight, obesity and severe obesity in 447,925 Swedish adults, 1995-2017. *Scand J Public Health*. 2020;1403494820914802. <https://doi.org/10.1177/1403494820914802>.
13. Bergström E, Blomquist HKs. Is the prevalence of overweight and obesity declining among 4-year-old Swedish children? *Acta Paediatr*. 2009;98:1956-1958.
14. Eills LJ, Rees K, Brown T, et al. Interventions for treating children and adolescents with overweight and obesity: an overview of Cochrane reviews. *Int J Obes (2005)*. 2018;42:1823-1833.
15. Hagman E, Danielsson P, Lindberg L, Marcus C. Paediatric obesity treatment during 14 years in Sweden: Lessons from the Swedish Childhood Obesity Treatment Register-BORIS. *Pediatr Obes*. 2020;e12626.<https://doi.org/10.1111/ijpo.12626>.
16. Waling M, Bäcklund C, Lind T, Larsson C. Effects on metabolic health after a 1-year-lifestyle intervention in overweight and obese children: a randomized controlled trial. *J Nutr Metab*. 2012;2012:913965. <https://doi.org/10.1155/2012/913965>.
17. Hasson RE, Hsu Y-WJ, Davis JN, Goran MI, Spruijt-Metz D. The influence of parental education on dietary intake in Latino youth. *J Immigr Minor Health*. 2018;20:250-254.
18. Spruijt-Metz D, Lindquist CH, Birch LL, Fisher JO, Goran MI. Relation between mothers' child-feeding practices and children's adiposity. *Am J Clin Nutr*. 2002;75:581-586.
19. Perry R, Daniels L, Baur L, Magarey A. Impact of a 6-month family-based weight management programme on child food and activity behaviours: short-term and long-term outcomes of the PEACH™ intervention. *Pediatr Obes*. 2018;13:744-751.
20. Ek A, Chamberlain KL, Ejderhamn J, et al. The More and Less Study: a randomized controlled trial testing different approaches to treat obesity in preschoolers. *BMC Public Health*. 2015;15:735.<https://doi.org/10.1186/s12889-015-1912-1>.
21. Ek A, Lewis Chamberlain K, Sorjonen K, et al. A parent treatment program for preschoolers with obesity: a randomized controlled trial. *Pediatrics*. 2019;144:e20183457.<https://doi.org/10.1542/peds.2018-3457>.
22. Loveman E, Al-Khudairy L, Johnson RE, et al. Parent-only interventions for childhood overweight or obesity in children aged 5 to 11 years. *Cochrane Database Syst Rev*. 2015;(12):<https://doi.org/10.1002/14651858.CD012008>.
23. Hedman E, Ljótsson B, Lindefors N. Cognitive behavior therapy via the Internet: a systematic review of applications, clinical efficacy and cost-effectiveness. *Expert Rev Pharmacoecon Outcomes Res*. 2012;12:745-764.
24. Delamater AM, Pulgaron ER, Rarback S, et al. Web-based family intervention for overweight children: a pilot study. *Child Obes*. 2013;9:57-63.
25. Barlow SE, Dietz WH. Obesity evaluation and treatment: expert committee recommendations. *Pediatrics*. 1998;102(3):e29.<https://doi.org/10.1542/peds.102.3.e29>.
26. Antwi F, Fazylova N, Garcon MC, Lopez L, Rubiano R, Slyer JT. The effectiveness of web-based programs on the reduction of childhood obesity in school-aged children: A systematic review. *JBI Libr Syst Rev*. 2012;10:1-14.
27. Waling M, Larsson C. Improved dietary intake among overweight and obese children followed from 8 to 12 years of age in a randomised controlled trial. *J Nutr Sci*. 2012;1:e16.<https://doi.org/10.1017/jns.2012.17>.
28. Internetstiftelsen. *The Swedes and the Internet [Web page]*. Internetstiftelsen; 2019 Available from: <https://internetstiftelsen.se/kunskap/rapporter-och-guider/svenskarna-och-internet-2018>
29. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240-1243.
30. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes*. 2012;7:284-294.
31. Folkhälsoinstitut S. FaR - individanpassad skriftlig ordination av fysisk aktivitet. Report. 2011.
32. Sweden Phao. EUPAP - A European model for Physical Activity on Prescription 2020 [Available from: <https://www.folkhalsomyndigheten.se/the-public-health-agency-of-sweden/living-conditions-and-lifestyle/physical-activity/eupap-a-european-model-for-physical-activity-on-prescription/>
33. Karlberg J, Luo Z, Albertsson-Wikland K. Body mass index reference values (mean and SD) for Swedish children. *Acta Paediatr*. 2001;90:1427-1434.
34. Arain M, Campbell MJ, Cooper CL, Lancaster GA. What is a pilot or feasibility study? A review of current practice and editorial policy. *BMC Med Res Methodol*. 2010;10:67.<https://doi.org/10.1186/1471-2288-10-67>.
35. Donkor HM, Grundt JH, Juliusson PB, et al. A family-oriented intervention programme to curtail obesity from five years of age had no effect over no intervention. *Acta Paediatrica*. 2020;109(6):1243-1251. <https://doi.org/10.1111/apa.15080>.
36. Forsell C, Gronowitz E, Larsson Y, Kjellberg BM, Friberg P, Marild S. Four-year outcome of randomly assigned lifestyle treatments in primary care of children with obesity. *Acta Paediatrica (Oslo, Norway: 1992)*. 2019;108:718-724.
37. Dalby S, Vahlkvist S, Østergaard I, Park Jørgensen J, Bogh JC. Two-year treatment programme showed that younger age and initial weight loss predicted better results in overweight and obese children aged 2-16 years. *Acta Paediatr*. 2020;00:1-7. <https://doi.org/10.1111/apa.15145>.
38. Waling M. *Dietary and Metabolic Effects of a 2-Year Lifestyle Intervention in Overweight and Obese Children*. Umeå University; 2011.
39. Moraeus L, Lissner L, Yngve A, Poortvliet E, Al-Ansari U, Sjöberg A. Multi-level influences on childhood obesity in Sweden: societal

- factors, parental determinants and child's lifestyle. *Int J Obes* (2005). 2012;36:969-976.
40. Larsson C, Hernell O, Lind T. Moderately elevated body mass index is associated with metabolic variables and cardiovascular risk factors in Swedish children. *Acta Paediatrica (Oslo, Norway: 1992)*. 2011;100:102-108.
41. Ball GDC, Kebbe M. Preventing and managing paediatric obesity: a special edition on randomized controlled trials. *Pediatr Obes*. 2018;13:635-638.

How to cite this article: Thorén A, Janson A, Englund E, Silfverdal S-A. Development, implementation and early results of a 12-week web-based intervention targeting 51 children age 5–13 years and their families. *Obes Sci Pract*. 2020;6:516–523. <https://doi.org/10.1002/osp4.440>