



LPG stove and fuel intervention among pregnant women reduce fine particle air pollution exposures in three countries: Pilot results from the HAPIN trial[☆]

Jiawen Liao^{a,b,1}, Miles A. Kirby^{a,c,1}, Ajay Pillarisetti^{a,d}, Ricardo Piedrahita^e, Kalpana Balakrishnan^f, Sankar Sambandam^f, Krishnendu Mukhopadhyay^f, Wenlu Ye^a, Ghislaine Rosa^g, Fiona Majorin^g, Ephrem Dusabimana^h, Florian Ndagijimana^h, John P. McCracken^{i,j}, Erick Mollinedo^{i,j}, Oscar de Leon^{a,i}, Anaité Díaz-Artigaⁱ, Lisa M. Thompson^{a,k}, Katherine A. Kearns^j, Luke Naeher^j, Joshua Rosenthal^{l,2}, Maggie L. Clark^m, Kyle Steenland^a, Lance A. Waller^a, William Checkley^{n,o}, Jennifer L. Peel^m, Thomas Clasen^a, Michael Johnson^{e,*}, on behalf of the HAPIN Investigators³

^a Gangarosa Department of Environmental Health, Emory University Rollins School of Public Health, Atlanta, GA, USA

^b Department of Population and Public Health Sciences, Keck School of Medicine of the University of Southern California, Los Angeles, CA, USA

^c Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Boston, MA, USA

^d School of Public Health, University of California, Berkeley, CA, USA

^e Berkeley Air Monitoring Group, Berkeley, CA, USA

^f SRU-ICMR Center for Advanced Research on Air Quality, Climate and Health, Sri Ramachandra Institute of Higher Education and Research, Chennai, India

^g London School of Hygiene and Tropical Medicine, London, UK

^h Eagle Research Center, Kigali, Rwanda

ⁱ Center for Health Studies, Universidad del Valle De Guatemala, Guatemala City, Guatemala

^j College of Public Health, University of Georgia, Athens, GA, USA

^k Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, GA, USA

^l Fogarty International Center, National Institutes of Health, Bethesda, MD, USA

^m Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO, USA

ⁿ Division of Pulmonary and Critical Care, School of Medicine, Johns Hopkins University, Baltimore, MD, USA

^o Center for Non-Communicable Disease Research and Training, School of Medicine, Johns Hopkins University, Baltimore, MD, USA

ARTICLE INFO

Keywords:

Household air pollution

Cookstove

Clean cooking fuel

Intervention

PM_{2.5}

Personal exposure

ABSTRACT

The Household Air Pollution Intervention Network trial is a multi-country study on the effects of a liquefied petroleum gas (LPG) stove and fuel distribution intervention on women's and children's health. There is limited data on exposure reductions achieved by switching from solid to clean cooking fuels in rural settings across multiple countries. As formative research in 2017, we recruited pregnant women and characterized the impact of the intervention on personal exposures and kitchen levels of fine particulate matter (PM_{2.5}) in Guatemala, India, and Rwanda. Forty pregnant women were enrolled in each site. We measured cooking area concentrations of and personal exposures to PM_{2.5} for 24 or 48 h using gravimetric-based PM_{2.5} samplers at baseline and two follow-ups over two months after delivery of an LPG cookstove and free fuel supply. Mixed models were used to estimate PM_{2.5} reductions. Median kitchen PM_{2.5} concentrations were 296 µg/m³ at baseline (interquartile range, IQR: 158–507), 24 µg/m³ at first follow-up (IQR: 18–37), and 23 µg/m³ at second follow-up (IQR: 14–37). Median personal exposures to PM_{2.5} were 134 µg/m³ at baseline (IQR: 71–224), 35 µg/m³ at first follow-up (IQR: 23–51),

[☆] This paper has been recommended for acceptance by Payam Dadvand.

* Corresponding author.

E-mail address: mjohnson@berkeleyair.com (M. Johnson).

¹ These authors contributed equally to this work.

² The views expressed in this article are those of the authors and do not reflect official positions or views of the National Institutes of Health or the U.S. Government.

³ The HAPIN investigators is reference to the HAPIN investigators presented in the acknowledgements section.

<https://doi.org/10.1016/j.envpol.2021.118198>

Received 15 June 2020; Received in revised form 14 September 2021; Accepted 16 September 2021

Available online 21 September 2021

0269-7491/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

and $32 \mu\text{g}/\text{m}^3$ at second follow-up (IQR: 23–47). Overall, the LPG intervention was associated with a 92% (95% confidence interval (CI): 90–94%) reduction in kitchen $\text{PM}_{2.5}$ concentrations and a 74% (95% CI: 70–79%) reduction in personal $\text{PM}_{2.5}$ exposures. Results were similar for each site.

Conclusions: The intervention was associated with substantial reductions in kitchen and personal $\text{PM}_{2.5}$ overall and in all sites. Results suggest LPG interventions in these rural settings may lower exposures to the WHO annual interim target-1 of $35 \mu\text{g}/\text{m}^3$. The range of exposure contrasts falls on steep sections of estimated exposure-response curves for birthweight, blood pressure, and acute lower respiratory infections, implying potentially important health benefits when transitioning from solid fuels to LPG.

1. Introduction

Household air pollution due to use of solid fuels such as wood and charcoal for cooking, heating and lighting is a major risk factor for morbidity and premature mortality (GBD, 2017 Risk Factor Collaborators, 2018). Over 3 billion households rely on solid fuels for cooking, primarily in low-income settings (Bonjour et al., 2013; Health Effects Institute, 2019; WHO, 2016). To date, solid fuel cooking interventions have focused on improved combustion efficiency or ventilation and have largely failed to demonstrate substantial exposure reductions or health benefits (Baumgartner et al., 2019; Kirby et al., 2019; Mortimer et al., 2020, 2016; Pope et al., 2017; Quansah et al., 2017).

Thus, clean cooking fuels, such as electricity, ethanol, biogas, and liquefied petroleum gas (LPG), are widely considered necessary for obtaining exposure reductions and accompanying health and climate benefits (Grieshop et al., 2011). Scale-up of these clean technologies would likely contribute to reaching several Sustainable Development Goals (Rosenthal et al., 2018), including Good health and well-being [SDG 3], Gender equality [SDG 5], Affordable and clean energy [SDG 7], and Climate action [SDG13]. LPG, in particular, is amenable to large-scale dissemination and uptake, and clean cooking programs focusing on LPG are being scaled up worldwide (Gould and Urpelainen, 2018; Quinn et al., 2018).

LPG stoves in lab settings have been shown to reach or surpass the WHO emission rate target guidance for $\text{PM}_{2.5}$ emissions ($0.23 \text{ mg}/\text{min}$) (Johnson et al., 2014; Shen et al., 2018; WHO, 2014) and perform similarly in real-world field conditions (Johnson et al., 2019). However, relatively few studies have assessed exposure reductions that can be obtained by transitioning to cleaner fuels like LPG, particularly in rural settings, where access to these fuels is limited (Pillarsetti et al., 2019b; Pope et al., 2017). The potential for clean cooking fuels to improve health likely depends primarily on high adherence (Johnson and Chiang, 2015; Snider et al., 2018) and on the ability to reduce exposure to levels below or near the WHO annual $\text{PM}_{2.5}$ Interim Target-1 (IT-1) of $35 \mu\text{g}/\text{m}^3$ (Bruce et al., 2015; Burnett et al., 2014; WHO, 2014). While it is believed that switching fuel types can result in these reductions in personal exposure and yield health benefits, there is very little evidence that it can be achieved in real-world settings (Steenland et al., 2018). For example, in a recent stepped wedge trial in rural Nepal, households with LPG had a mean 20-h kitchen $\text{PM}_{2.5}$ concentration of $442 \mu\text{g}/\text{m}^3$ and no differences in birth outcomes compared to houses with solid fuel stoves, likely due to continued use of solid fuels, air pollution from neighboring houses, and ambient pollution influenced in part by dust (Katz et al., 2020). Another recent trial in rural Ghana found that mothers randomized to receive LPG and cylinder refills had 32% lower mean 48-h $\text{PM}_{2.5}$ personal exposure concentrations compared to women in the control arm (solid fuel) (Chillrud et al., 2021). However, the mean in the LPG arm was $52 \pm 29 \mu\text{g}/\text{m}^3$, and two thirds of post-intervention maternal measurements in the LPG arm were above $35 \mu\text{g}/\text{m}^3$, possibly indicating the influence of ambient air quality and persistent solid fuel use (Chillrud et al., 2021).

The Household Air Pollution Intervention Network (HAPIN) Trial is a randomized controlled trial seeking to evaluate the health impacts of an LPG stove and fuel supply intervention on birthweight, stunting and severe pneumonia among children ≤ 1 year of age, and blood pressure

among older women. The study is taking place in Guatemala, India, Peru, and Rwanda, and is currently ongoing (<https://clinicaltrials.gov/ct2/show/NCT02944682> (Clasen et al., 2020)). As an efficacy trial, the aim is to maximize intervention uptake and use by providing the stove and fuel free of charge and by conducting comprehensive promotion and behavior change measures.

As part of formative research for the trial, numerous ambient, kitchen and personal $\text{PM}_{2.5}$ concentration estimates have been conducted to assist in trial site selection and intervention design, as reported previously (Clasen et al., 2020; Sambandam et al., 2020). Other theory-grounded formative work has been conducted on participants' perceptions of LPG, cooking behavioral differences across study sites, and the implementation components of the HAPIN trial, as well as development of strategies to monitor, promote and reinforce exclusive use of LPG for all cooking needs (Williams et al., 2020). Here we present results from a HAPIN pilot study in India, Rwanda, and Guatemala to estimate exposures to $\text{PM}_{2.5}$ among pregnant women before and after a short-term LPG intervention, with the aim of characterizing baseline exposures and contrasts that might be expected as a result of the intervention in the main trial.

2. Methods

2.1. Study site and population

The HAPIN pilot intervention study was conducted in the countries where the main trial was proposed. Potential HAPIN sites were short-listed by country lead investigators in communities in India, Guatemala, Peru, and Rwanda. In Peru, the pilot intervention was conducted within the context of the Cardiopulmonary outcomes and Household Air Pollution trial (<https://clinicaltrials.gov/ct2/show/NCT02994680>) (Fandiño-Del-Río et al., 2017), and the results have been published separately (Checkley et al., 2020).

Study sites were selected in rural areas, based primarily on the estimated proportion of households using biomass fuel; we additionally selected areas with relatively low levels of ambient air pollution, as determined during formative scoping activities preceding this pilot work. In India, we recruited from Villupuram and Nagapattinam Districts in Tamil Nadu; in Guatemala, from San Pedro Pinula and Jalapa municipalities from the Jalapa Department; and in Rwanda, from the Kabare, Murama and Kabarondo sectors in Kayonza District, Eastern Province.

2.2. Inclusion criteria

We enrolled 40 pregnant women in each study site. Pregnant women seeking routine antenatal care were screened at health centers and hospitals within study sites. The inclusion criteria were similar to those proposed for the main trial: 1) non-smoking pregnant women aged 18–34 years with gestational age 9–20 weeks confirmed through ultrasound (singleton pregnancy, without fetal anomalies or pregnancy complications identified at first ultrasound), 2) households using solid fuels as their main cooking fuel, and 3) no plans to move out of current household. Household selection among eligible houses was based on convenience and proximity to study offices in each study site. Details of

household enrollment and follow-up are included in [supplementary information \(SI\)](#).

2.3. Study design and data collection

As part of HAPIN pilot phase, this study was designed as an LPG stove and fuel “before-and-after” single-arm, non-randomized intervention study (Sedgwick, 2014). For each household, we conducted one pre-intervention baseline visit and two follow-up visits (at approximately one- and two-months post-intervention), between July and November 2017. During each visit, we measured PM_{2.5} levels for 24 or 48 h (described below) and conducted household surveys using REDCap (Vanderbilt University, Nashville, TN, USA) to assess social demographic information, cooking practices and adherence of LPG intervention, compliance wearing the monitoring equipment, and to identify other potential sources of air pollution.

2.4. Intervention

Approximately 1 week after enrollment and baseline measurements, households received a new 2 or 3-burner gas stove, a regulator and hose, and a cylinder of gas that was refilled for free, upon demand, for two months. Cylinder fuel quantities were 15 kg in Rwanda, 14.2 kg in India, 11.3 kg in Guatemala. Upon receipt of the intervention, households completed a pledge to use it exclusively (Williams et al., 2020). Due to differences in perceptions of LPG use in Guatemala and Rwanda as compared to study sites in India, the intervention in Guatemala and Rwanda also included a behavior change component with messages tailored to address motivators and barriers to LPG use that was delivered in the local language, a demonstration of how to use the LPG stove safely, supervised practice with the stove, locally appropriate cooking tips, standardized videos showing how to use it, and printed educational and promotional materials that households could keep. Households were visited every 2–4 weeks (1) to answer any questions related to LPG stove use, (2) to reinforce behavior change messages, and (3) to provide cylinder refills as needed.

2.5. Air pollutant measurements

In each household, we conducted 24- or 48-h sampling of PM_{2.5}

kitchen concentrations and personal exposures, once at baseline and twice at approximately around one- and two-months post-intervention, using active air samplers to estimate PM_{2.5} mass concentrations. Based on feasibility, 24-h sampling was conducted in Guatemala and Rwanda, while 48-h sampling was conducted in India. For kitchen area samples, instruments were placed in the same location each time for 24 or 48 h unless the kitchen location changed post-intervention: 1.5 m above ground, 1 m away from any doors and windows, and 1 m away from the combustion zone of the primary cookstove. For personal exposure samples, an apron or vest was designed at each site to carry the sampling devices, and participants were advised to wear this during the whole measurement period of 24 or 48 h. When bathing or sleeping, participants were instructed to place the apron/vest with the instruments less than 1 m away from them.

Due to logistical constraints, we used multiple instruments and size selective inlets to sample PM_{2.5} on either 15 mm (PT15-AN-PF02, MTL LLC., Minneapolis, MN USA) or 37 mm PTFE filters (Pall Life Sciences, Port Washington, NY, USA). We measured pump flow rates before and after sampling using a flow meter (Table 1). Filters were pre-weighed at temperature and humidity-controlled labs at Emory University (Atlanta GA, USA), University of Georgia (Athens GA, USA), Sri Ramachandra Institute of Higher Education and Research (Chennai, India), and Harvard University (Cambridge MA, USA). After sampling, filters were stored in a freezer and transported back under cold conditions to the same weighing lab where they are pre-weighed. In the total of 999 PM_{2.5} samples, we excluded PM_{2.5} samples whose average flow rate deviated from the desired flow rate by more than 10% (n = 7, 1%), whose sample duration deviated by more than 10% of the required duration (n = 38, 4%), and all filters with damage, such as holes or tears (n = 39, 4%), resulting in a total of 915 eligible gravimetric PM_{2.5} samples (kitchen area n = 413; personal exposure n = 502). Most of the eligible PM_{2.5} samples (n = 463, 51%) were collected on 15 mm filters using the Enhanced Children’s MicroPEM (ECM, RTI International, Durham NC, USA), a robust, light gravimetric PM_{2.5} sampling instrument for the HAPIN trial with similar size to a cell phone. Additionally, 373 samples were collected on 37 mm filters using gravimetric pumps (SKC XR5000, Casella Air Tuff, and Casella TuffPro) and 79 samples on 37 mm filters using Access Sensor Technologies’ (AST) Ultrasonic Personal Aerosol Sampler (UPAS) (Volckens et al., 2017). Detailed information regarding PM_{2.5} gravimetric instruments, filter weighing, blank filters, and sample

Table 1

Details of gravimetric PM_{2.5} sampling instrument, number of samples, and filter weighing facilities in India, Guatemala, and Rwanda.

	India			Guatemala		Rwanda	
Instrument	SKC XR5000 pump, SKC PCXR8, Casella	AST UPAS ^a	RTI ECM	Casella Tuff pump	RTI ECM	Casella Tuff 3 Pro IS pump	RTI ECM
Size Selector	BGI Triplex cyclone	Built-in cyclone	Built-in impactor	BGI Triplex cyclone	Built-in impactor	H-PEM impactor (3243–3229)	Built-in impactor
Nominal flow rate (LPM)	1.5	1	0.3	1.5	0.3	1.8	0.3
Flow meter	Gilian Gilibrator-2 (Sensidyne) and TSI 41401			Gilian Gilibrator-2 (Sensidyne)		Gilian Gilibrator-2 (Sensidyne)	
Sampling Time (hour)	48	48	48	24	24	24	24
Filter diameter (mm)	37	37	15	37	15	37	15
Number of total eligible samples	8 (kitchen:8)	79 (kitchen:37; personal: 42)	133 (kitchen:57; personal: 76)	216 (kitchen:114; personal: 102)	230 (kitchen:121; personal: 109)	149 (kitchen:43; personal: 106)	100 (kitchen:33; personal: 67)
Number of unique samples ^b	0	68 (kitchen:31; personal: 37)	133 (kitchen:57; personal: 76)	40 (kitchen:15; personal: 25)	193 (kitchen:104; personal: 89)	57 (kitchen:11; personal: 46)	100 (kitchen:33; personal: 67)
Sample duration (minutes)	2880			1440		1440	
Weighing facility	Sri Ramachandra Institute of Higher Education and Research (SRIHER)			Emory University, University of Georgia, Harvard University		Emory University, University of Georgia	

^a UPAS: Ultrasonic Personal Air Sampler; ECM: Enhanced Children’s MicroPEM; LPM: liters per minute.

^b The difference between number unique samples and eligible samples is due to removal of co-located samples, with *a priori* using ECM instrument, gravimetric pumps (SKC and Casella) and then UPAS instruments.

quality assurance is in the [supplementary information \(SI\)](#).

Among all samples, 299 (72%) of 413 kitchen area samples were co-located samples (using two side-by-side sampling devices with the same operating time in the same location); 306 (61%) of 502 personal samples were co-located and collected with side-by-side PM_{2.5} samplers during the same household visit. ECMs were deemed as our *a priori* sampler and we only use ECM measurements from co-located samples, therefore, most of samples (N = 426) are from ECMs compared to other instruments (N = 165, [Table 1](#)). For any given household visit without ECM samples, we used samples collected by gravimetric pumps or UPAS instruments. Overall, we had 591 unique samples for following statistical analysis, comprised of kitchen area PM_{2.5} concentrations (N = 251) and pregnant women's personal exposure (N = 340). Among these samples, most were from ECM instruments (N = 426, 72%) and around a quarter were from gravimetric pumps (N = 97, 16%) and UPAS (N = 68, 12%).

The PM_{2.5} mass concentration for each sample was calculated using the average measured flow rates, the sample duration, and net change in mass of the filters (equation (1)).

$$PM\ Conc. = \frac{(WT_{post} - WT_{pre}) - WTC_{BN}}{Flow \times Duration} \times 1000 \quad (1)$$

PM Conc. is the gravimetric PM_{2.5} concentration in µg/m³; WT_{post} and WT_{pre} are the post- and pre-weight of the filters in µg. WTC_{BN} is the net filter mass change for blank filters (blanks were collected at each site in sampling pumps when they are turned-off). Flow is the sampler flow rate in L/min. In situations where the pre- or post-flow rate was not measured using a primary flow standard for ECMs, we used the average real-time flow rate logged by that ECM (N = 11, 2%). Duration is the sampling duration in minutes as recorded by field workers or logged by sampling devices.

We also collected 71 field blank filters spread out over the study period (India: 5 for 15 mm filters, 11 for 37 mm filters; Guatemala: 10 for 15 mm filters, 14 for 37 mm filters; Rwanda: 14 for 15 mm filters and 17 for 37 mm filters) to account for changes in filter mass due to handling and transportation. The median value of field blanks in Guatemala were 1.0 µg for 15 mm filters and -0.6 µg for 37 mm filters; in India, 0.3 µg for 15 mm filters and 0.75 µg for 37 mm filters; and in Rwanda, -0.25 µg for 15 mm filters and 10 µg for 37 mm filters.

2.6. Statistical analysis

Median, first, and third quartiles of kitchen gravimetric PM_{2.5} concentrations and personal PM_{2.5} exposures by each visit were summarized for 591 unique samples. We then used mixed effects linear models to estimate the reduction of kitchen area and personal PM_{2.5} after introduction of the LPG intervention. As kitchen PM_{2.5} measurements are log-normally distributed, we first log-transformed PM_{2.5} measurements, and checked normality of the distribution based on a quantile-quantile plot ([Figure S1, SI](#)). Then, we applied a linear mixed-effect model with household random intercepts to assess the reduction of kitchen PM_{2.5} concentrations and personal PM_{2.5} after LPG intervention. We included LPG intervention status (pre vs post), household characteristics, reported and observed sources of other air pollution during each household visit (self-reported smoke from cigarette smoking, trash burning, kerosene lighting, generators, smoking meat, and burning crops), and rainfall and temperature in the model (equation-2). Daily rainfall (in cm) and average temperature on the first day of each visit were determined by Climate Hazards Group InfraRed Precipitation with Station Data (CHIRPS) 2.0 ([Funk et al., 2015](#)) and Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA -2, <https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/>) for the geographical centroid of each study site, respectively.

$$\log(PM_{ij}) = \beta_0 + \beta_1 LPG_{ij} + \beta_2 X_{ij} + \beta_3 W_i + u_i + e_{ij} \quad (2)$$

PM_{ij} is kitchen area PM_{2.5} or personal PM_{2.5} exposure for household *i* at time *j*; β₀ is the overall intercept, LPG_{ij} indicates whether there is a LPG intervention for the measurement in household *i* at time *j*; X_{ij} includes time-dependent explanatory variables, including temperature, rainfall, and self-reported other sources of pollution during the measurement period; W_i are time-independent explanatory variables, including study site, pregnant woman's age and years of education, her husband's years of education, and the number of people in the household. u_i is the household random intercept of normal distribution with mean 0 and variance σ_b², and e_{ij} is the model residual following a normal distribution with mean 0 and variance σ_e². The random intercepts adjust for repeated measures from the same household where σ_b² refers to the between household variance and σ_e² refers to within household variance.

Subgroup stratified analysis were conducted for each site, and reductions of kitchen area and personal PM_{2.5} exposure were presented separately for each site. Additionally, we conducted sensitivity analysis for restriction to only households where kitchens locations were not changed after the intervention was delivered (SI). Data manipulation and processing were performed in R (version 3.6.0) and statistical analyses were performed in Stata (version 14.2). All relevant data and codebooks are publicly available ([Liao et al., 2021](#)).

2.7. Reduction of associated health effects

We modeled the potential health benefits for three of the primary outcomes assessed as part of the main HAPIN trial ([Clasen et al., 2020](#)): systolic blood pressure (SBP), birth weight, and as a proxy for severe pneumonia, acute lower respiratory infection (ALRI) by evaluating changes in outcomes or risk associated with reductions in the median personal PM_{2.5} exposure before and after LPG intervention based on state-of-the-science exposure-response curves ([GBD, 2017 Risk Factor Collaborators, 2018](#); [Steenland et al., 2018](#)). Uncertainty interval (UI) bounds around these projected health effects were based on the interquartile range (IQR) of exposure estimates and 95% confidence interval of exposure-response curves.

2.8. Ethics

This pilot study received ethical approval from Emory IRB (00089,799), Universidad del Valle de Guatemala (146-08-2016/11-2016) and the Guatemalan Ministry of Health National Ethics Committee (11-2016), the Indian Council of Medical Research-Health Ministry Screening Committee (5/8/4-30/(Env)/Indo-US/2016-NCD-I), Sri Ramachandra Institute of Higher Education and Research (IEC-N1/16/JUL/54/49), London School of Hygiene and Tropical Medicine (11,664-2), and the Rwandan National Ethics Committee (No.148/RNEC/2017). Prior to participation, details of the study were provided to prospective participants and informed written consent was obtained.

3. Results

One hundred and twenty pregnant women from 120 households, were enrolled across all three countries and were followed from July through November 2017. Enrolled households were visited at baseline and for two follow-up visits after LPG fuel and stove intervention, with the exception of 1 household in Rwanda which exited the study prior to the first post-LPG exposure measurement visit.

3.1. Participant and household characteristics

Across the three countries, the mean age of pregnant women was 26.2 years (standard deviation 4.3) ([Table 2](#)). All participants were the primary cooks in their households. In Rwandan and Indian households, most women worked as farmers, while in Guatemala, housework was listed as the main occupation. At baseline, most of the households had

Table 2
Characteristics of enrolled pregnant women and households at baseline.

Variables	India	Guatemala	Rwanda	Overall
Characteristics of women participants				
	N = 40	N = 40	N = 40	N = 120
Age in years, mean (SD)	24.8 (3.3)	26.2 (4.2)	27.9 (4.3)	26.2 (4.3)
Years of schooling, pregnant woman, mean (SD)	7.5 (5.1)	3.65 (3.1)	5.6 (3.3)	5.6 (4.2)
Years of schooling, male partner, mean (SD)	7.4 (4.1)	4.15 (3.0)	6.6 (4.1)	6.1 (4.0)
Number of children under 12 years of age, mean (SD)	0.73 (0.7)	1.85 (1.4)	1.10 (0.95)	1.21 (1.1)
Number of adults in household, mean (SD)	2.93 (1.25)	2.70 (1.7)	2.58 (1.2)	2.74 (1.4)
Employment status of pregnant woman				
Employed, n (%)	4 (10%)	38 (95%)	2 (5%)	44 (37%)
Unemployed or farmer, n (%)	36 (90%)	2 (5%)	38 (95%)	76 (63%)
Exposure to other sources of air pollution ^a , n (%)	11 (27.5%)	25 (62.5%)	15 (37.5%)	51 (42.5%)
Household Characteristics				
Kitchen type, n (%)				
Fully enclosed (roof with 4 walls)	32 (80%)	32 (82%)	22 (55%)	86 (71.7%)
Not enclosed/outdoor cooking	8 (20%)	8 (18%)	18 (20%)	34 (28.3%)
Fully enclosed kitchen measurements				
Kitchen size in m ² , mean (SD)	9.9 (4.1)	12.5 (4.7)	5.2 (1.6)	10.0 (4.8)
Kitchen height in m, mean (SD)	2.8 (0.6)	2.53 (0.3)	2.58 (0.3)	2.64 (0.5)
# of doors, mean (SD)	1.40 (0.5)	1.15 (0.4)	1.14 (0.35)	1.24 (0.4)
# of windows or other openings, mean (SD)	1.82 (1.6)	7.06 (6.9)	1.23 (1.3)	3.63 (5.1)
Location of kitchen changed, n (%)	24 (60%)	10 (25%)	36 (90%)	70 (58%)
# of stoves, mean (SD)	1.31 (0.5)	1.47 (0.5)	1.33 (0.6)	1.37 (0.5)
Primary stove type, n (%)				
Open fire stove (three-stone)	40 (100%)	35 (87.5%)	11 (27.5%)	86 (71.7%)
Biomass stove with chimney Rondereza (in-built wood-burning stove without chimney in Rwanda)	0	5 (12.5%)	0	5 (4.2%)
Portable stove (wood or charcoal)	0	0	10 (25%)	10 (8.3%)
Other	0	0	1 (7.5%)	1 (2.4%)
Primary stove fuel type, n (%)				
Grass/Shrubs	0	0	9 (22.5%)	9 (7.5%)
Agricultural waste	0	0	1 (2.5%)	1 (0.8%)
Wood	40 (100%)	40 (100%)	25 (62.5%)	105 (87.5%)
Charcoal	0	0	5 (12.5%)	5 (4.2%)
# of burners of primary stove, n (%)				
1	4 (9.7%)	28 (70%)	29 (72.5%)	61 (50.1%)
2	36 (90.3%)	4 (10%)	9 (22.5%)	49 (40.8%)
≥3	0	8 (20%)	2 (5%)	10 (4.2%)
Meteorological Characteristics				
Average Rainfall (cm)	6.1	5.7	1.5	4.5
Average temperature (°C)	29.3	22.9	22.4	25.0

^a Self-reported exposure to second-hand smoke, mosquito coil burning, trash burning, kerosene burning, incense burning, electricity generator, smoking meat, or crop burning over the past 24 h.

fully enclosed kitchens; 28.3% (n = 34) of the households had unenclosed kitchens or cooked outdoors. Most households cooked with an open fire or built-in biomass stove without chimney (n = 104, 86.7%), or portable stoves with wood or charcoal (n = 10, 8.3%) (Table 2). All households used biomass (wood, grass/shrubs, charcoal or agriculture residue) as their primary fuel at baseline. After the LPG intervention, all open, outdoor kitchens were moved indoors (n = 70), typically within the house or to a separated kitchen. This change of kitchen location was mainly due to household preference and the feasibility of indoor LPG stove installation.

3.2. LPG use

Households had high compliance with the LPG intervention, according to field worker observation (at end of measurement period) and self-reported fuel use during the 24/48-h measurement period. Only 3 out of 226 household visits indicated biomass use after receipt of the intervention, and all were from the second follow-up approximately two months after receipt (2 households from Guatemala and 1 from Rwanda).

3.3. PM_{2.5} concentrations

We show the distribution of log-transformed 24/48-h kitchen area and personal PM_{2.5} concentrations by study site and LPG intervention status in Fig. 1. Only Guatemala and India conducted baseline kitchen PM_{2.5} measurements (in a total of 65 households). Ninety-four and 92 kitchen area measurements were collected from all three sites, from follow-up 1 and follow-up 2 visits, respectively.

Baseline personal PM_{2.5} exposure measurements were conducted in three sites. We found that the LPG fuel and stove intervention resulted in a substantial reduction of both kitchen level and personal PM_{2.5} concentrations, in 591 samples removing duplicates (Table 3). The median (Q1-Q3) kitchen level PM_{2.5} concentrations were 296 (158–507) µg/m³ at baseline, 24 (18–37) µg/m³ at follow-up 1 and 23 (14–37) µg/m³ at follow-up 2 after LPG intervention. The median (Q1 - Q3) personal PM_{2.5} exposure were 134 (71–224) µg/m³ at baseline, 35 (23–51) µg/m³ at follow-up 1 and 32 (23–47) µg/m³ at follow-up 2. The complete samples of gravimetric PM_{2.5} measurements (N = 915) with duplicates showed similar reductions of kitchen area PM_{2.5} and personal PM_{2.5} exposure (Table S1 in SI).

3.4. Reduction of personal PM_{2.5} exposure from LPG intervention

In Table 4, we show, using mixed effect linear models, that the LPG intervention is associated with a 92% (95% CI: 90%–94%) reduction of kitchen PM_{2.5} concentrations and a 74% (95% CI: 70%–79%) reduction of personal PM_{2.5} exposures. Reductions of kitchen and personal PM_{2.5} after LPG intervention are similar across sites. We observed a higher reduction of kitchen PM_{2.5} concentrations compared to personal PM_{2.5} exposures, with the highest reductions of PM_{2.5} in Guatemalan kitchens. In the sensitivity analysis including only households where kitchen locations have not changed (N = 50), the reduction due to the LPG intervention is similar (SI).

As shown in Fig. 2, observed personal exposures after LPG intervention suggest transitioning to LPG in these settings could result in health benefits. Given the exposure reductions we observed, we estimate a reduction of 5.7 mmHg (uncertainty interval (UI): 5.1–6.1) in SBP after LPG, a 250 g (UI 205–315) increase in birth weight; and a reduction in ALRI relative risk from 1.8 (1.6–2.1) to 1.4 (1.3–1.5).

4. Discussion

4.1. Context of HAP exposures

The introduction of LPG stoves and fuel in primarily biomass-using




Fig. 1. Distributions of kitchen PM_{2.5} concentrations and personal PM_{2.5} exposure before (biomass) and after LPG intervention for 591 Samples.

Table 3

Summary statistics (median, first quartile, third quartile) of kitchen and personal PM_{2.5} levels (µg/m³) for 591 unique samples by each visit and site.

		Baseline			Follow-up 1			Follow-up 2		
		N	Median	Q1 – Q3	N	Median	Q1 – Q3	N	Median	Q1 – Q3
India ^a	Kitchen (48 h)	26	191	96–368	30	20	14–31	32	32	23–38
	Personal (48 h)	40	71	50–134	38	26	20–37	35	28	24–38
Guatemala	Kitchen (24hr)	39	433	193–511	40	27	19–37	40	15	7–24
	Personal (24 h)	39	151	86–345	37	38	30–45	38	30	13–60
Rwanda	Kitchen (24hr)				24	26	20–36	20	25	19–37
	Personal (24 h)	35	175	112–276	39	46	26–67	39	36	26–49
All	Kitchen	65	296	158–507	94	24	18–37	92	23	14–37
	Personal	114	134	71–224	114	35	23–51	112	32	23–47

N: number of measurements, Q1: first quartile, Q3: third quartile; grey shaded area indicates no measurements were conducted.

^aEstimates presented here are derived from the same database as those reported by Sambandam et al. (2020). The number of measurement (N) diverges slightly due to different exclusion criteria, as we excluded samples that deviated more than 20% of target length of time or samples with filter damage, differing slightly to exclusion criteria used by Sambandam et al. (2020).

Table 4

Reduction (95% confidence interval) of kitchen PM_{2.5} concentration and personal PM_{2.5} exposure with LPG intervention.^a

Model	All three sites	Stratified by site		
		India	Guatemala	Rwanda
Kitchen PM _{2.5}	92% (90%–94%)	90% (86%–93%)	97% (90%–99%)	n/a
Personal PM _{2.5}	74% (70%–79%)	58% (45%–67%)	83% (76%–88%)	70% (60%–77%)

^a Based on linear mixed effects models, adjusted by study site, pregnant woman’s age and years of education, husband’s year of education, the number of people in the household, temperature, rainfall, and self-reported other sources of air pollution during the measurement period. Kitchen PM_{2.5} concentrations and personal PM_{2.5} exposures are log-transformed; grey shaded area indicates no analysis was conducted.

households, accompanied by behavior change strategies, was associated with substantial reductions in both kitchen concentrations of and personal exposures to PM_{2.5} in three countries. Effects were consistent for two follow-up measurements one month apart, and overall the intervention yielded median kitchen and personal concentrations at, or lower than the annual WHO interim target-1 (IT-1) of 35 µg/m³ for the first and second follow-ups. The effect was most pronounced in Guatemala for both kitchen concentrations and personal exposures, potentially due

to low ambient pollution and the limited impacts of other sources of exposure, as well as high baseline PM_{2.5} levels. Only 3 households (2.5%) reported biomass stove use after receiving the LPG intervention. Although relying on reported stove and fuel use is not as accurate as direct measurement using stove use monitors, the lack of observed traditional stove use when field workers arrived and anecdotal reports from field staff suggests that LPG was likely used nearly exclusively.

Variability in kitchen concentrations and exposures was evident between field sites, with higher PM_{2.5} levels in Rwanda and Guatemala, compared to those measured in India. This type of variation has been reported in systematic HAP review papers (Pope et al., 2017; Quansah et al., 2017), and is similar to the pattern observed in the recent Prospective Urban and Rural Epidemiological (PURE) study, which also reported higher PM_{2.5} exposures for women in biomass-using homes in Africa (geomean = 153 µg/m³) compared to India (geomean = 89 µg/m³) (Shupler et al., 2020). There are several reasons for regional variation in HAP exposures due to differences in environmental and behavioral factors, such as location of cooking (indoors/outdoors), quantity of cooking and/or fuel use, proximity to neighbors, and others. Here, we speculate that all these factors likely contributed to the differences of HAP we observed.

The relative reductions in PM_{2.5} were greatest for kitchens, which is unsurprising given stoves are the dominant source in that environment. Our observed kitchen reductions are similar to a study in Cameroon (LACE-1), which found significant differences between exclusively




Fig. 2. Potential effects of changes in $PM_{2.5}$ personal exposure on systolic blood pressure (SBP) and birth weight (BW) (both adapted from Steenland et al., 2018) and relative risk (RR) of acute lower respiratory infection (ALRI) (adapted (GBD, 2017 Risk Factor Collaborators, 2018). Across all panels, the modeled exposure response curve is shown as a solid black line with uncertainty intervals in light grey. The pink dashed line is the median personal exposure during the baseline period; the green solid line is the median personal exposure during the follow-up period after LPG intervention. The correspondingly colored shaded areas are the IQR of measurements during that period. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

biomass using households and those using LPG, with median kitchen concentrations of $447.9 \mu\text{g}/\text{m}^3$ and $21.1 \mu\text{g}/\text{m}^3$ respectively (Bruce et al., 2018). Personal exposures among women from the same study were $46.9 \mu\text{g}/\text{m}^3$ and $14.0 \mu\text{g}/\text{m}^3$ respectively, although the relatively low exposures observed may have been due to season and behavior, including minimal time spent in kitchen (Bruce et al., 2018). In general, personal exposure levels were slightly higher than kitchen levels during the post-intervention period, and kitchens had more dramatic reductions, perhaps indicating other sources of exposure experienced outside of kitchens. Similarly, a recent study in India found that switching to LPG resulted in an 85% reduction in kitchen $PM_{2.5}$, but mean levels were only reduced to $76 \mu\text{g}/\text{m}^3$ (approximately twice the annual interim target), potentially due to unmeasured traditional stove use and ambient air pollution (Pillariseti et al., 2019b). A larger proportional reduction in kitchen rather than personal exposure was also observed in a wood-burning chimney stove intervention in Guatemala (Smith et al., 2010). We observed a similar trend (higher personal vs kitchen after intervention) most prominently in Rwanda at the first round of follow-up, although the second round saw further reductions in personal exposure which approached kitchen levels.

In settings of low ambient air pollution and near-exclusive stove uptake, both kitchen and personal exposure levels of $PM_{2.5}$ can approach or meet health-based WHO targets, as observed in a recent LPG trial in Peru where 23% of kitchens and 78% of personal measurements were

less than $35 \mu\text{g}/\text{m}^3$ in the intervention arm (Checkley et al., 2020). By contrast, clean fuel use within a context of non-exclusive use, neighboring traditional stove use, and higher ambient levels can limit exposure reduction potential, as observed in the GRAPHS trial where two thirds of post intervention maternal measurements were above $35 \mu\text{g}/\text{m}^3$ in the intervention arm (Chillrud et al., 2021). Similarly, a trial in Nigeria found limited personal exposure contrasts between mothers assigned to ethanol compared to kerosene and wood-burning stoves, with mean 72-h $PM_{2.5}$ means of 118 (SD 166) $\mu\text{g}/\text{m}^3$ vs. 102 (SD 102) $\mu\text{g}/\text{m}^3$ respectively in the dry season, and 61 (SD 74) vs 66 (SD 82) $\mu\text{g}/\text{m}^3$ respectively in the rainy season (Alexander et al., 2018). More dramatically in rural Nepal, households assigned to LPG had a mean 20-h kitchen concentration above $400 \mu\text{g}/\text{m}^3$ (Katz et al., 2020).

Our $PM_{2.5}$ findings further suggest the capacity to reach the WHO targets with a clean fuel intervention, and are generally similar to, or lower than other estimates of $PM_{2.5}$ among LPG-using households in these regions. Recent modeled global and regional estimates for kitchen and personal $PM_{2.5}$ across different fuel types in rural areas indicate that our results are similar or in some cases lower (Shupler et al., 2018). Globally, it is estimated that transitioning from traditional cooking with wood to gas would reduce mean (95% CI) of kitchen concentrations from 395 (148, 1039) $\mu\text{g}/\text{m}^3$ to 105 (39, 273) $\mu\text{g}/\text{m}^3$ (Shupler et al., 2018), and the PURE study, conducted across 8 countries, found that mean kitchen concentrations among gas-using households was $45 \mu\text{g}/\text{m}^3$

(95% CI 43–48), 53 $\mu\text{g}/\text{m}^3$ among electricity-using households, and above 90 $\mu\text{g}/\text{m}^3$ among charcoal, wood, and agricultural/crop waste-burning households, and over 200 $\mu\text{g}/\text{m}^3$ among animal dung and shrub/grass-burning households (Shupler et al., 2020). Encouragingly, a recent stepped wedge intervention trial in Honduras comparing traditional wood-burning cooking to a wood stove with an improved combustion chamber and chimney (*Justa*) had substantially lower kitchen $\text{PM}_{2.5}$ and personal exposure concentrations with means below 80 and 40 respectively, and with over 40% of personal measures falling below 37.5 $\mu\text{g}/\text{m}^3$ (Benka-Coker et al., 2021).

4.2. Implications for HAPIN trial

Exposure reductions fell on steep parts of currently estimated exposure-response curves for several important health outcomes (Bruce et al., 2015; Burnett et al., 2014; Steenland et al., 2018). Therefore, based on the exposure-response curves, we should be able to observe health benefits associated with LPG intervention if the reduction of personal $\text{PM}_{2.5}$ is consistently maintained across a longer observational period. Furthermore, based on these exposure-response curves, even reductions of personal $\text{PM}_{2.5}$ exposures above 35 $\mu\text{g}/\text{m}^3$, the annual IT-1 guideline, would also be expected to deliver some health benefits, particularly for blood pressure, birthweight, and ALRI (Steenland et al., 2018). Another published $\text{PM}_{2.5}$ -mortality hazard ratio function based on outdoor air pollution in multiple countries suggests these reductions may lower risk of mortality as well (Burnett et al., 2018). However, reducing kitchen and personal exposures below WHO interim guidelines through a stove intervention does not guarantee health benefits will be observed, as reported in a recent trial in rural Peru which found no impacts among 180 women aged 25–64 on systolic or diastolic blood pressure, peak expiratory flow, or respiratory symptoms (Checkley et al., 2020). Furthermore, there was no exposure-response relationship found between personal exposure to HAP and primary blood pressure outcomes (Checkley et al., 2020). Reasons for this may include small sample size, a relatively healthy population that under studied, and limited length of time during which exposure reductions were sustained and health effects were studied. This suggests that longer follow-up may be necessary, especially for chronic disease outcomes (Checkley et al., 2020; Kaali et al., 2020). Furthermore, persistence of other pollutants present when cooking with gas, such as NO_2 , may minimize potential health benefits (Kephart et al., 2021). Given the variability and uncertainty in exposure-response relationships, exposures may need to be much lower than WHO interim targets and sustained over longer periods of the time in order to sufficiently achieve meaningful health effects (Kaali et al., 2020).

Thus, while this pilot study's $\text{PM}_{2.5}$ exposure contrasts fell on promising sections of published exposure-response curves (e.g. Fig. 2), the potential for clean cooking interventions to meaningfully improve health remains to be demonstrated and these curves may be updated as more observational studies and trials are conducted. Trials such as HAPIN are thus needed to address important gaps in the evidence for health benefits of cleaner cooking, as well as help better characterize exposure-response curves for several health outcomes that contribute substantially to morbidity and mortality globally (Clasen et al., 2020).

4.3. Limitations

This study has a number of limitations. Perhaps most important, as pilot work, the sample sizes were relatively small, and although we found significant differences, the kitchen concentration and exposure estimates are not highly robust. Further, the convenience sample and before-and-after single-arm, non-randomized intervention design without a parallel control group leaves open the possibility that effects were due in part to selection bias and unmeasured time-varying confounders. Particularly, seasonal differences in environmental conditions and household practices could have contributed to observed differences,

although we controlled for temperature and precipitation in an effort to account for these changes, and there was little variation between baseline and follow-up in each study site. The short length of follow-up time over which air quality monitoring was conducted is another limitation, even though this can minimize impact of season and other temporal trends of ambient and household $\text{PM}_{2.5}$ levels. Use of the intervention stove may decrease over time (Pillarsetti et al., 2014), and traditional stove use may increase, especially for certain cooking tasks where traditional stoves may be preferred (Ruiz-Mercado and Masera, 2015). Certain cooking tasks and stove preferences may vary seasonally, and other seasonal activities such as heating can also impact exposure. A study in rural China found that personal exposure was only 37% lower among those who primarily used LPG or electricity compared to biomass users, although this was only observed in summer when heating was not as common (Ni et al., 2016). We speculate that unmeasured confounders across sites and visit periods contributed to relative differences in kitchen concentrations of and personal exposures to $\text{PM}_{2.5}$ post-intervention; however, we are not able to test these hypotheses due to limited contextual data we were able to feasibly collect, and small sample sizes.

In this study, we did not systematically measure traditional stove usage with stove use monitors, which could strengthen the evidence that LPG uptake was near-exclusive and sustained, although the lack of observed traditional stove use by field workers supports high uptake of the intervention, which was also supported by low self-reported traditional stove post-intervention. It is possible that self-reported traditional stove use is under-reported due to response bias, which could partially explain the small number of higher $\text{PM}_{2.5}$ concentrations, especially for kitchen measurements. Those higher kitchen concentrations and exposure estimates during follow-up measurements may also be the result of other factors such as exposure to neighbor's smoke, trash-burning, or others. It is also possible that the measurements are not representative of normal exposures due to the Hawthorne effect, with more exclusive use of LPG during the 24–48 h of monitoring compared when HAP monitoring wasn't occurring (Lozier et al., 2016). Additionally, real-time compliance of wearing the exposure monitoring equipment was also not measured. The HAPIN main trial will provide an opportunity to look at LPG and traditional stove use over an 18-month period using sensors, along with exposure equipment-wearing compliance using accelerometers, and will thus be able to more objectively and accurately characterize stove use and exposure reductions observed over time (Johnson et al., 2020).

Another limitation of our study is the application of different $\text{PM}_{2.5}$ measurement instrumentation at the field sites. While the different instrumentation was a necessity given the resources and timing of the pilot study, protocols were the same across research sites in terms of filter handling, flow checks, microbalance specifications and weighing procedures, and others key to the collection of gravimetric $\text{PM}_{2.5}$ samples. Moreover, instrument comparison studies, including one by the HAPIN study team, demonstrated similar performance between ECMs, UPASes, and pump and filter systems (Arku et al., 2018; Burrows et al., 2020; Du et al., 2019; Pillarsetti et al., 2019a; Volckens et al., 2017).

We only measured primary cooking area $\text{PM}_{2.5}$ concentrations and only personal exposure experienced by pregnant woman. While these are expected to be representative of what a clean stove intervention could achieve in terms of kitchen and personal $\text{PM}_{2.5}$ reductions among primary cooks, future work will incorporate monitoring of multiple household environments, children and other household members in order to provide better $\text{PM}_{2.5}$ exposure estimates (Johnson et al., 2020; Liao et al., 2019).

5. Conclusion

The pilot results of this LPG intervention as implemented in three different countries show promise for reducing exposure to levels expected to deliver health benefits (Burnett et al., 2014; Steenland et al.,

2018). Ongoing and soon-to-be published clean cookstove studies conducted in Ghana will provide further evidence of the potential of similar interventions to improve health and sustain uptake (Carrión et al., 2018; Jack et al., 2015). Further research is needed to examine if effects and near-exclusive use are sustained over time (Rosenthal, 2015), if health benefits are in fact achieved (Ezzati and Baumgartner, 2016; Thakur et al., 2018), and how barriers to clean cooking can be reduced at household, community and national levels (Puzzolo et al., 2016), particularly among the poorest (Brooks et al., 2016).

Author statement

Jiawen Liao: Conceptualization, Methodology, Software, Formal analysis, Data Curation, Writing, Writing - Review & Editing, Miles Kirby: Conceptualization, Methodology, Software, Formal analysis, Investigation, Data, Writing Curation, Writing - Review & Editing, Supervision, Ajay Pillarisetti: Conceptualization, Methodology, Data Curation, Writing, Writing - Review & Editing, Ricardo Piedrahita: Methodology, Data Curation, Writing, Kalpana Balakrishnan: Conceptualization, Methodology, Writing - Review & Editing, Supervision, Project administration, Funding acquisition, Sankar Sambandam: Investigation, Data Curation, Supervision, Krishnendu Mukhopadhyay: Investigation, Data Curation, Supervision, Wenlu Ye: Formal analysis, Data Curation, Ghislaine Rosa: Conceptualization: Investigation, Supervision, Project administration, Funding acquisition, Fiona Majorin: Investigation, Supervision, Ephrem Dusabimana: Investigation, Supervision, Florian Ndagijimana: Investigation, Supervision, John P. McCracken: Conceptualization, Methodology, Writing - Review & Editing, Supervision, Project administration, Erick Mollinedo: Investigation, Data Curation, Oscar de Leon: Formal analysis, Investigation, Data Curation, Anaité Díaz-Artiga: Investigation, Supervision, Lisa M. Thompson: Writing - Review & Editing, Supervision, Project administration, Katherine Kearns: Data Curation, Luke Naeher: Methodology, Writing - Review & Editing, Joshua Rosenthal: Conceptualization, Writing - Review & Editing, Maggie L. Clark: Writing, Writing - Review & Editing, Kyle Steenland: Methodology, Writing, Writing - Review & Editing, Lance A. Waller: Formal analysis, Writing - Review & Editing, William Checkley: Conceptualization, Writing - Review & Editing, Project administration, Funding acquisition, Jennifer L. Peel: Conceptualization, Writing - Review & Editing, Project administration, Funding acquisition, Thomas Clasen: Conceptualization, Writing - Review & Editing, Project administration, Funding acquisition, Michael Johnson: Conceptualization, Methodology, Writing, Writing - Review & Editing, Project administration.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The HAPIN trial is funded by the U.S. National Institutes of Health (cooperative agreement 1UM1HL134590) in collaboration with the Bill & Melinda Gates Foundation [OPP1131279].

We would like to express our deep and sincere gratitude to the households that graciously invited us into their homes for this study. We also would like to thank the field teams, which worked diligently to collect the data presented here.

A multidisciplinary, independent Data and Safety Monitoring Board (DSMB) appointed by the National Heart, Lung, and Blood Institute (NHLBI) monitors the quality of the data and protects the safety of patients enrolled in the HAPIN trial. NHLBI DSMB: Nancy R Cook, Stephen Hecht, Catherine Karr (Chair), Joseph Millum, Nalini Sathiakumar, Paul K Whelton, Gail G Weinmann (Executive Secretary). Program

Coordination: Gail Rodgers, Bill & Melinda Gates Foundation; Claudia L Thompson, National Institute of Environmental Health Science; Mark J. Parascandola, National Cancer Institute; Marion Koso-Thomas, Eunice Kennedy Shriver National Institute of Child Health and Human Development; Joshua P Rosenthal, Fogarty International Center; Conception R Nierras, NIH Office of Strategic Coordination Common Fund; Katherine Kavounis, Dong- Yun Kim, Antonello Punturieri, and Barry S Schmetter, NHLBI. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the US National Institutes of Health or Department of Health and Human Services.

The study protocol has been reviewed and approved by institutional review boards (IRBs) or Ethics Committees at Emory University (00089799), Johns Hopkins University (00007464), Sri Ramachandra Institute of Higher Education and Research (IEC-N1/16/JUL/54/49) and the Indian Council of Medical Research – Health Ministry Screening Committee (5/8/4–30/(Env)/Indo-US/2016-NCD-I), Universidad del Valle de Guatemala (146-08-2016) and Guatemalan Ministry of Health National Ethics Committee (11–2016), the London School of Hygiene and Tropical Medicine (11664–5) and the Rwandan National Ethics Committee (No.357/RNEC/2018), and Washington University in St. Louis (201611159). The HAPIN trial has been registered with [ClinicalTrials.gov](https://clinicaltrials.gov) (Identifier NCT02944682).

[†]HAPIN Investigators: Vigneswari Aravindalochanan, Gloriose Bankundiye, Dana Boyd Barr, Alejandra Bussalleu, Devan Campbell, Eduardo Canuz, Adly Castañaza, Howard Chang, Yunyun Chen, Marilú Chiang, Rachel Craik, Victor G. Davila-Roman, Lisa de las Fuentes, Lisa Elon, Juan Gabriel Espinoza, Sarada Garg, Sarah Hamid, Stella Hartinger, Steven A. Harvey, Mayari Hengstermann, Ian Hennessee, Phabiola M. Herrera, Shakir Hossen, Penelope P. Howards, Lindsay Jaacks, Shirin Jabbarzadeh, Jacob Kremer, Pattie Lenzen, Amy E. Lovvorn, Jane Mbabazi, Eric McCollum, Rachel Meyers, Lawrence Moulton, Alexie Mukeshimana, Bernard Mutariyani, Durairaj Natesan, Azhar Nizam, Jean de Dieu Ntivuguruzwa, Aris Papageorghiou, Naveen Puttaswamy, Elisa Puzzolo, Ashlinn Quinn, Karthikeyan Dharmapuri Rajamani, Usha Ramakrishnan, Rengaraj Ramasami, Alexander Ramirez, P. Barry Ryan, Sudhakar Saidam, Jeremy A. Sarnat, Suzanne Simkovich, Sheela S. Sinharoy, Kirk R. Smith, Damien Swearing, Gurusamy Thangavel, Ashley Toenjes, Lindsay Underhill, Viviane Valdes, Kendra N. Williams, Bonnie N. Young.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envpol.2021.118198>.

References

- Alexander, D.A., Northcross, A., Karrison, T., Morhasson-Bello, O., Wilson, N., Atalabi, O. M., Dutta, A., Adu, D., Ibigbami, T., Olamijulo, J., Adepoju, D., Ojengbede, O., Olopade, C.O., 2018. Pregnancy outcomes and ethanol cook stove intervention: a randomized-controlled trial in Ibadan, Nigeria. *Environ. Int.* 111, 152–163. <https://doi.org/10.1016/J.ENVINT.2017.11.021>.
- Arku, R.E., Birch, A., Shupler, M., Yusuf, S., Hystad, P., Brauer, M., 2018. Characterizing exposure to household air pollution within the Prospective Urban Rural Epidemiology (PURE) study. *Environ. Int.* 114, 307–317. <https://doi.org/10.1016/j.envint.2018.02.033>.
- Baumgartner, J., Clark, S., Carter, E., Lai, A., Zhang, Y., Shan, M., Schauer, J.J., Yang, X., 2019. Effectiveness of a household energy package in improving indoor air quality and reducing personal exposures in rural China. *Environ. Sci. Technol.* 53, 9306–9316. <https://doi.org/10.1021/acs.est.9b02061>.
- Benka-Coker, M.L., Young, B.N., Keller, J.P., Walker, E.S., Rajkumar, S., Volckens, J., Good, N., Quinn, C., L'Orange, C., Weller, Z.D., Africano, S., Osorto Pinel, A.B., Peel, J.L., Clark, M.L., 2021. Impact of the wood-burning Justa cookstove on fine particulate matter exposure: a stepped-wedge randomized trial in rural Honduras. *Sci. Total Environ.* 767, 144369. <https://doi.org/10.1016/j.scitotenv.2020.144369>.
- Bonjour, S., Adair-Rohani, H., Wolf, J., Bruce, N.G., Mehta, S., Prüss-Ustün, A., Lahiff, M., Rehfuess, E.A., Mishra, V., Smith, K.R., 2013. Solid fuel use for household cooking: country and regional estimates for 1980–2010. *Environ. Health Perspect.* 121, 784–790. <https://doi.org/10.1289/ehp.1205987>.

- Brooks, N., Bhojvaid, V., Jeuland, M.A., Lewis, J.J., Patange, O., Pattanayak, S.K., 2016. How much do alternative cookstoves reduce biomass fuel use? Evidence from North India. *Resour. Energy Econ.* 43, 153–171. <https://doi.org/10.1016/j.reneneeco.2015.12.001>.
- Bruce, N., de Cuevas, R.A., Cooper, J., Enonchong, B., Ronzi, S., Puzzolo, E., MBatchou, B., Pope, D., 2018. The Government-led initiative for LPG scale-up in Cameroon: programme development and initial evaluation. *Energy Sustain. Dev.* 46, 103–110. <https://doi.org/10.1016/j.esd.2018.05.010>.
- Bruce, N.G., Pope, D., Rehfuess, E., Balakrishnan, K., Adair-Rohani, H., Dora, C., 2015. WHO indoor air quality guidelines on household fuel combustion: strategy implications of new evidence on interventions and exposure-risk functions. *Atmos. Environ.* 106, 451–457. <https://doi.org/10.1016/j.atmosenv.2014.08.064>.
- Burnett, R., Chen, H., Szyszko, M., Fann, N., Hubbell, B., Pope, C.A., Apté, J.S., Brauer, M., Cohen, A., Weichenthal, S., Coggins, J., Di, Q., Brunekreef, B., Frostad, J., Lim, S.S., Kan, H., Walker, K.D., Thurston, G.D., Hayes, R.B., Lim, C.C., Turner, M.C., Jerrett, M., Krewski, D., Gapstur, S.M., Diver, W.R., Ostro, B., Goldberg, D., Crouse, D.L., Martin, R.V., Peters, P., Pinault, L., Tjepkema, M., van Donkelaar, A., Villeneuve, P.J., Miller, A.B., Yin, P., Zhou, M., Wang, L., Janssen, N.A.H., Marra, M., Atkinson, R.W., Tsang, H., Quoc Thach, T., Cannon, J.B., Allen, R.T., Hart, J.E., Laden, F., Cesaroni, G., Forastiere, F., Weinmayr, G., Jaensch, A., Nagel, G., Concin, H., Spadaro, J.V., 2018. Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. In: *Proc. Natl. Acad. Sci. U. S. A.*, vol. 115, pp. 9592–9597. <https://doi.org/10.1073/pnas.1803222115>.
- Burnett, R.T., Pope, C.A.I., Ezzati, M., Olives, C., Lim, S.S., Mehta, S., Shin, H.H., Singh, G., Hubbell, B., Brauer, M., Anderson, H.R., Smith, K.R., Balmes, J.R., Bruce, N.G., Kan, H., Laden, F., Prüss-Ustün, A., Turner, M.C., Gapstur, S.M., Diver, W.R., Cohen, A., 2014. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ. Health Perspect.* <https://doi.org/10.1289/ehp.1307049>.
- Burrows, V.J., Piedrahita, R., Pillarisetti, A., Underhill, L.J., Fandiño-Del-Río, M., Johnson, M., Kephart, J.L., Hartinger, S.M., Steenland, K., Naehrer, L., Kearns, K., Peel, J.L., Clark, M.L., Checkley, W., Nambajimana, A., Verma, A., Lovvorn, A., Diaz, A., Papageorghiou, A., Toenjes, A., Quinn, A., Nizam, A., Ryan, B., Young, B., Barr, D., Goodman, D., Canuz, E., Puzzolo, E., McCollum, E., Mollinedo, E., Majorin, F., Ndagijimana, F., Rosa, G., Thangavel, G., Chang, H., Fuentes, I.S.P., Miranda, J.J., Ntuguruzwa, J. de D., Uwiziyimana, J., Peel, J., Sarnat, J., Liao, J., McCracken, J., Rosenthal, J., Espinoza, J.G., Campbell, J.M.P., Balakrishnan, K., Williams, K., Smith, K., Mukhopadhyay, K., Waller, L., Moulton, L., Jaacks, L., Elon, L., Thompson, L., Laws, M., Chiang, M., Howard, M., Crocker, M., Kirby, M., Puttaswamy, N., De Leon, O., Herrera, P., Craik, R., Merrick, R., Sambandam, S., Garg, S., Rajkumar, S., Gupton, S., Hossein, S., Sinharoy, S., Jabbarzadeh, S., Harvey, S., Simkovich, S., Clasen, T., Ramakrishnan, U., Davila-Roman, V.G., Aravindalochanan, V., Chen, Y., Sakas, Z., 2020. Comparison of next-generation portable pollution monitors to measure exposure to PM_{2.5} from household air pollution in Puno, Peru. *Indoor Air* 30, 445–458. <https://doi.org/10.1111/ina.12638>.
- Carrion, D., Dwommoh, R., Tawiah, T., Agyei, O., Agbokey, F., Twumasi, M., Mujtaba, M., Jack, D., Asante, K.P., 2018. Enhancing LPG adoption in Ghana (ELAG): a factorial cluster-randomized controlled trial to Enhance LPG Adoption & Sustained use. *BMC Publ. Health* 18, 689. <https://doi.org/10.1186/s12889-018-5622-3>.
- Checkley, W., Williams, K.N., Kephart, J.L., Fandiño-Del-Río, M., Steenland, K., Gonzales, G.F., Naehrer, L.P., Harvey, S.A., Moulton, L.H., Davila-Roman, V.G., Goodman, D., Tarazona-Meza, C., Miele, C.H., Simkovich, S., Chiang, M., Chartier, R.T., Koehler, K., 2020. Effects of a cleaner energy intervention on cardiopulmonary outcomes in Peru: a randomized controlled trial. *Am. J. Respir. Crit. Care Med.* <https://doi.org/10.1164/rccm.202006-2319oc>.
- Chillrud, S.N., Ae-Ngibise, K.A., Gould, C.F., Owusu-Agyei, S., Mujtaba, M., Manu, G., Burkart, K., Kinney, P.L., Quinn, A., Jack, D.W., Asante, K.P., 2021. The effect of clean cooking interventions on mother and child personal exposure to air pollution: results from the Ghana Randomized Air Pollution and Health Study (GRAPHS). *J. Expo. Sci. Environ. Epidemiol.* 1–16. <https://doi.org/10.1038/s41370-021-00309-5>.
- Clasen, T., Checkley, W., Peel, J.L., Balakrishnan, K., McCracken, J.P., Rosa, G., Thompson, L.M., Barr, D.B., Clark, M.L., Johnson, M.A., Waller, L.A., Jaacks, L.M., Steenland, K., Miranda, J.J., Chang, H.H., Kim, D.-Y., McCollum, E.D., Davila-Roman, V.G., Papageorghiou, A., Rosenthal, J.P., 2020. Design and rationale of the HAPIN study: a multicountry randomized controlled trial to assess the effect of liquefied petroleum gas stove and continuous fuel distribution. *Environ. Health Perspect.* 128, 047008. <https://doi.org/10.1289/EHP6407>.
- Du, Y., Wang, Q., Sun, Q., Zhang, T., Li, T., Yan, B., 2019. Assessment of PM_{2.5} monitoring using MicroPEM: a validation study in a city with elevated PM_{2.5} levels. *Ecotoxicol. Environ. Saf.* 171, 518–522. <https://doi.org/10.1016/j.ecoenv.2019.01.002>.
- Ezzati, M., Baumgartner, J.C., 2016. Household energy and health: where next for research and practice? *Lancet* 63–72. [https://doi.org/10.1016/S0140-6736\(16\)32506-5](https://doi.org/10.1016/S0140-6736(16)32506-5).
- Fandiño-Del-Río, M., Goodman, D., Kephart, J.L., Miele, C.H., Williams, K.N., Moazzami, M., Fung, E.C., Koehler, K., Davila-Roman, V.G., Lee, K.A., Nangia, S., Harvey, S.A., Steenland, K., Gonzales, G.F., Checkley, W., 2017. Effects of a liquefied petroleum gas stove intervention on pollutant exposure and adult cardiopulmonary outcomes (CHAP): study protocol for a randomized controlled trial. *Trials* 18, 518. <https://doi.org/10.1186/s13063-017-2179-x>.
- Funk, C., Peterson, P., Landsfeld, M., Pedrerros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., Michaelsen, J., 2015. The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. *Sci. Data* 2, 150066. <https://doi.org/10.1038/sdata.2015.66>.
- GBD 2017 Risk Factor Collaborators, 2018. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Stu. *Lancet* 392, 1923–1994. [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6).
- Gould, C.F., Urpelainen, J., 2018. LPG as a clean cooking fuel: adoption, use, and impact in rural India. *Energy Pol.* 122, 395–408. <https://doi.org/10.1016/j.enpol.2018.07.042>.
- Grieshop, A.P., Marshall, J.D., Kandlikar, M., 2011. Health and climate benefits of cookstove replacement options. *Energy Pol.* 39, 7530–7542. <https://doi.org/10.1016/j.enpol.2011.03.024>.
- Health Effects Institute, 2019. *State of Global Air 2019. Special Report* (Boston, MA).
- Jack, D.W., Asante, K.P., Wylie, B.J., Chillrud, S.N., Whyatt, R.M., Ae-Ngibise, K.A., Quinn, A.K., Yawson, A.K., Boamah, E.A., Agyei, O., Mujtaba, M., Kaali, S., Kinney, P., Owusu-Agyei, S., 2015. Ghana randomized air pollution and health study (GRAPHS): study protocol for a randomized controlled trial. *Trials* 16, 420. <https://doi.org/10.1186/s13063-015-0930-8>.
- Johnson, M., Edwards, R., Morawska, L., Smith, K.R., Nicas, M., 2014. WHO Guidelines for Indoor Air Quality: Household Fuel Combustion - Review 3: Model for Linking Household Energy Use with Indoor Air Quality.
- Johnson, M.A., Chiang, R.A., 2015. Quantitative guidance for stove usage and performance to achieve health and environmental targets. *Environ. Health Perspect.* 123, 820–826. <https://doi.org/10.1289/ehp.1408681>.
- Johnson, M.A., Garland, C.R., Jagoe, K., Edwards, R., Ndemere, J., Weyant, C., Patel, A., Kithinji, J., Wasirwa, E., Nguyen, T., Khoi, D.D., Kay, E., Scott, P., Nguyen, R., Yagnaraman, M., Mitchell, J., Derby, E., Chiang, R.A., Pennise, D., 2019. In-home emissions performance of cookstoves in asia and Africa. *Atmosphere* 10, 290. <https://doi.org/10.3390/atmos10050290>.
- Johnson, M.A., Steenland, K., Piedrahita, R., Clark, M.L., Pillarisetti, A., Balakrishnan, K., Peel, J.L., Naehrer, L.P., Liao, J., Wilson, D., Sarnat, J., Underhill, L.J., Burrows, V., McCracken, J.P., Rosa, G., Rosenthal, J., Sambandam, S., de Leon, O., Kirby, M.A., Kearns, K., Checkley, W., Clasen, T., 2020. Air pollutant exposure and stove use assessment methods for the household air pollution intervention network (HAPIN) trial. *Environ. Health Perspect.* 128, 47009. <https://doi.org/10.1289/EHP6422>.
- Kaali, S., Jack, D., Lee, A.G., 2020. The conundrum of cleaner cookstove interventions: necessary but insufficient? *Am. J. Respir. Crit. Care Med.* <https://doi.org/10.1164/rccm.202012-4353ed>.
- Katz, J., Tielsch, J.M., Khatry, S.K., Shrestha, L., Breyse, P., Zeger, S.L., Kozuki, N., Checkley, W., LeClerq, S.C., Mullany, L.C., 2020. Impact of improved biomass and liquid petroleum gas stoves on birth outcomes in rural Nepal: results of 2 randomized trials. *Glob. Health Sci. Pract.* 8. <https://doi.org/10.9745/GHSP-D-20-00011>.
- Kephart, J.L., Fandiño-Del-Río, M., Williams, K.N., Malpartida, G., Lee, A., Steenland, K., Naehrer, L.P., Gonzales, G.F., Chiang, M., Checkley, W., Koehler, K., 2021. Nitrogen dioxide exposures from LPG stoves in a cleaner-cooking intervention trial. *Environ. Int.* 146, 106196. <https://doi.org/10.1016/j.envint.2020.106196>.
- Kirby, M.A., Nagel, C.L., Rosa, G., Zambrano, L.D., Musafiri, S., Ndirabeya, J. de D., Thomas, E.A., Clasen, T., 2019. Effects of a large-scale distribution of water filters and natural draft rocket-style cookstoves on diarrhoea and acute respiratory infection: a cluster-randomized controlled trial in Western Province, Rwanda. *PLoS Med.* 16, e1002812. <https://doi.org/10.1371/journal.pmed.1002812>.
- Liao, Jiawen, Kirby, Miles A., Pillarisetti, Ajay, Piedrahita, Ricardo, Balakrishnan, Kalpana, Sambandam, Sankar, Mukhopadhyay, Krishnendu, Ye, Wenlu, Rosa, Ghislaine, Majorin, Fiona, Dusabimana, Ephrem, Ndagijimana, Florian, McCracken, John P., Mollinedo, Erick, de Leon, Oscar, Díaz-Artiga, Anaïté, Thompson, Lisa M., Kearns, Katherine A., Naehrer, Luke, Rosenthal, Joshua, Clark, Maggie L., Steenland, Kyle, Waller, Lance A., Checkley, William, Peel, Jennifer L., Clasen, Thomas, 2021. Replication Data for: LPG stove and fuel intervention among pregnant women reduce fine particle air pollution exposures in three countries: Pilot results from the HAPIN trial. UNC Dataverse V1 (UNF:6:Crazcb4/sjPKKcdNpmOYCO== [fileUNF]). <https://doi.org/10.15139/S3/XSGXUM>.
- Liao, J., McCracken, J.P., Piedrahita, R., Thompson, L., Mollinedo, E., Canuz, E., De León, O., Díaz-Artiga, A., Johnson, M., Clark, M., Pillarisetti, A., Kearns, K., Naehrer, L., Steenland, K., Checkley, W., Peel, J., Clasen, T.F., 2019. The use of bluetooth low energy Beacon systems to estimate indirect personal exposure to household air pollution. *J. Expo. Sci. Environ. Epidemiol.* 1–11. <https://doi.org/10.1038/s41370-019-0172-z>.
- Lozier, M.J., Sircar, K., Christensen, B., Pillarisetti, A., Pennise, D., Bruce, N.G., Stanistreet, D., Naehrer, L., Pilishvili, T., Farrar, J.L., Sage, M., Nyagol, R., Muoki, J., Wofchuck, T., Yip, F., 2016. Use of temperature sensors to determine exclusivity of improved stove use and associated household air pollution reductions in Kenya. *Environ. Sci. Technol.* 50, 4564–4571. <https://doi.org/10.1021/acs.est.5b06141>.
- Mortimer, K., Lesosky, M., Semple, S., Malava, J., Katundu, C., Crampin, A., Wang, D., Weston, W., Pope, D., Havens, D., Gordon, S.B., Balmes, J., 2020. Pneumonia and exposure to household air pollution in children under the age of 5 Years in rural Malawi: findings from the cooking and pneumonia study. *Chest* 158, 501–511. <https://doi.org/10.1016/j.chest.2020.03.064>.
- Mortimer, K., Ndamala, C.B., Naunje, A.W., Malava, J., Katundu, C., Weston, W., Havens, D., Pope, D., Bruce, N.G., Nyirenda, M., Wang, D., Crampin, A., Grigg, J., Balmes, J., Gordon, S.B., 2016. A cleaner burning biomass-fuelled cookstove intervention to prevent pneumonia in children under 5 years old in rural Malawi (the Cooking and Pneumonia Study): a cluster randomised controlled trial. *Lancet* 389, 167–175. [https://doi.org/10.1016/S0140-6736\(16\)32507-7](https://doi.org/10.1016/S0140-6736(16)32507-7).
- Ni, K., Carter, E., Schauer, J.J., Ezzati, M., Zhang, Y., Niu, H., Lai, A.M., Shan, M., Wang, Y., Yang, X., Baumgartner, J., 2016. Seasonal variation in outdoor, indoor,

- and personal air pollution exposures of women using wood stoves in the Tibetan Plateau: baseline assessment for an energy intervention study. *Environ. Int.* 94, 449–457. <https://doi.org/10.1016/j.envint.2016.05.029>.
- Pillarsetti, A., Carter, E., Rajkumar, S., Young, B.N., Benka-Coker, M.L., Peel, J.L., Johnson, M., Clark, M.L., 2019a. Measuring personal exposure to fine particulate matter (PM_{2.5}) among rural Honduran women: a field evaluation of the Ultrasonic Personal Aerosol Sampler (UPAS). *Environ. Int.* <https://doi.org/10.1016/j.envint.2018.11.014>.
- Pillarsetti, A., Ghorpade, M., Madhav, S., Dhongade, A., Roy, S., Balakrishnan, K., Sankar, S., Patil, R., Levine, D.I., Juvekar, S., Smith, K.R., 2019b. Promoting LPG usage during pregnancy: a pilot study in rural Maharashtra, India. *Environ. Int.* 127, 540–549. <https://doi.org/10.1016/j.envint.2019.04.017>.
- Pillarsetti, A., Vaswani, M., Jack, D., Balakrishnan, K., Bates, M.N., Arora, N.K., Smith, K.R., 2014. Patterns of stove usage after introduction of an advanced cookstove: the long-term application of household sensors. *Environ. Sci. Technol.* 48, 14525–14533. <https://doi.org/10.1021/es504624c>.
- Pope, D., Bruce, N., Dherani, M., Jagoe, K., Rehfuess, E., 2017. Real-life effectiveness of ‘improved’ stoves and clean fuels in reducing PM_{2.5} and CO: systematic review and meta-analysis. *Environ. Int.* 101, 7–18. <https://doi.org/10.1016/j.envint.2017.01.012>.
- Puzzolo, E., Pope, D., Stanistreet, D., Rehfuess, E.A., Bruce, N.G., 2016. Clean fuels for resource-poor settings: a systematic review of barriers and enablers to adoption and sustained use. *Environ. Res.* 146, 218–234. <https://doi.org/10.1016/j.envres.2016.01.002>.
- Quansah, R., Semple, S., Ochieng, C.A., Juvekar, S., Armah, F.A., Luginaah, I., Emina, J., 2017. Effectiveness of interventions to reduce household air pollution and/or improve health in homes using solid fuel in low-and-middle income countries: a systematic review and meta-analysis. *Environ. Int.* 103, 73–90. <https://doi.org/10.1016/j.envint.2017.03.010>.
- Quinn, A.K., Puzzolo, E., Dickinson, K., Sturke, R., Jack, D.W., Mehta, S., Shankar, A., Sherr, K., Rosenthal, J.P., 2018. An analysis of efforts to scale up clean household energy for cooking around the world. *Energy Sustain. Dev.* 46, 1–10. <https://doi.org/10.1016/j.esd.2018.06.011>.
- Rosenthal, J., 2015. The real challenge for cookstoves and health: more evidence. *EcoHealth* 12, 8–11. <https://doi.org/10.1007/s10393-014-0997-9>.
- Rosenthal, J., Quinn, A., Grieshop, A.P., Pillarsetti, A., Glass, R.I., 2018. Clean cooking and the SDGs: integrated analytical approaches to guide energy interventions for health and environment goals. *Energy Sustain. Dev.* 42, 152–159. <https://doi.org/10.1016/j.esd.2017.11.003>.
- Ruiz-Mercado, I., Masera, O., 2015. Patterns of stove use in the context of fuel–device stacking: rationale and implications. *EcoHealth* 12, 42–56. <https://doi.org/10.1007/s10393-015-1009-4>.
- Sambandam, S., Mukhopadhyay, K., Sendhil, S., Ye, W., Pillarsetti, A., Thangavel, G., Natesan, D., Ramasamy, R., Natarajan, A., Aravindalochanan, V., Vinayagamoorathi, A., Sivavadevel, S., Uma Maheswari, R., Balakrishnan, L., Gayatri, S., Nargunanathan, S., Madhavan, S., Puttaswamy, N., Garg, S.S., Quinn, A., Rosenthal, J., Johnson, M., Liao, J., Steenland, K., Piedrahita, R., Peel, J., Checkley, W., Clasen, T., Balakrishnan, K., 2020. Exposure contrasts associated with a liquefied petroleum gas (LPG) intervention at potential field sites for the multi-country household air pollution intervention network (HAPIN) trial in India: results from pilot phase activities in rural Tamil Nadu. *BMC Publ. Health* 20, 1–13. <https://doi.org/10.1186/s12889-020-09865-1>.
- Sedgwick, P., 2014. Before and after study designs. *BMJ* 349, g5074. <https://doi.org/10.1136/bmj.g5074>.
- Shen, G., Hays, M.D., Smith, K.R., Williams, C., Faircloth, J.W., Jetter, J.J., 2018. Evaluating the performance of household liquefied petroleum gas cookstoves. *Environ. Sci. Technol.* 52, 904–915. <https://doi.org/10.1021/acs.est.7b05155>.
- Shupler, M., Godwin, W., Frostad, J., Gustafson, P., Arku, R.E., Brauer, M., 2018. Global estimation of exposure to fine particulate matter (PM_{2.5}) from household air pollution. *Environ. Int.* 120, 354–363. <https://doi.org/10.1016/j.envint.2018.08.026>.
- Shupler, M., Hystad, P., Birch, A., Miller-Lionberg, D., Jeronimo, M., Arku, R.E., Chu, Y. L., Mushtaha, M., Heenan, L., Rangarajan, S., Seron, P., Lanas, F., Cazor, F., Lopez-Jaramillo, P., Camacho, P.A., Perez, M., Yeates, K., West, N., Ncube, T., Ncube, B., Chifamba, J., Yusuf, R., Khan, A., Hu, B., Liu, X., Wei, L., Tse, L.A., Mohan, D., Kumar, P., Gupta, R., Mohan, I., Jayachitra, K.G., Mony, P.K., Rammohan, K., Nair, S., Lakshmi, P.V.M., Sagar, V., Khawaja, R., Iqbal, R., Kazmi, K., Yusuf, S., Brauer, M., 2020. Household and personal air pollution exposure measurements from 120 communities in eight countries: results from the PURE-AIR study. *Lancet Planet. Health* 4, e451–e462. [https://doi.org/10.1016/S2542-5196\(20\)30197-2](https://doi.org/10.1016/S2542-5196(20)30197-2).
- Smith, K.R., Mccracken, J.P., Thompson, L., Edwards, R., Shields, K.N., Canuz, E., Bruce, N.G., 2010. Personal child and mother carbon monoxide exposures and kitchen levels: methods and results from a randomized trial of woodfired chimney cookstoves in Guatemala (RESPIRE). *J. Expo. Sci. Environ. Epidemiol.* 20, 406–416. <https://doi.org/10.1038/jes.2009.30>.
- Snider, G., Carter, E., Clark, S., Tseng, J., Tzu, W., Yang, X., Ezzati, M., Schauer, J.J., Wiedinmyer, C., Baumgartner, J., 2018. Impacts of stove use patterns and outdoor air quality on household air pollution and cardiovascular mortality in southwestern China. *Environ. Int.* 117, 116–124. <https://doi.org/10.1016/j.envint.2018.04.048>.
- Steenland, K., Pillarsetti, A., Kirby, M., Peel, J., Clark, M., Checkley, W., Chang, H.H.H., Clasen, T., 2018. Modeling the potential health benefits of lower household air pollution after a hypothetical liquefied petroleum gas (LPG) cookstove intervention. *Environ. Int.* 111, 71–79. <https://doi.org/10.1016/j.envint.2017.11.018>.
- Thakur, M., Nuyts, P.A.W., Boudewijns, E.A., Flores Kim, J., Faber, T., Babu, G.R., Van Schayck, O.C.P., Been, J.V., 2018. Impact of improved cookstoves on women’s and child health in low and middle income countries: a systematic review and meta-analysis. *Thorax* 73, 1026–1040. <https://doi.org/10.1136/thoraxjnl-2017-210952>.
- Volckens, J., Quinn, C., Leith, D., Mehaffy, J., Henry, C.S., Miller-Lionberg, D., 2017. Development and evaluation of an ultrasonic personal aerosol sampler. *Indoor Air* 27, 409–416. <https://doi.org/10.1111/ina.12318>.
- WHO, 2016. *Burning Opportunity: Clean Household Energy for Health, Sustainable Development, and Wellbeing of Women and Children* (Geneva, Switzerland).
- WHO, 2014. *Indoor Air Quality Guidelines: Household Fuel Combustion* (Geneva, Switzerland).
- Williams, K.N., Thompson, L.M., Sakas, Z., Hengstermann, M., Quinn, A., Díaz-Artiga, A., Thangavel, G., Puzzolo, E., Rosa, G., Balakrishnan, K., Peel, J., Checkley, W., Clasen, T.F., Miranda, J.J., Rosenthal, J.P., Harvey, S.A., 2020. Designing a comprehensive behaviour change intervention to promote and monitor exclusive use of liquefied petroleum gas stoves for the Household Air Pollution Intervention Network (HAPIN) trial. *BMJ Open* 10, e037761. <https://doi.org/10.1136/bmjopen-2020-037761>.