

Mineral micronutrient status and spatial distribution among the Ethiopian population

Adamu Belay^{1,2}, Dawd Gashu^{1*}, Edward J. M. Joy³, R. Murray Lark⁴, Christopher Chagumaira⁴, Dilnesaw Zerfu², E. Louise Ander⁵, Scott D. Young⁴, Elizabeth H. Bailey⁴ & Martin R. Broadley^{4,6}

¹Center for Food Science and Nutrition, Addis Ababa University, P.O.Box 1176, Addis Ababa, Ethiopia

²Food Science and Nutrition Research Directorate, Ethiopian Public Health Institute, Gulele Sub City, P.O. Box 1242, Addis Ababa, Ethiopia

³Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK

⁴School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, UK

⁵Inorganic Geochemistry, Centre for Environmental Geochemistry, British Geological Survey, Nottingham NG12 5GG, UK

⁶Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK

***Corresponding author:** Dr Dawd Gashu; dawd.gashu@aau.edu.et

Short title: Mineral micronutrient nutrition



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114522000319

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Multiple micronutrient deficiencies are widespread in Ethiopia. However, the distribution of selenium and zinc deficiency risks has previously shown evidence of spatially dependent variability, warranting the need to explore this aspect for wider micronutrients. Here, blood serum concentrations for calcium (Ca), magnesium (Mg), cobalt (Co), copper (Cu), and molybdenum (Mo) were measured (n 3,102) on samples from the Ethiopian National Micronutrient Survey. Geostatistical modelling was used to test spatial variation of these micronutrients for women of reproductive age (WRA), who represent the largest demographic group surveyed (n 1,290). Median serum concentrations were 8.6 mg dL⁻¹ for Ca, 1.9 mg dL⁻¹ for Mg, 0.4 µg L⁻¹ for Co, 98.8 µg dL⁻¹ for Cu, and 0.2 µg dL⁻¹ for Mo. The prevalence of Ca, Mg and Co deficiency was 41.6%, 29.2% and 15.9%, respectively; Cu and Mo deficiency prevalence was 7.6% and 0.3%, respectively. A higher prevalence of Ca, Cu and Mo deficiency was observed in north western, Co deficiency in central, and Mg deficiency in north eastern parts of Ethiopia. Serum Ca, Mg, and Mo concentrations show spatial dependencies up to 140–500 km; however, there was no evidence of spatial correlations for serum Co and Cu concentrations. These new data indicate the scale of multiple mineral micronutrient deficiency in Ethiopia and the geographical differences in the prevalence of deficiencies suggesting the need to consider targeted responses during the planning of nutrition intervention programs.

Keywords: Ethiopia, Geospatial prediction, Micronutrient surveys, Mineral micronutrients, Spatial variation, Variogram analysis

Background

Multiple micronutrient deficiencies remain widespread globally with particular attention on Fe, folate, vitamin A, iodine, and Zn through the analysis of biomarkers for population-level surveillance ⁽¹⁾. For many other micronutrients, there is limited knowledge of the prevalence of deficiencies based on biomarkers. This is likely due to the analytical challenges of measuring multiple micronutrients on small sample volumes, and a lack of agreed thresholds/cut-offs and potential adjustments (e.g. for inflammation) for some micronutrients ^(2,3). However, deficiencies of multiple micronutrients that are not routinely measured in national micronutrient surveys are likely to be widespread. For example, the deficiency risk of Ca based on food supply data is greater (54%) than for Zn (40%) in sub-Saharan Africa ^(2,4). On the other hand, the deficiency risk for Cu is 1% and less for Mg based on food supply data ⁽²⁾.

Biomarker based studies reporting mineral micronutrient status of the Ethiopian population except for Zn and Fe are limited. However, available reports suggest presence of multiple mineral micronutrient deficiency. A study analyzing serum concentration of Mg, Zn, Se, Ca, Fe, Co and Mo shows presence of high prevalence of Se (62%) and Zn (47%) deficiency among school age children from Northwest Ethiopia. However, no or few children had Mg (2%), Ca, Fe, Co, Mo and Cu/Zn deficiency⁽⁵⁾. In addition, a nationally representative study reported that 72% of the Ethiopian population are Zn deficient ⁽⁶⁾.

Nutrient deficiency is not always due to lack of adequate intakes. Minerals may also antagonize the absorption and transportation of other minerals. Higher intake levels of one micronutrient in the diet or from supplements may competitively inhibit absorption of other micronutrients. This often occurs between nutrients possessing similar chemical characteristics as they can share similar receptors or transporting proteins⁽⁷⁾. In addition, deficiency of a micronutrient can affect the biological function of another micronutrient through disruption of the metabolic pathways. For example, Se deficiency can reduce the activity of the Se-dependant enzyme involved in the conversion of the biologically inactive thyroxine (T₄) in to triiodothyronine (T₃) ⁽⁸⁾ indicating the need to concomitantly address deficiency of these two micronutrients. On the other hand, high dietary Ca concentration in the luminal environment reduces the rate of Mg absorption ⁽⁹⁾, while Zn and Cu compete for absorption in the intestine ⁽¹⁰⁾. Such available evidence of nutrient-nutrient interactions

suggests that data on multiple micronutrient status of populations are important to understand not only the magnitude of the nutritional problem but also the context of the nutritional problem including presence of antagonistic or synergistic interactions to effectively design interventions.

The aim of this study was to determine the prevalence of multiple micronutrient deficiencies in Ethiopia and explore potential spatial dependencies of this variation based on each of the serum biomarkers. Prevalence of a deficiency is an important statistic to inform national policies, but spatial dependence is also significant. In the presence of spatial dependence, it may be expected that prevalence of deficiency changes from one location to another, and if spatial dependence is seen over sufficiently long distances, then this could imply that effective interventions differ between regions of a country, and that spatial information is needed to target them ^(6,11,12). The focus is Ethiopia, where a National Micronutrient Survey (ENMS) was conducted in 2015. From the ENMS, the prevalence of Fe deficiency among young children (YC), school age children (SAC), and women of reproductive age (WRA), as measured by ferritin and adjusted for inflammation, were 17.8%, 9.1% and 10.0%, respectively ⁽¹³⁾. Subsequently, multi-elemental analyses of serum samples were reported, both for Se⁽¹¹⁾ and for Zn which included adjustments for inflammation ⁽⁶⁾. These two studies provided strong additional evidence that Se and Zn deficiencies are widespread in Ethiopia, consistent with previous cross-sectional studies ⁽¹⁴⁾. Furthermore, the prevalence of Se and Zn deficiency was spatially dependent, based on data for women of reproductive age (WRA), who represent the largest demographic group in the ENMS ^(6,11). Thus, people living in some areas are likely to have a greater risk of micronutrient deficiency. For Se, spatial dependencies up to 200 km were reported ⁽¹¹⁾. Serum Se concentrations correlated positively with grain Se concentration, including a strong trend of increasing Se status from west to east Amhara linked to soil properties and landscape factors ⁽¹²⁾. For Zn, spatial dependencies was found over shorter distances, of up to 45 km ⁽⁶⁾. In the current study, the focus is on Ca, Mg, Co, Cu, and Mo. This is because these elements could be reliably analysed in multi-element ICP-MS, and because there are accepted thresholds to indicate deficiency risks ⁽¹⁵⁾. In addition, serum Zn concentration data reported previously ⁽⁶⁾ were used to quantify Cu:Zn ratios, as a potential indicator for infection ^(16,17).

Materials and Methods

Study design and sample population

The ENMS was designed to cover all regions and administrative cities (Addis Ababa and Dire Dawa) of Ethiopia (Figure 1). The design of the ENMS is explained in detail elsewhere ^(6,11,13,18). Briefly, the ENMS was a population-based, cross-sectional survey including Young Children (YC, aged 6–59 months, n=1,100), School Age Children (SAC, aged 5–15 years, n=1,500), non-pregnant (self-reported) Women of Reproductive Age (WRA, aged 15–49 years, n=1,600) and Men (aged 15–54 years, n=500) conducted between March and July 2015. The ENMS enumeration areas (EAs), or clusters, are geographic areas defined by the Central Statistics Agency (CSA) for the Ethiopia Population and Housing Census ⁽¹⁹⁾. For each region, the EAs were selected based on standard probability proportional to size (PPS). EAs contain on average 181 households (150 to 200) ⁽¹⁹⁾. Within each selected EA, 11 households were randomly selected for enumeration.

This study used socio-demographic data collected from households in the ENMS, and newly-generated data on micronutrients including Ca, Mg, Co, Cu, and Mo concentrations measured in archived serum samples from the ENMS. The new micronutrient data were obtained only for those individuals from the ENMS survey for whom socio-demographic data were available, and where there was at least 0.3 mL of archived serum sample. Serum Zn concentration data were taken from

Belay et al. ⁽⁶⁾. A total of 3,102 sample were included in the analysis.

Ethics

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the National Research Ethics Review Committee at the Ministry of Science and Technology, Ethiopia (Reference 3.10/433/06). Written informed consent and assent were obtained for adult and child participants, respectively. This study was also approved by the Research Ethical Review Committee at the Ethiopian Public Health Institute (Protocol EPHI-IRB-140-2018). Archived serum samples were transferred from storage at EPHI to the University of Nottingham, UK for analysis under a Material Transfer Agreement.

Data collection and analysis

Socio-demographic data

The ENMS collected socio-demographic information using pre-tested questionnaires. The data collectors were trained for three weeks on data collection and quality control following pilot testing in a cluster which was not included in the final ENMS. Questionnaires were refined after the pilot testing and before the survey data collection began.

Collection, processing and analysis of serum micronutrients

Detail information about blood collection has been described previously^(11,13,18). Briefly, trained phlebotomist collected blood samples from the antecubital veins of participants following World Health Organization (WHO) blood collection guidelines⁽²⁰⁾. Blood and serum samples were aliquoted for a range of analyses⁽¹³⁾. Backup sample aliquots were kept at -80°C to enable analytical cross-checks and additional analyses such as those reported previously^(6,11) and in this study.

Serum micronutrient analysis

Elemental concentrations of Ca, Mg, Co, Cu, and Mo in serum samples were determined using inductively coupled plasma-mass spectrometry (ICP-MS; Thermo Fisher Scientific iCAPQ, Thermo Fisher Scientific, Bremen, Germany) as reported previously^(6,11). Samples were introduced, via a single line, from an autosampler incorporating an ASXpress™ rapid uptake module (Cetac ASX-520, Teledyne Technologies Inc., Omaha, NE, USA) through a perfluoroalkoxy (PFA) Microflow PFA-ST nebulizer (Thermo Fisher Scientific, Bremen, Germany). All samples and external multi-element calibration standards were diluted as 0.3 mL added to 6 mL of a solution containing (i) 0.5% HNO₃ (Primar Plus grade), (ii) 2.0% methanol (Fisher Scientific UK Ltd, Loughborough, UK) and (iii) three internal standards including ⁷²Ge (10 µg L⁻¹), ¹⁰³Rh (5 µg L⁻¹), ¹⁹³Ir (5 µg L⁻¹) (SPEX Certiprep Inc., Metuchen, NJ, USA). Calibration standards included: (i) Co, Cu, and Mo (0, 20, 40, 100 µg L⁻¹; Claritas-PPT grade CLMS-2; SPEX Certiprep Inc., Metuchen, NJ, USA); Ca and Mg (0, 10, 20, 30 mg L⁻¹; PlasmaCAL, SCP Science, France). The ICP-MS was operated in 'collision-reaction cell mode', with kinetic energy discrimination, using He for all elements. The quadrupole dwell time was 0.1 s and 150 scans were used to obtain an average measurement of signal intensity (CPS; counts-per-second).

The limit of detection (LOD) for all elements was measured as $3 \times$ standard deviation of 10 operational blanks; the limits of detection (LOD) of Ca, Mg, Co, Cu, and Mo were 91.3, 3.12, 0.013, 0.407, and $0.064 \mu\text{g L}^{-1}$, respectively. The limit of quantification (LOQ) was calculated as $10 \times$ standard deviation. Values of LOQ ($\mu\text{g L}^{-1}$) were 304 for Ca, 10.4 for Mg, 0.044 for Co, 1.36 for Cu, and 0.212 for Mo.

Accuracy was verified by the use of two appropriate certified reference materials (CRM) (Seronorm™ L-1 (Lot 1801802) and Seronorm™ L-2 (Lot 1801803)); Nycomed Pharma AS, Billingstad, Norway); these were typically run at the start and the end of sample batch runs and they were prepared identically to samples and calibration standards. A total of 24 individual CRM analyses were undertaken for both L-1 and L-2. The average recoveries were recorded for Seronorm™ L-1 and Seronorm™ L-2. The average recoveries (%) were 89 for Ca, 100 for Mg, 100 for Co, 88 for Cu, and 113 for Mo in Seronorm™ L-1 CRM and in Seronorm™ L-2 CRM, the average recoveries (%) were 92 for Ca, 102 for Mg, 101 for Co, 85 for Cu, and 95 for Mo. All micronutrients in CRM were accredited except for Mo.

Data analysis

From the total of 3,373 samples, 271 observations were excluded due to missing GPS coordinates, missing information on time of food intake during data collection, or a lack of socio-demographic data and analytical outlier datum ($3 \times \text{IQR}$)⁽²¹⁾. Thus, 3,102 observations were included for descriptive statistical analyses of the prevalence of micronutrient deficiencies using STATA (Version 14.0, StataCorp LLP, Texas, USA). Prevalence of Zn deficiency among all demographic group in Ethiopia were previously determined⁽⁶⁾ as suggested by King et al.⁽²²⁾. Cut off values for Ca, Mg, Co, Cu and Mo deficiency were defined at their serum levels of 8.4 mg dL^{-1} , 1.8 mg dL^{-1} , $0.22 \mu\text{g L}^{-1}$, $75 \mu\text{g dL}^{-1}$ and $0.02 \mu\text{g dL}^{-1}$, respectively, for all demographic groups⁽¹⁵⁾. Ratio of Ca:Mg⁽⁹⁾ and Cu:Zn⁽¹⁶⁾ were calculated as indicators for chronic disease and presence of inflammation in the population, respectively. Pearson's correlation coefficients were estimated for exploration of association between pairs of elements.

Spatial determination of micronutrient deficiencies

Geostatistical modelling was undertaken to reveal the spatial dependence of micronutrient deficiencies, using methods described previously^(6,11,23), and more detail on the analytical methods is presented in the Supplementary Information S1. Briefly, ordinary kriging (OK) was used to predict serum concentrations at unsampled location by interpolation from the

measured data. In OK the interpolation is a weighted linear combination of the observations, with weights computed to minimize the expected squared prediction error (kriging variance). This measure of uncertainty of the predictions was calculated (Supplementary Information Table S1). The OK prediction is the mean of the prediction distribution at the target location, and the kriging variance is the variance of the prediction distribution. OK predictions were computed on the nodes of a 60-metre square grid. These predictions, and the kriging variances, were then presented as a map using ArcGIS (10.4.1). For all micronutrients, the spatial analysis is focused on WRA because there were greater numbers of WRA and cover a large number of EAs (320 out of a total of 346 EAs). The statistical summary for serum micronutrients of WRA is shown in Supplementary Information Figure S1.

Results

Population demographic and micronutrient data

This study included 474 YC, 935 SAC, 403 men and 1,290 WRA, with highest participants from Oromia region and lowest number of participants from Dire Dawa Administrative city. The mean age of YC, SAC, men and WRA was 3.1 ± 1.0 , 9.6 ± 2.7 , 30.1 ± 10.5 and 28.2 ± 8.9 year, respectively. The detailed demographic information is reported elsewhere ^(6,11).

The median serum concentration of micronutrients in the study groups are shown in Table 1. Median serum concentrations of Ca, Mg, Co, Cu, and Mo were 8.6 mg dL^{-1} , 1.9 mg dL^{-1} , $0.4 \mu\text{g L}^{-1}$, $98.8 \mu\text{g dL}^{-1}$, and $0.2 \mu\text{g dL}^{-1}$, respectively; ratios of Ca:Mg and Cu:Zn were 4.4 and 1.8, respectively. The median serum concentrations of micronutrients differed between the regions, with the highest serum Ca and Mg concentrations observed in Dire Dawa followed by Amhara. Serum Ca:Mg ratio was high (>4) in all regions. Serum Co concentration was highest in Benishangul-Gumuz and lowest in Addis Ababa. Serum Cu concentration was highest in Gambela and lowest in SNNP. Younger children had lower serum concentrations of all micronutrients except Cu compared to their counterparts. Young children had the largest serum Cu:Zn ratio. Urban residents had larger serum micronutrient concentrations than rural residents, except for Cu. There were little differences in median micronutrient status between groups based on the educational status of the head of household.

Of the total study population serum samples 41.6%, 29.2%, 15.9%, 7.6% and 0.3% were Ca, Mg, Co, Cu and Mo deficient, respectively (Table 2). Among the study groups 34.9% of the

population was deficient in at least one of the micronutrients, 22.4% and 20.7% of study population showed two and three coexisting micronutrients deficiencies, respectively. Among demographic groups, a high micronutrient deficiency burden was observed in young children, except for Mo.

The correlation coefficients indicated presence of association between pairs of micronutrients, namely Ca-Mg ($r=0.70$), Cu-Mg ($r=0.26$), Zn-Mg ($r=0.31$), Ca-Zn ($r=0.42$), Cu-Zn ($r=0.12$) as indicated in Table 3.

Spatial analysis of WRA serum micronutrient data

The WRA comprised the largest demographic group within the analysed samples, so these were used to explore whether spatial variation exists within the survey data. Variogram values were estimated with the alternative estimators and plotted against lag distance, along with models fitted by weighted least squares using R software. If the exponential variogram model for estimates by a particular estimator had a smaller value of AIC than the corresponding pure nugget model, then this exponential model was subject to cross-validation. A model (from one of the estimators) was selected by reference to the value of the median standardized squared prediction error. There was no evidence to support a model of spatial dependence for the data on Co, Cu and the Cu:Zn ratio. However, the AIC for the spatially dependent model was smaller than for the pure nugget model in the case of Ca, Mg, Mo and the Ca: Mg ratio. Furthermore, for all these variables the model fitted to the estimates by Matheron's estimator were selected as the median SSPE for the cross validation of this model fell within the 95% confidence interval for this statistic under a valid model in all cases. This indicates that there is no basis for mapping spatial variation in the Co, Cu and Cu:Zn serum data, and these data are not further presented in this section. In contrast, Ca, Mg, Mo and the Ca: Mg ratio showed spatial dependence at distances from 140–500 km, with Ca:Mg having the shortest range of spatial dependence, at which the variogram is approximately flat (150 km) and Ca showing spatial dependence over the longest distances (over 500 km).

The kriging interpolation maps (Figure 2a-d) show higher serum Ca and Mg concentrations in north western and south eastern areas, including large parts of Amhara, Dire Dawa and eastern Oromia regions; shaded in deep purple colors. Lower concentrations of these elements were predicted in south western, north eastern areas, and along the Rift Valley, including parts of Tigray, Benishangul-Gumuz, Gambella, and western and southern parts of

Oromia regions; shaded in light purple colours. The highest values of serum Ca:Mg ratio were observed in north eastern, south western, south eastern and central parts of Ethiopia, appearing as disconnected patches of deep purple colors. Higher serum Mo concentrations can be seen in the Rift Valley area. The lower serum Mo concentrations were observed in north western and south eastern parts of the country including Amhara, Benishangul-Gumuz, east and south Oromia regions.

The kriging variance of serum Ca, Mg, Ca:Mg ratio and Mo concentrations for WRA are shown in Figure 3a-d; these are the expected kriging variance computed based on the variogram model. These are the measures of prediction uncertainty which is minimized by ordinary kriging. Interpolation error is expected because of the spatial variation of our target variables. These kriging variances allow us to visualize the uncertainties and avoid arbitrary decisions as to where spatial prediction is reliable from a particular distribution of observations. For all micronutrients, there is greater uncertainty in some parts of the country, shaded in light grey, because of the sparse distribution of observations ⁽²⁴⁾. In these circumstances, the predicted value tends to the mean of the observations, and the kriging variance becomes large (light grey areas). If decisions on nutrition intervention programs are needed where the kriging variance is large, then local sampling would be essential.

The combined predicted concentration and uncertainty estimations outputs from the geostatistical analysis allows us to visualize occurrence that the probability of both risk micronutrient deficiency or that there are high micronutrients concentration at a national scale. Figure 4a-d shows the probability that the mean serum Ca, Mg, Ca:Mg ratio and Mo concentration of WRA in an EA is below a threshold of 8.4 mg dL⁻¹, 1.8 mg dL⁻¹, 4 and 0.02 µg dL⁻¹, respectively. Because continuous probabilities are not always easily interpreted by diverse stakeholders, we also present these probabilities using the “calibrated phrases” of the Intergovernmental Panel for Climate Change (IPCC) ⁽²⁵⁾, which are designed for the communication of uncertain information to data users unfamiliar with probability. This approach has been used elsewhere ^(11,23,26).

From the Ca and Mg probability maps (Figure 4a-d), we can see that in the north west (Amhara region) and east of Oromia and Dire Dawa regions, WRA are ‘unlikely’ through to ‘exceptionally unlikely’ to be Ca and Mg deficient. However, the likelihood of Ca and Mg deficiencies are generally greater in some north eastern and western areas of Ethiopia, which

may increase the risk of some diseases like pre-eclampsia and hypomagnesemia, respectively. In addition, from the Ca:Mg ratio probability map we observed that WRA in most parts of the country are 'unlikely' through 'exceptionally unlikely' to have a low Ca:Mg ratio (<4). Beyond the Rift Valley areas, WRA are 'likely' to 'virtually certain' to have low serum Mo concentrations.

Discussion

This study revealed a high prevalence of deficiencies of Ca, Mg and Co but low prevalence of deficiencies of Cu and Mo in Ethiopia. The prevalence of micronutrient deficiencies varied spatially for Ca, Mg and Mo with a high burden of deficiency observed in young children. A greater prevalence of micronutrients deficiencies was observed among those living in rural areas, except for Co.

In the previous study based on food supplies data, the mean risk of Ca and Mg deficiencies for Eastern parts of Africa was 69% and 1.2%, respectively⁽²⁾. Furthermore, the Ethiopian food consumption survey result revealed that the mean Ca intake for women was 501 mg d⁻¹ at the national level which is far less than the cutoff (900 mg d⁻¹) defined for a low-intake indicating that about 88% of the population are at risk of Ca-deficiency⁽²⁷⁾. In the present study, 41% and 29% of the study population were deficient in Ca and Mg, respectively, based on serum concentration thresholds. However, it is known that biomarker data of micronutrients is more reliable compared to estimation of deficiency risk using food supply data. Young children had the highest prevalence of Ca (90%) and Mg (76.1%) deficiency, although these are based on adult thresholds. Countries with low dietary intake of dairy products have low Ca status⁽²⁸⁾. In Ethiopia, milk and dairy product consumption is relatively low⁽²⁷⁾. Rural residents were more likely to be below the thresholds for serum Ca and Mg concentrations than urban residents, which may reflect the intake of flesh, dairy and eggs which are good source of these minerals are higher in urban than in rural areas⁽²⁹⁾. Serum Ca and Mg shows strong positive association (r=0.70).

Calcium and Mg are essential for bone formation and for numerous kinds of metabolisms in human body⁽³⁰⁾. They are intimately related to each other and have been reported to collectively influence metabolic functions such as hypertension and insulin resistance^(31,32). A study in a Chinese population revealed that individuals with metabolic syndrome had higher whole blood levels of Mg but lower Ca compared to healthy controls⁽³⁰⁾. A study in

Nigeria also shows that pre-eclampsia prevalence was higher in women with a Ca:Mg ratio greater than 2⁽³³⁾. In our study, the mean and median Ca:Mg ratio was greater than 4, which may increase the risk of developing chronic diseases including hypertension, cardio-vascular and diabetes^(34,35).

Copper and Zn are involved in numerous aspects of cellular metabolism because they allow many critical enzymes to function properly⁽¹⁰⁾. Copper deficiency is relatively low among all of the demographic groups in Ethiopia based on this threshold, and consistent with food supply data⁽²⁾. In contrast, Zn deficiency is widespread in the country⁽⁶⁾, with YC more likely to be Zn deficient than SC, men and WRA. This observation agrees with Ethiopian Food Consumption survey which reported 95% of YC were susceptible for Zn deficiency based on food intake⁽³⁶⁾.

An elevated serum Cu:Zn ratio indicates the presence of inflammation in addition to C-reactive protein (CRP) and α 1-acid glycoprotein (AGP) biomarkers. This is because inflammation and infection can reduce serum Zn during the acute phase response due to the redistribution of the serum Zn into liver and other tissues. Furthermore, acute infection leads to increased serum Cu concentration⁽¹⁶⁾. Both responses cause serum Cu:Zn ratio to increase. In a previous study on school age children in Ethiopia, the Cu:Zn ratio was greater than 2 indicating a high prevalence of infection⁽⁵⁾. Similarly, in the present study, the Cu:Zn ratio of YC was >2, which is consistently indicating the presence of a high burden of infection in young children⁽³⁷⁾.

Cobalt is a key component of cobalamin (vitamin B12) and is required to produce haemoglobin and red blood cells⁽³⁸⁾. It also plays important roles in the formation of amino acid and neurotransmitters⁽³⁹⁾. The median concentration of serum Co in Ethiopia was 0.4 $\mu\text{g L}^{-1}$. There was no difference among different demographic groups and their residential area. We observed that 15.9% of the study population had Co deficiency with highest prevalence in Addis Ababa and Dire Dawa administrative areas. Among demographic groups, the highest prevalence of Co deficiency was in WRA (20.7%). The ENMS reported that 15.1% of the surveyed WRA were deficient in Vitamin B12 with the highest prevalence observed in Dire Dawa (39.2%, n=1374)⁽¹³⁾. This Co deficiency in women might be associated with disturbance in Vitamin B12 synthesis, as a result it might trigger anemia⁽⁴⁰⁾.

Molybdenum is an important component of enzymes such as xanthine dehydrogenase, aldehyde oxidase and sulfite oxidase^(41,42). The median concentration of serum Mo in

Ethiopia was $0.2 \mu\text{g dL}^{-1}$ within the range of $0.01\text{--}2.1 \mu\text{g dL}^{-1}$. This is high compared with other countries like Japan, Belgium, USA, UK, Sweden, Hungary, Germany and Zaire, that serum Mo concentration is in the range of $0.12\text{--}0.22 \mu\text{g dL}^{-1}$ ⁽⁴²⁾. Similar to the distribution of Co, there was no significant difference in serum Mo status among demographic groups and their residential area. However, there were spatial differences between regions, with Mo deficiency prevalence was 1.1% in Amhara and 0.4% in the Somali region. Food groups with high Mo concentrations are plant foods such as cereals, pulses, nuts and their product ^(41,42). The national food consumption survey reported that most of the food sourced in Ethiopia are plant based ⁽³⁶⁾.

Mineral micronutrients of similar properties share same transporting and receptor proteins hence exhibit interaction at the site of absorption or during metabolic processes. For example, Ca and Mg interact and a change in Ca may affect Mg balance and vice-versa ⁽⁴³⁾. Similarly, in the present study there was an association between the mineral micronutrients. The present study has a hierarchical sampling hence the variation comprises contributions from variations among individuals within households, from households within enumeration areas and from enumeration areas. In addition, the variation varies between scales, and correlations will very likely do the same. The relative contribution of the different sources of variation to mineral concentrations are determined entirely by the sampling decisions (number of households and enumeration areas), and so these correlations do not only reflect the actual processes at the different scales, but those decisions too. As a result, these correlations can only be exploratory, and should be interpreted with caution. The significance levels are very questionable, thus are not included.

Regarding the geostatistical output, the variogram model shows spatial variability of serum Ca concentration, followed by Mg, Mo and Ca:Mg ratio. Several factors may contribute to variations in serum micronutrients concentrations. These include physiological status including presence of infection, and influences including dietary patterns, age and sex ^(44,45). In addition, in Ethiopia, spatial factors linked to soil types and landscape features appear to be strong drivers of longer-range variation in micronutrients status ^(2,14,23). Food systems are highly localized in Ethiopia, particularly in rural areas, with a large proportion of dietary intakes met through subsistence production or purchases of locally-produced food ⁽⁴⁶⁾. Thus, individuals' status reflects the soil types and landscapes where they reside. While the

variogram model represents no spatial structure for serum Cu, Cu:Zn ratio and Co concentrations (Supplementary Table S1).

The information produced in this study can be used for a baseline assessment for further studies, it can also help to target the most affected areas for appropriate food-based interventions. The information used to apply further sampling requirements may need if the kriging variance is very large as seen in eastern part of the country where the sampling data is sparse.

The main study strengths include (i) the analysis of different micronutrient concentrations in the serum of different demographic groups; (ii) the large sample size; (iii) the use of ICP-MS by which multiple elements can be reliably analyzed simultaneously and with high sensitivity and (iv) the application of geostatistical models to predict the unsampled area with the probability map and visualize the uncertainty of the prediction. However, the present study lacks detailed information on socioeconomic status, data on dietary intake, soil and crops data at national level. But, soil and crop data in Amhara regions shows a strong correlation Gashu et al. ⁽¹²⁾. These data may help us to explore true determinants of micronutrient status and deficiency in the population studied. We apply the same cut-off for all demographic groups except for Zn may increase high deficiency rate in children. In addition, serum calcium concentration is not a reliable marker of calcium status as it is subjected to strong homeostatic regulation and changes only after chronic calcium deprivation ^(47,48) hence, the calcium result of the present should be interpreted in caution.

Conclusion

This is the first national study to demonstrate the serum micronutrient concentration in Ethiopia. There is a high prevalence of Ca, Mg and Co deficiencies, with YC likely to be most affected with micronutrient deficiency. However, there is a need to establish deficiency thresholds for all demographic groups to gain a better estimation of micronutrient deficiency prevalence, as recommended by Ewers et al. ⁽⁴⁹⁾. The variogram models show that Ca, Mg, Ca:Mg ratio and Mo have spatial dependence in the distance range 140–500 km, whereas Co, Cu and Cu:Zn ratio shown spatially independent. These data contribute on the serum micronutrients status of Ethiopian population and can be helpful in the subsequent studies as baseline as well as for the cost-effective intervention program to mitigate micronutrients deficiency in high burden area.

Supplementary Materials: Supplementary information (Supplementary Information 1, Supplementary Table S1, Supplementary Figure S1) are available in the following link)

Acknowledgment

The authors would like to thank data collectors and study participants

Financial support

This work was supported, in part, by the Bill & Melinda Gates Foundation [INV-009129]. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic License has already been assigned to the Author Accepted Manuscript version that might arise from this submission. The funder had no role in the design, execution, analyses or interpretation of the data.

Conflict of Interest

The authors declare no conflicts of interest.

Authorship

A.B., D.G., E.J.M.J., R.M.L. and M.R.B. designed the research; A.B. and D.Z. supervised data collection; A.B., S.D.Y., E.H.B. conducted the laboratory analysis; A.B., E.L.A., and R.M.L. performed statistical analysis; R.M.L. and C.C. prepared Figs. 1, 2, 3, 4, and 5; All authors reviewed the manuscript.

References

1. Al-fartusie FS & Mohssan SN (2017) Essential trace elements and their vital roles in human body. *Indian J Adv Chem Sci* **5**, 127–36.
2. Joy EJM, Ander EL, Young SD, *et al.* (2014) Dietary mineral supplies in Africa. *Physiol Plant* **151**, 208–29.
3. Gombart AF, Pierre A, & Maggini S (2020) A review of micronutrients and the immune system—working in harmony to reduce the risk of infection. *Nutrients* **12**, 1–36.
4. Kumssa DB, Joy EJM, Ander EL, *et al.* (2015) Dietary calcium and zinc deficiency risks are decreasing but remain prevalent. *Sci Rep* **5**, 1–11.
5. Amare B, Moges B, Fantahun B *et al.* (2012) Micronutrient levels and nutritional status of school children living in northwest Ethiopia. *Nutr J* **11**, 1–8.
6. Belay A, Gashu D, Joy EJM, *et al.* (2021) Zinc deficiency is highly prevalent and spatially dependent over short distances in Ethiopia. *Sci Rep* **11**, 6510.
<https://doi.org/10.1038/s41598-021-85977-x>
7. Sandström B (2001) Micronutrient interactions: effects on absorption and bioavailability. *Br J Nutr*, **85**:S181.
8. Arthur JR, Beckett GJ, & Mitchell JH (1999) The interactions between selenium and iodine deficiencies in man and animals. *Nutr Res Rev*, **12**:55–73.
9. Dai Q, Shu X, Deng X, *et al.* (2013) Modifying effect of calcium / magnesium intake ratio and mortality : a population- based cohort study. *BMG open*, **3**:1–7.
10. Osredkar J & Sustar N (2011) Copper and zinc, biological role and significance of copper/zinc imbalance. *J Clin Toxicol* **S3**, 1–18.
11. Belay A, Joy EJM, Chagumaira C, *et al.* (2020) Selenium deficiency is widespread and spatially dependent in Ethiopia. *Nutrients* **12**, 1–17.
12. Gashu D, Lark RM, Milne AE, *et al.* (2020) Spatial prediction of the concentration of selenium (Se) in grain across part of Amhara Region, Ethiopia. *Sci Total Environ.* **733**:139231.

13. UNICEF; World Bank; World Food Program; USAID ENGIN (2016) Ethiopian National Micronutrient Survey Report. Available online: https://www.ephi.gov.et/images/pictures/download2009/National_MNS_report.pdf.
14. Gashu D, Marquis GS, Bougma K, *et al.* (2018) Spatial variation of human selenium in Ethiopia. *Biol Trace Elem Res* **189**, 354–60.
15. Sauberlich, H.E. (1999) Laboratory Tests for the Assessment of Nutritional Status (2nd ed.) Routledge, Boca Raton, FL: CRC Press. <https://doi.org/10.1201/9780203749647>.
16. Laine JT, Pekka T, Jukka T, *et al.* (2020) Serum copper-to-zinc-ratio and risk of incident infection in men : the kuopio ischaemic heart disease risk factor study. *Eur J Epidemiol* **35**, 1149–56.
17. Emokpae MA, & Fatimehin EB (2020) Copper-to-zinc ratio as an inflammatory marker in patients with sickle cell disease. *Sci* **2**, 1–9.
18. Tessema M, Groote H De, Brouwer ID, *et al.* (2019) Soil zinc is associated with serum zinc but not with linear growth of children in Ethiopia. *Nutrients* **11**, 221.
19. Population Census Commission (2008) Summary and Statistical Report of the 2007 Population and Housing Census. Population Size by Age and Sex; Population Census Commission: Addis Ababa, Ethiopia.
(file:///C:/Users/afomt/Downloads/2007%20Census%20Admin%20Report%20(1).pdf)
20. WHO (2010) WHO Guidelines on Drawing Blood: Best Practices in Phlebotomy. Geneva, Switzerland.
21. Tukey JW (1977) Exploratory Data Analysis. Addison-Wesley: Reading, MA, USA.
22. King JC, Brown KH, Gibson RS, *et al.* (2016) Biomarkers of nutrition for development (BOND)—Zinc Review. *J Nutr* **146**, 858S–85S.
23. Phiri FP, Ander EL, Elizabeth HB, *et al.* (2019) The risk of selenium deficiency in Malawi is large and varies over multiple spatial scales. *Sci Rep* **9**, 6566.
24. Lark RM (2000) A comparison of some robust estimators of the variogram for use in soil survey. *Eur J Soil Sci* **51**, 137 – 157.

25. Mastrandrea MD, Field CB, Stocker TF *et al.* (2010) Guidance Note for Lead Authors of the IPCC Fifth Assessment Report on Consistent Treatment of Uncertainties; Intergovernmental Panel on Climate Change (IPCC): Geneva, Switzerland. Available online:
https://www.ipcc.ch/site/assets/uploads/2017/08/AR5_Uncertainty_Guidance_Note.
26. Lark RM, Ander EL, Cave MR *et al.* (2014) Mapping trace element deficiency by cokriging from regional geochemical soil data : A case study on cobalt for grazing sheep in Ireland. *Geoderma* **226**, 64–78.
27. Tesfaye B, Sinclair K, Wuehler SE *et al.* (2018) Applying international guidelines for calcium supplementation to prevent pre-eclampsia : simulation of recommended dosages suggests risk of excess intake in Ethiopia. *Public Health Nutr* **22**, 531-541.
28. United Nations Administrative Committee on Coordination, Sub-committee on Nutrition (1997) 3rd Report on the World Nutrition Situation. Geneva: ACC/SCN Secretariat; available at https://www.unscn.org/web/archives_resources/files/rwns3.pdf.
29. Evang EC, Habte TY, Owino WO *et al.* (2020) The nutritional and micronutrient status of urban schoolchildren with moderate anemia is better than in a rural area in Kenya. *Nutrients* **12**, 207.
30. Paper O (2016) Higher levels of magnesium and lower levels of calcium in whole blood are positively correlated with the metabolic syndrome in a Chinese population : a case–control study. *Ann Nutr Metab* **69**, 125–34.
31. Resnick L (1999) The cellular ionic basis of hypertension and allied clinical conditions. *Prog Cardiovasc Dis* **42**, 1–22.
32. Resnick LM, Gupta RK, Bhargava KK *et al.* (1991) Cellular ions in hypertension, diabetes, and obesity. *Hypertension* **17**, 951–7.
33. Okoror CEM, Enabudoso EJ, Okoror OT *et al.* (2020) Serum calcium-magnesium ratio in women with pre-eclampsia at a tertiary hospital in Nigeria. *Int J Gynecol Obstet* **149**, 354–8.

34. Sahmoun AE & Singh BB (2010) Does a higher ratio of serum calcium to magnesium increase the risk for postmenopausal breast cancer? *Med Hypotheses* **75**, 315–8.
35. Sato H, Takeuchi Y, & Matsuda K (2018) Evaluation of the predictive value of the serum calcium-magnesium ratio for all-cause and cardiovascular mortality in incident dialysis patients. *Cardiorenal Med* **8522**, 50–60.
36. Hafebo AS, Wuehler S, Gibbs M *et al.* (2015) Overview of the Ethiopian national food consumption survey (NFCS): Implications for fortification programs. *Eur J Nutr Food Saf* **5**, 961–962.
37. Lu HD, Wang ZQ, Pan YR *et al.* (1999) Comparison of serum Zn, Cu and Se contents between healthy people and patients in high, middle and low incidence areas of gastric cancer of Fujian Province. *World J Gastroenterol* **5** 84-86.
38. Barceloux DG & Barceloux D (1999). Cobalt. *Journal of Toxicology: Clinical Toxicology* **37**, 201–216. doi:10.1081/clt-100102420.
39. Prashanth L, Kattapagari KK, Chitturi RT, *et al.* (2015) A review on role of essential trace elements in health and disease. *J Ntr Univ Heal Sci* **4**, 75–85.
40. Battaglia V, Compagnone A, Bandino A *et al.* (2009) Cobalt induces oxidative stress in isolated liver mitochondria responsible for permeability transition and intrinsic apoptosis in hepatocyte primary cultures. *Int J Biochem Cell Biol* **41**, 586–94.
41. Rajagopalan K.V (1988) MOLYBDENUM : An Essential Trace Element in Human Nutrition. *Ann Rev Nutr*, **8**:401–27.
42. Yoshida M (2012) Molybdenum in biological samples and clinical significance of serum molybdenum. In: Za'ray SC and G, editor. First edit, New Jersey: John Wiley & Sons, Inc. p. 397–408.
43. Rosanoff A, Dai Q, & Shapses SA (2016) Essential nutrient interactions: Does low or suboptimal magnesium status interact with vitamin D and/or calcium status? *Adv Nutr*, **7**:25–43.
44. Bui VQ, Stein AD, Digirolamo AM *et al.* (2012) Associations between serum c-reactive protein and serum zinc , ferritin , and copper in guatemalan school children. *Biol Trace Elem Res* **148**, 154–60.

45. Paz-tal O, Can A, Marko R *et al.* (2013) Dynamics of magnesium, copper, selenium and zinc serum concentrations for 2-year dietary intervention. *ESPEN J* **8**, e100–7.
46. Central Statistical Agency of Ethiopia (2012) Household Consumption and Expenditure Survey 2010/11: Analytical Report. Statistical Bulletin 563; Central Statistical Agency of Ethiopia: Addis Ababa, Ethiopia.
47. McLean FC & Hastings AB (1935) The state of calcium in the fluids of the body. *J Biol Chem* **108**, 285–322.
48. Gibson RS (2005) Principles of nNutritional Assessment, 2nd ed., pp. 645. New York: Oxford University Press.
49. Ewers U, Krause C, Schulz C *et al.* (1999) Reference values and human biological monitoring values for environmental toxins. Report on the work and recommendations of the commission on human biological monitoring of the German federal environmental agency. *Int Arch Occup Environ Health* **72**, 255–60.

Table 1. Median [Q1, Q3] serum concentrations of micronutrients among the Ethiopian population, according to region, demographic group, location of residence, and educational status.

Variable	n	Ca, mg dL ⁻¹ †	Mg, mg dL ⁻¹ †	Ca:Mg ratio†	Co, µg L ⁻¹ †	Cu, µg dL ⁻¹ †	Zn, µg dL ⁻¹ †	Cu:Zn ratio†	Mo, µg dL ⁻¹ †
Region									
Addis Ababa	184	8.5[8.2,8.8]	2.0[1.9,2.1]	4.3[4.0,4.5]	0.26[0.18,0.41]	110.5[94.7,1,24.4]	67.9[57.9,72.5]	1.8[1.5,1.9]	0.17[0.12,0.25]
Afar	266	8.5[7.9,8.9]	1.9[1.8,2.0]	4.5[4.2,4.6]	0.36[0.22,0.54]	102.4[89.9,118.6]	58.6[50.7,65.9]	1.8[1.5,2.2]	0.19[0.14,0.27]
Amhara	480	9.0[8.5,9.3]	2.0[1.9,2.2]	4.4[4.1,4.7]	0.52[0.36,0.75]	101.4[90.3,113.2]	57.4[49.9,64.4]	1.8[1.5,2.1]	0.11[0.07,0.17]
Benishangul-Gumuz	211	8.3[7.9,8.7]	1.9[1.8,2.0]	4.4[4.1,4.7]	0.59[0.38,0.95]	96.6[83.5,112.2]	57.6[49.9,65.1]	1.7[1.4,2.0]	0.14[0.10,0.21]
Dire Dawa	156	9.2[8.8,9.6]	2.0[1.9,2.2]	4.5[4.2,4.7]	0.39[0.18,0.60]	99.2[87.7,118.7]	60.8[55.4,67.4]	1.7[1.4,1.9]	0.19[0.14,0.26]
Gambela	192	8.4[7.9,8.9]	1.9[1.8,2.1]	4.3[4.1,4.5]	0.42[0.29,0.68]	115.2[99.4,128.3]	58.3[52.1,65.5]	1.9[1.6,2.3]	0.20[0.14,0.30]
Harari	167	8.8[7.8,9.2]	1.9[1.8,2.1]	4.5[4.2,4.7]	0.39[0.24,0.64]	98.1[84.1,116.6]	55.7[46.6,63.7]	1.9[1.6,2.2]	0.19[0.15,0.28]
Oromia	515	8.4[7.8,8.8]	1.9[1.7,2.0]	4.4[4.2,4.7]	0.45[0.28,0.81]	96.9[85.2,109.5]	54.5[46.3,62.2]	1.8[1.5,2.2]	0.15[0.10,0.23]

SNNP		8.4[7.9,8.8	1.9[1.7,2.0	4.4[4.2,4.7	0.32[0.22,0.56	96.2[84.6,110.1]	57.3[49.4,63.5	1.7[1.4,2.1	0.23[0.16,0.34
	390]]]]]]]
Somali		8.7[8.3,9.1	1.9[1.8,2.1	4.5[4.2,4.8	0.47[0.22,0.78	110.6[99.2,127.7]	59.7[53.2,67.1	1.9[1.6,2.2	0.16[0.12,0.22
	201]]]]]]]
Tigray		8.3[7.7,8.6	1.9[1.8,2.0	4.3[4.1,4.5	0.42[0.28,0.63	97.4[86.6,111.9]	54.0[46.7,60.8	1.9[1.5,2.2	0.18[0.13,0.25
	340]]]]]]]
Total	3,10	8.6[8.0,9.0	1.9[1.8,2.1	4.4[4.2,4.7	0.43[0.27,0.73	98.8[87.3,112.7]	56.2[48.6,63.5	1.8[1.5,2.1	0.16[0.10,0.24
	2]]]]]]]
Demographic group									
YC		7.3[6.9,7.7	1.7[1.6,1.8	4.4[4.1,4.6	0.43[0.30,0.69	101.8[89.1,116.1]	46.1[40.4,53.3	2.2[1.9,2.6	0.15[0.11,0.22
	474]]]]]]]
SAC		8.6[8.1,9.9	1.9[1.8,2.1	4.4[4.2,4.6	0.45[0.31,0.71	100.9[89.5,113.0]	55.9[48.9,61.9	1.8[1.6,2.1	0.16[0.11,0.26
	935]]]]]]]
Men		8.8[8.4,9.1	1.9[1.8,2.0	4.5[4.3,4.8	0.39[0.25,0.67	91.7[83.2,100.4]	61.1[52.6,67.3	1.5[1.3,1.8	0.15[0.10,0.22
	403]]]]]]]
WRA	1,29	8.7[8.3,9.2	1.9[1.8,2.1	4.5[4.2,4.7	0.43[0.25,0.76	99.7[87.3,113.3]	59.6[51.4,65.7	1.7[1.4,2.0	0.16[0.10,0.24
	0]]]]]]]
Residence									
Rural	2,28	8.5[7.9,9.0	1.9[1.8,2.1	4.4[4.2,4.7	0.45[0.29,0.76	98.7[86.9,111.9]	55.4[47.6,62.5	1.8[1.5,2.2	0.16[0.10,0.24
	9]]]]]]]

Urban	813	8.7[8.3,9.1]	2.0[1.8,2.1]	4.4[4.2,4.7]	0.35[0.22,0.55]	101.1[88.8,115.6]	61.7[54.9,67.5]	1.7[1.4,2.0]	0.15[0.10,0.22]
Educational status									
Literate	1,548	8.5[8.1,9.0]	1.9[1.8,2.1]	4.4[4.2,4.7]	0.41[0.26,0.69]	98.4[86.6,112.6]	57.6[49.6,64.7]	1.7[1.5,2.1]	0.16[0.10,0.24]
Illiterate	1,554	8.6[7.9,9.0]	1.9[1.8,2.1]	4.4[4.2,4.7]	0.47[0.29,0.75]	99.5[87.6,112.8]	55.4[47.7,62.3]	1.8[1.5,2.2]	0.16[0.10,0.24]

Note: SNNP: Southern Nations, Nationalities, and Peoples; YC: young children; SAC: School Age Children; WRA: Women of reproductive age; † values in brackets are Q1: 25th percentile and Q3: 75th percentile.

Table 2. Micronutrient deficiency prevalence (%) among the Ethiopian population, according to region, demographic group, location of residence, and educational status.

Variable	n	Ca	Mg	Co	Cu	Zn	Mo
Region							
Addis Ababa	184	39.1	15.1	36.4	5.3	27.2	0.00
Afar	266	40.3	27.7	23.5	8.9	63.2	0.00
Amhara	480	21.1	17.1	6.5	2.9	74.1	1.14
Benishangul-Gumuz	211	54.7	28.7	9.4	12.9	69.9	0.00
Dire Dawa	156	9.2	5.5	33.2	7.5	66.4	0.00
Gambela	192	49.5	20.3	11.5	2.7	64.4	0.00
Harari	167	37.5	27.6	20.5	10.6	71.8	0.00
Oromia	515	51.9	37.2	16.0	8.5	80.0	0.00
SNNP	390	46.7	33.6	25.8	11.5	73.2	0.00
Somali	201	30.4	23.3	24.4	2.6	66.5	0.43
Tigray	340	59.5	29.5	12.5	5.8	78.4	0.00
Total	3,102	41.6	29.2	15.9	7.6	75.1	0.33
Demographic group							
YC	474	90.5	76.1	12.1	7.9	92.1	0.00
SAC	935	38.2	20.9	11.2	7.6	75.8	0.95
Men	403	23.1	23.4	17.7	7.7	73.9	0.00
WRA	1,290	32.2	20.5	20.7	6.3	67.2	0.04

Residence							
Rural	2,289	42.9	30.7	14.2	7.6	78.0	0.43
Urban	813	33.9	20.4	25.5	4.5	58.1	0.00
Household head educational status							
Educated	1,548	40.5	29.5	18.7	7.2	72.5	0.00
Non educated	1,554	42.6	28.9	13.6	7.1	77.3	0.61

Note: SNNP: Southern Nations, Nationalities, and Peoples; YC, young children; SAC, School Age Children and WRA, Women of reproductive age

Table 3. Pearson's correlation coefficients between serum micronutrients of the Ethiopian population

Micronutrient	Mg	Ca	Co	Cu	Zn	Mo
Mg	1.00					
Ca	0.70	1.00				
Co	0.00	0.01	1.00			
Cu	0.26	0.21	0.02	1.00		
Zn	0.31	0.42	0.03	0.12	1.00	
Mo	0.04	0.00	0.01	0.01	0.04	1.00




Figure 1. Locations of the centroids of $n=346$ Enumeration Areas from which study participants were recruited ^(6,11).




Figure 2. Predicted serum (a) Ca, (b) Mg, (c) Ca:Mg ratio and (d) Mo concentration (the mean of the prediction distribution) for women of reproductive age in Ethiopia.




Figure 3. Serum micronutrient kriging variance for women of reproductive age in Ethiopia a) Ca, b) Mg, c) Ca:Mg ratio and d) Mo




Figure 4. Probability that serum micronutrient concentration of women of reproductive age would fall below threshold for serum (a) Ca, (b) Mg, (c) Ca:Mg, (d) Mo