





BMJ Open Indicators of optimal diabetes care and burden of diabetes complications in Africa: a systematic review and meta-analysis

Davis Kibirige ¹, Nyasatu Chamba,^{2,3} Irene Andia-Biraro,^{4,5} Kajiru Kilonzo,^{2,3} Sweetness Naftal Laizer,³ Isaac Sekitoleko,⁶ Andrew Peter Kyazze,⁴ Sandra Ninsiima,⁷ Phillip Ssekamatte ⁷, Felix Bongomin ⁸, Lucy Elauteri Mrema,⁹ Willyhelmina Olomi,¹⁰ Theodora D Mbunda,⁹ Nyanda Elias Ntinginya,⁹ Issa Sabi,¹¹ Katrina Sharples,¹² Philip Hill,¹² Lindsey te Brake,¹³ Josephine VandeMaat,¹⁴ Reinout vanCrevel,^{15,16} Julia Alison Critchley ¹⁷, on behalf of PROTID consortium

To cite: Kibirige D, Chamba N, Andia-Biraro I, *et al.* Indicators of optimal diabetes care and burden of diabetes complications in Africa: a systematic review and meta-analysis. *BMJ Open* 2022;**12**:e060786. doi:10.1136/bmjopen-2022-060786

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-060786>).

Received 06 January 2022
Accepted 20 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Davis Kibirige;
kibirigedavis@gmail.com

ABSTRACT

Objective Contemporary data on the attainment of optimal diabetes treatment goals and the burden of diabetes complications in adult populations with type 2 diabetes in Africa are lacking. We aimed to document the current status of attainment of three key indicators of optimal diabetes care and the prevalence of five diabetes complications in adult African populations with type 2 diabetes.

Methods We systematically searched Embase, PubMed and the Cochrane library for published studies from January 2000 to December 2020. Included studies reported any information on the proportion of attainment of optimal glycated haemoglobin (HbA1c), blood pressure (BP) and low-density lipoprotein cholesterol (LDLC) goals and/or prevalence of five diabetes complications (diabetic peripheral neuropathy, retinopathy, nephropathy, foot ulcers and peripheral arterial disease). Random effect model meta-analysis was performed to determine the pooled proportion of attainment of the three treatment goals and the prevalence of five diabetes complications.

Results In total, 109 studies with a total of 63 890 participants (53.3% being females) were included in the meta-analysis. Most of the studies were conducted in Eastern African countries (n=44, 40.4%). The pooled proportion of attainment of an optimal HbA1c, BP and LDLC goal was 27% (95% CI 24 to 30, $I^2=94.7%$), 38% (95% CI 30 to 46, $I^2=98.7%$) and 42% (95% CI 32 to 52, $I^2=97.4%$), respectively. The pooled prevalence of diabetic peripheral neuropathy, retinopathy, diabetic nephropathy, peripheral arterial disease and foot ulcers was 38% (95% CI 31 to 45, $I^2=98.2%$), 32% (95% CI 28 to 36, $I^2=98%$), 31% (95% CI 22 to 41, $I^2=99.3%$), 19% (95% CI 12 to 25, $I^2=98.1%$) and 11% (95% CI 9 to 14, $I^2=97.4%$), respectively.

Conclusion Attainment of optimal diabetes treatment goals, especially HbA1c, in adult patients with type 2 diabetes in Africa remains a challenge. Diabetes complications, especially diabetic peripheral neuropathy

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ To our knowledge, it is the first systematic review and meta-analysis to simultaneously investigate the status of attainment of the three key diabetes treatment goals and the burden of five common diabetes complications in an adult indigenous African population with type 2 diabetes.
- ⇒ The systematic review and meta-analysis included a large number of studies that assessed the extent of attainment of diabetes treatment goals and the prevalence of diabetes complications based on recommendations or definitions by internationally recognised associations.
- ⇒ There was high heterogeneity among the studies included in the meta-analysis.
- ⇒ A relative number of studies included in the meta-analysis had low to moderate quality on assessment.

and retinopathy, are highly prevalent in adult populations with type 2 diabetes in Africa.

INTRODUCTION

Globally, the burden of diabetes mellitus (DM) continues to exponentially rise to epidemic proportions, disproportionately affecting low-income and middle-income countries. The recent 2021 International Diabetes Federation (IDF) estimates show that about 24 million adults (1 in 22 adults) live with DM in Africa. The IDF also predicts that the greatest future increase in the prevalence of DM will occur in Africa because of the predicted ageing of Africa's currently very young populations, as well as increasing urbanisation and associated lifestyle changes.¹ This will ultimately lead to an immense strain




Figure 1 PRISMA flow diagram of selection of eligible studies. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

on weak healthcare systems that are poorly structured and inadequately financed to manage non-communicable diseases (NCDs) like DM.²

In addition, the rates of undiagnosed DM continue to increase in Africa. Among the IDF regions, Africa has the highest proportion of undiagnosed diabetes: about 54% of all cases.¹ The majority of patients are diagnosed late with coexisting debilitating complications, and suboptimal diabetes care remains common in most clinical settings in Africa.³ This could be explained by low awareness about DM, healthcare systems that are structured mainly to manage communicable diseases as opposed to NCD, low screening rates of DM to ensure early diagnosis, low availability of affordable essential diagnostic tests and medicines for DM and knowledge–practice gaps among healthcare practitioners.^{2 4–6}

Published diabetes treatment guidelines by most international organisations like the IDF and American Diabetes Association (ADA) recommend targets of glycated haemoglobin (HbA1c) level of <7% (53 mmol/mol), blood pressure (BP) <140/90 mm Hg and low-density lipoprotein cholesterol (LDLC) <2.6 mmol/L (100 mg/dL) as key indicators of optimal diabetes care.^{7–9} Attainment of these treatment goals in diabetes care ultimately translates to reduced risk of onset and progression of diabetes complications and mortality.

Despite the increasing burden of DM and its related complications, late diagnosis of diabetes and prevalent suboptimal diabetes care in clinical settings in Africa, there is an information gap regarding the current status of attainment of the recommended diabetes treatment goals and the prevalence of common diabetes complications to inform targeted strategies or interventions to reduce diabetes-related morbidity and mortality. This systematic review and meta-analysis aimed to document

the proportion of attainment of optimal HbA1c, BP and LDLC goals and the prevalence of five diabetes complications (diabetic peripheral neuropathy, nephropathy, retinopathy, foot ulcers and peripheral arterial disease) in adult native populations with type 2 diabetes in Africa.

METHODS

This systematic review and meta-analysis was conducted according to the criteria outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.¹⁰ The PRISMA checklist is available as an online supplemental table 1. The study protocol was registered in the PROSPERO International Prospective Register of systematic reviews (CRD42020215576).

Search strategy

We searched Embase, PubMed and the Cochrane library for published studies from January 2000 to December 2020. The following search terms were used after discussion with a medical librarian: “Quality of diabetes care” OR “Indicators of diabetes care” OR “status of diabetes care” OR “diabetes care” OR “glycaemic control” OR “blood pressure control” OR “lipid profile control” OR “screening of diabetes complications” OR “diabetes complications” OR “screening for diabetic retinopathy” OR “screening for diabetic peripheral nephropathy” OR screening for diabetic neuropathy” OR screening for diabetic foot ulcers OR “screening for peripheral arterial disease” OR “prevalence of diabetic retinopathy” OR “prevalence of diabetic peripheral nephropathy” OR “prevalence of diabetic peripheral neuropathy” OR “prevalence of diabetic foot ulcers” OR “prevalence of peripheral arterial disease”, AND “type 2 diabetes mellitus” OR “type 2 diabetes” AND Algeria OR Angola OR Benin OR Botswana OR “Burkina Faso” OR Burundi OR Cameroon OR “Cape Verde” OR “Central African Republic” OR Chad OR Comoros OR “Democratic Republic of Congo” OR Djibouti OR Egypt OR “Equatorial Guinea” OR Eritrea OR Ethiopia OR Gabon OR Gambia OR Ghana OR Guinea OR “Guinea Bissau” OR “Ivory Coast” OR “Cote d’Ivoire” OR Kenya OR Lesotho OR Liberia OR Libya OR Libya OR Madagascar OR Malawi OR Mali OR Mauritania OR Mauritius OR Morocco OR Mozambique OR Namibia OR Niger OR Nigeria OR Rwanda OR “Sao Tome” OR Senegal OR Seychelles OR “Sierra Leone” OR Somalia OR “South Africa” OR “South Sudan” OR Sudan OR Swaziland OR Tanzania OR Togo OR Tunisia OR Uganda OR Zaire OR Zambia OR Zimbabwe OR “Central Africa” OR “West Africa” OR “Western Africa” OR “East Africa” OR “Eastern Africa” OR “North Africa” OR “Northern Africa” OR “Southern Africa” OR “sub Saharan Africa” OR “sub-Saharan Africa” OR Africa.

In addition, references of included articles were hand-searched for any other original articles. The search and selection were restricted to studies written only in the English language.

Table 1 General characteristics of all participants (n=63 890) included in the systematic review and meta-analysis

Characteristic	Cumulative value	Number of studies
Age in years (mean±SD)	54.9±4.7	88
Gender – females (% , 95% CI)	55.3, 52.7 to 57.8	101
Smokers (% , 95% CI)	9.9, 0.5 to 55.6	44
Participants on OHA (% , 95% CI)	65.0, 34.0 to 96.6	51
Participants on insulin (% , 95% CI)	31.3, 26.3 to 36.2	52
Participants on lipid-lowering agents (% , 95% CI)	25.7, 0.5 to 86.7	14
Participants on anti-hypertensive agents (% , 95% CI)	73.3, 64.1 to 82.5	18
BMI in kg/m ² (mean±SD)	27.9±0.5	40
HbA1c in % (mean±SD)	9.0±1.5	40
HbA1c in mmol/mol (mean±SD)	75.0±1.5	40

BMI, body mass index; HbA1c, glycated haemoglobin; OHA, oral hypoglycaemic agents.

Study selection criteria

The preliminary screening of titles and abstracts to identify potentially eligible articles was done by two independent reviewers (NC and DK). This was followed by removing all duplicates. After the initial screening, full texts of the potentially eligible studies were retrieved and closely reviewed for eligibility.

The inclusion criteria of studies were: cross-sectional, cohort or randomised controlled trials published between January 2000 and December 2020 in English language, studies reporting any data on proportion of adult patients with type 2 diabetes who attained the recommended optimal HbA1c, BP or LDLC targets and residing in African countries and studies reporting data on any of prevalence of diabetic nephropathy, peripheral neuropathy, retinopathy, foot ulcers or peripheral arterial disease in adult patients with type 2 diabetes in African countries.

Any disagreements that arose were resolved by consensus. We excluded retrospective studies, case series and reports, studies published in languages other than English and studies whose full texts could not be retrieved.

Data extraction

After identifying the eligible original studies, they were collated and sent to additional reviewers to extract the relevant study information using a Microsoft Excel 2016 form. The information of interest that was extracted from the eligible studies included: the last name of the first author and year of publication, country(ies) and region(s) of Africa where the study was conducted, type of study design, number of study participants, the mean age of study participants, the proportion of female participants, the proportion of participants with a current or history of smoking, the proportion of participants on oral hypoglycaemic agents, insulin, lipid-lowering agents (statins) and antihypertensive agents, mean body mass index (BMI) and HbA1c of study participants, the proportions of participants with optimal HbA1c, BP and LDLC targets and the prevalence of diabetic nephropathy, peripheral

neuropathy, retinopathy, foot ulcers and peripheral arterial disease.

Operational definitions

All included studies defined optimal targets of HbA1c, BP and LDLC as <7% (53 mmol/mol), <140/90 mm Hg and <2.6 mmol/L or 100 mg/dL, respectively, as recommended by the IDF and ADA diabetes treatment guidelines.^{9 11}

The definitions and measurements of diabetes complications greatly varied between studies. The following definitions were used for each diabetes complication by the various studies: micro/macroalbuminuria and/or an estimated glomerular filtration rate <60 mL/min/1.73 m² for the presence of diabetic nephropathy, signs and symptoms suggestive of peripheral neuropathy, use of neuropathy screening scores like neuropathy disability score, Michigan Neuropathy Screening Instrument, neuropathy symptom score and 10g monofilament testing for the presence of diabetic peripheral neuropathy, presence of lesions like soft or hard exudates, cotton wool spots, microaneurysms, neovascularisation and retinal haemorrhages on funduscopy for diabetic retinopathy, presence of foot ulcers on clinical inspection for diabetic foot ulcers and the presence of measured ankle brachial index <0.9 using Doppler studies for peripheral arterial disease.

Assessment of quality of studies

The quality of all eligible studies included in the systematic review and meta-analysis was assessed using the Newcastle-Ottawa Scale (NOS).¹² This was done by two independent authors (NC and SNL). The total score of the adapted scale is eight stars. Studies with more than six stars were considered high quality, while those with 5 and 6 stars, and <5 stars were considered of moderate and low quality.

Study outcomes

The study outcomes were the pooled proportions of attainment of the recommended optimal HbA1c, BP

**Table 2** Indicators of optimal glycated haemoglobin goal

Optimal glycated haemoglobin (HbA1c) goal (n=34 studies): **pooled rate of attainment of optimal HbA1c goal=27% (95% CI 24 to 30, I²=94.7%, 95% CI 93.6 to 95.8) and I² after meta-regression: 56.3%**

Attainment of the optimal HbA1c goal per region: **central: 20% (95% CI 16 to 23), Eastern: 23% (95% CI 15 to 34), Northern: 24% (95% CI 17 to 31), Southern: 31% (95% CI 28 to 34) and Western: 37% (95% CI 29 to 46)**

First author and year	Country(ies)	Region of Africa	No. of study participants	Mean age of participants	% of females	% with optimal HbA1c
Adentunji2006 ⁶⁰	Nigeria	Western	50	–	–	52.0
Agboghoroma 2020 ⁶¹	Nigeria	Western	200	–	–	19.0
Akalu 2020 ²⁰	Ethiopia	Eastern	378	–	38.6	40.7
Amod 2012 ¹⁰¹	South Africa	Southern	701	57.4	43.9	30.4
Amour 2019 ²¹	Tanzania	Eastern	238	57.2	65.7	9.2
Ashur 2016 ⁸⁴	Libya	Northern	523	54.4	47.0	21.8
Attoye 2020 ⁶³	Nigeria	Western	260	–	–	34.6
Awadalla 2017 ⁸⁷	Sudan	Northern	424	–	49.3	15.6
Balogun 2011 ⁶⁴	Nigeria	Western	40	59.4	62.5	52.5
Bentata 2015 ⁸⁸	Morocco	Northern	637	58.5	62.3	30.1
Blum 2020 ¹¹⁷	DRC	Central	319	–	33.5	14.1
Cairncross 2017 ¹⁰⁴	South Africa	Southern	203	–	72.5	31.3
Camara 2015 ⁵⁹	Cameroon and Guinea Conakry	Central and Western	1267	58.0	61.0	26.0
Chadli 2016 ⁹⁰	Morocco	Northern	498	58.0	62.4	26.8
Chamba 2017 ²³	Tanzania	Eastern	119	58.1	49.6	39.3
Chetoui 2019 ⁹²	Morocco	Northern	1456	56.2	73.4	33.7
Cohen 2010 ¹⁰⁵	Malawi	Southern	620	52.2	60.1	36.0
Diaf 2017 ⁹³	Algeria	Northern	210	55.6	65.0	51.4
Hall 2017 ¹²⁰	Cameroon	Central	261	56.0	56.3	27.2
Iwuala 2015 ⁷¹	Nigeria	Western	100	59.9	62.0	45.0
Kibirige 2017 ³⁵	Uganda	Eastern	425	–	67.0	26.5
Kimando 2017 ³⁶	Kenya	Eastern	385	62.1	65.5	39.5
Kisozi 2017 ³⁷	Uganda	Eastern	288	48.5	38.0	23.3
Mbwete 2020 ⁴⁴	Tanzania	Eastern	161	63.9	67.1	49.7
Megallaa 2019 ⁹⁷	Egypt	Northern	180	–	24.4	4.4
Molefe-Baikai 2018 ¹¹⁰	Botswana	Southern	289	50.7	66.1	29.4
Muddu 2019 ⁴⁶	Uganda	Eastern	175	46.0	48.6	8.1
Muddu, 2016 ⁴⁵	Uganda	Eastern	202	46.0	49.5	8.4
Mwebaze 2014 ⁴⁷	Uganda	Eastern	146	53.9	48.6	19.2
Mwita 2019 ¹¹¹	Botswana	Southern	500	58.9	66.0	32.3
Noor, 2016 ⁹⁸	Sudan	Northern	387	–	49.6	15.0
Omar 2018 ⁹⁹	Sudan	Northern	339	54.8	69.9	28.1
Sobngwi 2011 ³	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	2352	53.0	61.1	29.2
Uloko 2012 ⁶⁷	Nigeria	Western	531	57.1	60.5	32.4

Table 3 Indicators of optimal blood pressure (BP) goal

Optimal BP goal (n=26 studies): **pooled rate of attainment of optimal BP goal=38% (95% CI 30 to 46, I²=98.7%–95% CI 98.6 to 99.0), and I² after meta-regression: 95.4%.**
 Attainment of the optimal BP goal per region: **Western: 31% (95% CI 20 to 43), Eastern: 40% (95% CI 24 to 57), Southern: 40% (95% CI 26 to 55), Central: 41% (95% CI 38 to 45) and Northern: 42% (95% CI 24 to 61).**

Author and year	Country(ies)	Region of Africa	No. of study participants	Mean age of participants	% of females	% with optimal BP
Abdissa <i>et al</i> 2020 ¹⁸	Ethiopia	Eastern	229	–	40.4	31.0
Agboghroma <i>et al</i> 2020 ⁶¹	Nigeria	Western	200	–	–	30.0
Akalu <i>et al</i> 2020 ²⁰	Ethiopia	Eastern	378	–	38.6	57.7
Amour <i>et al</i> 2019 ²¹	Tanzania	Eastern	238	57.2	65.7	21.7
Awadalla <i>et al</i> 2017 ⁸⁷	Sudan	Northern	424	–	49.3	60.1
Balogun <i>et al</i> 2011 ⁶⁴	Nigeria	Western	40	59.4	62.5	55.0
Chadli <i>et al</i> 2016 ⁹⁰	Morocco	Northern	498	58.0	62.4	20.2
Chahbi <i>et al</i> 2018 ⁹¹	Morocco	Northern	300	–	93.0	32.6
Chisha <i>et al</i> 2017 ²⁴	Ethiopia	Eastern	270	–	48.9	85.9
Cohen <i>et al</i> 2010 ¹⁰⁵	Malawi	Southern	620	52.2	60.1	48.0
Hall <i>et al</i> 2017 ^{5 120}	Cameroon	Central	261	56.0	56.3	43.0
Hayfron-Benjamin <i>et al</i> 2019 ⁷⁰	Ghana	Western	206	52.9	68.9	37.9
Jingi <i>et al</i> 2015 ¹²¹	Cameroon	Central	407	54.2	41.8	40.4
Kahloun <i>et al</i> 2014 ⁹⁶	Tunisia	Northern	2320	54.5	60.2	62.5
Kimando <i>et al</i> 2017 ³⁶	Kenya	Eastern	385	62.1	65.5	50.4
Lewis <i>et al</i> 2018 ¹⁰⁷	Zambia	Southern	921	56.0	45.0	46.6
Lumu <i>et al</i> 2017 ³⁹	Uganda	Eastern	425	52.2	67.0	54.7
Magan <i>et al</i> 2019 ⁴¹	Uganda	Eastern	44	50.4	63.4	34.1
Megallaa <i>et al</i> 2019 ⁹⁷	Egypt	Northern	180	–	24.4	37.8
Muddu <i>et al</i> 2016 ⁴⁵	Uganda	Eastern	202	46.0	49.5	38.1
Mwebaze <i>et al</i> 2014 ⁴⁷	Uganda	Eastern	146	53.9	48.6	1.5
Mwita <i>et al</i> 2019 ¹¹¹	Botswana	Southern	500	58.9	66.0	54.2
Onakpoya <i>et al</i> 2015 ⁷⁷	Nigeria	Western	133	–	48.1	24.1
Rotchford <i>et al</i> 2002 ¹¹³	South Africa	Southern	253	56.5	73.1	14.0
Sobngwi <i>et al</i> 2011 ³	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	2352	53.0	61.1	21.0
Uloko <i>et al</i> 2012 ⁶⁷	Nigeria	Western	531	57.1	60.5	17.0

and LDLC goals and the pooled prevalence of diabetic nephropathy, peripheral neuropathy, retinopathy, foot ulcers and peripheral arterial disease in adult patients with type 2 diabetes in Africa.

Data analysis

All analyses were performed using STATA V.16.0 statistical software (Stata Corp, USA). The descriptive data of all eligible studies included in the systematic review and meta-analysis like age, gender, the proportion of participants on specific glucose-lowering agents, BMI and

HbA1c were summarised using frequencies and 95% CIs and mean±SD.

For the continuous variables, the average estimated value was obtained from each of the studies, and this was used in the final analysis, while for the categorical variables, the proportions were estimated for each of the studies and used in the final analysis.

The pooled proportions of achievement of optimal HbA1c, BP and LDLC goals and the prevalence of diabetic nephropathy, peripheral neuropathy, retinopathy, foot ulcers and peripheral arterial disease were determined using a random effect model meta-analysis and presented




Figure 2 Forest plot summarising studies on the proportion of attainment of an optimal low-density lipoprotein cholesterol goal in percentage. ES, effect size.

in forest plots. The DerSimonian and Laird method was used for pooling random effects estimates.¹³

The heterogeneity of studies was assessed using the I^2 value and corresponding 95% CIs. Based on the Cochrane collaboration guide, the I^2 values of 0%–40%, 30%–60%, 50%–90% and 75%–100% were considered not important, moderate, substantial and considerable levels of heterogeneity, respectively.¹⁴ To further explore heterogeneity effects across studies, we conducted a meta-regression analysis to assess whether the heterogeneity could be explained by the study level characteristics, that is, age, sex of participants and region, in which the study was conducted. The age, BMI and sex of the participants was defined as the estimated mean age and BMI of participants and the proportion of females from each of the




Figure 3 Forest plot summarising studies on the proportion of attainment of an optimal blood pressure goal in percentage. ES, effect size.

study, respectively. The region of the study was defined as the area (Northern, Southern, Eastern, Western, and Central Africa) where the study was conducted. One effect measure per study was considered in the meta-regression. All the variables were included in the model together to assess for variability.

We assessed the presence of publication bias using the Egger test of bias with $p < 0.05$ indicating significant publication bias.¹⁵ A narrative review was also used to present the study results. Information about all included studies was also summarised in tables.

We also performed a sensitivity analysis based on the NOS scores of the studies (excluding moderate and low-quality studies) and compared the analysis with all the eligible studies and with only high-quality studies to identify any differences in the pooled estimates of the rates of attainment of optimal diabetes treatment goals and the prevalence of the five diabetes complications.

Patient and public involvement

The main research question and outcomes of interest of the systematic review and meta-analysis were informed by the need to understand the burden of diabetes complications in patients with type 2 diabetes in Africa and the extent of attainment of optimal diabetes care to inform strategies aimed to improve optimal management of diabetes in the region. Because it was a systematic review and meta-analysis, we did not involve patients in its design, recruitment and conduct.

RESULTS

Figure 1 summarises the article selection in a PRISMA flow diagram.

The literature search returned a total of 835 articles. From these, 222 duplicates were removed. Titles and abstracts of the remaining 613 articles were reviewed, and 235 articles were identified for full-text retrieval. Of the 235 articles, 126 were excluded, and the remaining 109 articles were included in this systematic review and meta-analysis. A total of 48 and 89 eligible studies contained information on optimal diabetes treatment goals and diabetes complications, respectively, while 28 studies reported information on both.

The 126 excluded articles included five studies published in French language, 21 retrospective studies, six studies with general populations (not entirely patients with type 2DM), 18 studies whose full texts were unable to be retrieved and 76 studies that did not report outcomes of interest.

Characteristics of included studies

The majority of studies were performed in Eastern African countries (44, 40.4%).^{3 16–58} The proportion of studies conducted in Western, Northern, Southern and Central Africa was 22% ($n=24$ studies),^{3 59–80} 16.5% ($n=18$ studies),^{81–99} 15.6% ($n=17$ studies)^{100–116} and 8.3% ($n=9$ studies),^{3 59 117–123} respectively. Three studies were

conducted in more than one region of Africa (Western, Central and Eastern).^{3 58 59} Most of the studies were cross-sectional in design (100, 91.7%).

Considerable heterogeneity was noted across the studies with the I^2 value ranging from 97.4% to 99.3% for studies reporting the burden of diabetes complications and 94.7%–98.7% for studies reporting the extent of attainment of optimal diabetes treatment goals. However, on meta-regression after adjusting for age and sex of study participants, and region where each study was conducted, the heterogeneity based on I^2 of studies on the prevalence of diabetes complications decreased, ranging from 1.4% for studies on diabetic foot ulcers to 95.6% for studies on diabetic nephropathy. For studies on the proportion of attainment of optimal treatment goals, the heterogeneity also decreased, to 56.3%, 92.1% and 95.4%, for studies reporting optimal HbA1c, LDLC and BP goals.

Characteristics of study participants

Table 1 summarises the characteristics of all participants in the studies included in the systematic review and meta-analysis.

The studies had a total of 63 890 participants (ranging from 40 to 11 866) with 53.3% being female. The mean±SD age, BMI and HbA1c of the participants was 54.9±4.7 years (ranging from 40.5 to 63.9 years), 27.9±0.5 kg/m² (ranging from 20.6 to 42.9 kg/m²) and 9.0±1.5% (ranging from 6.5% to 13.9%), respectively. Among the studies that reported data on the type of glucose-lowering therapies used by participants, treatment with oral hypoglycaemic agents, insulin, statins and antihypertensives was reported in about 65% (95% CI 34 to 96.6), 31.3% (95% CI 26.3 to 36.2), 25.7% (95% CI 0.5




Figure 4 Forest plot summarising studies on the proportion of attainment of an optimal glycated haemoglobin goal in percentage. ES, effect size.

to 86.7) and 73.3% (95% CI 64.1 to 82.5) of participants, respectively.

Assessment of study quality and publication bias

The assessment of the quality of studies and funnel plots assessing publication bias are summarised in online

Table 4 Indicators of optimal LDLC goal

Optimal LDLC goal (n=11 studies)
Pooled rate of attainment of optimal LDLC goal=42% (95% CI 32 to 52, $I^2=97.4\%$ –95% CI 96.5 to 98.1) and I^2 after meta-regression-92.1%).
 Attainment of the optimal LDLC goal per region: **Southern: 27% (95% CI 24 to 30), Eastern: 37% (95% CI 30 to 45), Western: 51% (95% CI 43 to 58) and Northern: 53% (95% CI 32 to 74).**

Author and year	Country(ies)	Region of Africa	No. of study participants	Mean age of participants	% of females	% with optimal LDLC
Agboghroma et al 2020 ⁶¹	Nigeria	Western	200	–	–	50.5
Amour et al 2019 ²¹	Tanzania	Eastern	238	57.2	65.7	26.0
Awadalla et al 2017 ⁸⁷	Sudan	Northern	424	–	49.3	47.4
Chadli et al 2016 ⁹⁰	Morocco	Northern	498	58.0	62.4	38.6
Chamba et al 2017 ²³	Tanzania	Eastern	119	58.1	49.6	27.7
Elnasri et al 2008 ⁹⁴	Sudan	Northern	250	52.0	62.0	84.8
Kisozi et al 2017 ³⁷	Uganda	Eastern	288	48.5	38.0	37.0
Lumu et al 2017 ³⁹	Uganda	Eastern	425	52.2	67.0	38.9
Megallaa et al 2019 ⁹⁷	Egypt	Northern	180	–	24.4	37.8
Mwebaze et al 2014 ⁴⁷	Uganda	Eastern	146	53.9	48.6	48.6
Mwita et al 2019 ¹¹¹	Botswana	Southern	500	58.9	66.0	20.4

LDLC, low-density lipoprotein cholesterol.

**Table 5** Prevalence of diabetic nephropathy

Prevalence of diabetic nephropathy (n=40 studies): **pooled prevalence=31% (95% CI 22 to 41, I²=99.3% 95% CI 99.2 to 99.4) and I² after meta-regression: 95.6%**.
 Prevalence of diabetic nephropathy per region: **Central: 22% (95% CI 9 to 39), Eastern: 25% (95% CI 10 to 43), Southern: 28% (95% CI 18 to 40), Northern: 38% (95% CI 14 to 65) and Western: 47% (95% CI 25 to 69)**.

Author and year	No. of study participants	Country (ies)	Region of Africa	Mean age of participants	% of females	Prevalence of nephropathy, %
Abejew <i>et al</i> 2015 ¹⁹	216	Ethiopia	Eastern	45.0	42.6	2.2
Adeniyi <i>et al</i> 2020 ¹⁰⁰	327	South Africa	Southern	–	70.3	24.5
Adentunji <i>et al</i> 2006 ⁶⁰	50	Nigeria	Western	–	–	83.0
Ahmed <i>et al</i> 2017 ⁸²	316	Sudan	Northern	58.0	41.5	40.2
Albalawi <i>et al</i> 2020 ⁸³	159	Sudan	Northern	58.1	65.4	26.4
Alebiosu <i>et al</i> 2013 ⁶²	342	Nigeria	Western	53.4	–	28.4
Amour <i>et al</i> 2019 ²¹	315	Tanzania	Eastern	57.2	65.7	72.2
Balogun <i>et al</i> 2011 ⁶⁴	40	Nigeria	Western	59.4	62.5	90.0
Bello <i>et al</i> 2017 ⁶⁶	358	Nigeria	Western	57.8	61.7	53.4
Bentata <i>et al</i> 2015 ⁸⁸	637	Morocco	Northern	58.5	62.3	77.2
Blum <i>et al</i> 2020 ¹¹⁷	319	DRC	Central	–	33.5	38.6
Bouaziz <i>et al</i> 2012 ⁸⁹	73	Tunisia	Northern	59.3	–	11.0
Chahbi <i>et al</i> 2018 ⁹¹	300	Morocco	Northern	–	93.0	26.3
Cohen <i>et al</i> 2010 ¹⁰⁵	620	Malawi	Southern	52.2	60.1	34.7
Deribe <i>et al</i> 2014 ²⁷	216	Ethiopia	Eastern	50.7	40.3	8.8
Dzudie <i>et al</i> 2012 ¹¹⁸	420	Cameroon	Central	56.7	51.0	15.9
Efundem <i>et al</i> 2017 ¹¹⁹	162	Cameroon	Central	55.3	67.3	14.2
Eghan <i>et al</i> 2007 ⁶⁹	109	Ghana	Western	54.1	75.0	43.0
Fasil <i>et al</i> 2019 ²⁸	367	Ethiopia	Eastern	48.6	59.3	4.4
Gill <i>et al</i> 2008 ³⁰	105	Ethiopia	Eastern	41.0	30.0	51.0
Goro <i>et al</i> 2019 ³¹	208	Ethiopia	Eastern	54.8	47.1	26.0
Hayfron-Benjamin <i>et al</i> 2019 ⁷⁰	206	Ghana	Western	52.9	68.9	32.0
Janmohamed <i>et al</i> 2013 ³²	369	Tanzania	Eastern	54.0	53.4	83.7
Kahloun <i>et al</i> 2014 ⁹⁶	2320	Tunisia	Northern	–	60.2	3.4
Khalil <i>et al</i> 2019 ⁸⁶	506	Egypt	Northern	–	–	33.2
Lebeta <i>et al</i> 2017 ³⁸	344	Ethiopia	Eastern	40.5	42.7	11.4
Machingura <i>et al</i> 2017 ¹⁰⁸	260	Zimbabwe	Southern	57.6	72.7	45.4
Makwero <i>et al</i> 2018 ¹⁰⁹	150	Lesotho	Southern	58.2	80.7	6.7
Megallaa <i>et al</i> 2019 ⁹⁷	180	Egypt	Northern	–	24.4	86.1
Mohmad <i>et al</i> 2011 ⁸¹	71	Sudan	Central	–	42.0	50.7
Molefe-Baikai <i>et al</i> 2018 ¹¹⁰	289	Botswana	Southern	50.7	66.1	44.6
Muddu <i>et al</i> 2019 ⁴⁶	175	Uganda	Eastern	46.0	48.6	47.4
Neuhann <i>et al</i> 2001 ⁴⁸	474	Tanzania	Eastern	53.8	46.0	7.5
Olamoyegun <i>et al</i> 2015 ⁷⁶	90	Nigeria	Western	62.5	50.0	54.3
Rotchford <i>et al</i> 2002 ¹¹³	253	South Africa	Southern	56.5	73.1	46.4
Sobngwi <i>et al</i> 2011 ³	2352	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	53.0	61.1	2.4
Tesfaye <i>et al</i> 2015 ⁵³	247	Ethiopia	Eastern	–	40.5	6.5

Continued

Table 5 Continued

Prevalence of diabetic nephropathy (n=40 studies): **pooled prevalence=31% (95% CI 22 to 41, I²=99.3% 95% CI 99.2 to 99.4) and I² after meta-regression: 95.6%**.

Prevalence of diabetic nephropathy per region: **Central: 22% (95% CI 9 to 39), Eastern: 25% (95% CI 10 to 43), Southern: 28% (95% CI 18 to 40), Northern: 38% (95% CI 14 to 65) and Western: 47% (95% CI 25 to 69).**

Author and year	No. of study participants	Country (ies)	Region of Africa	Mean age of participants	% of females	Prevalence of nephropathy, %
Thinyane <i>et al</i> 2013 ¹¹⁴	80	Lesotho	Southern	49.0	49.0	6.0
Uloko <i>et al</i> 2012 ⁶⁷	531	Nigeria	Western	57.1	60.5	3.2
Worku <i>et al</i> 2010 ⁵⁷	305	Ethiopia	Eastern	44.4	37.1	15.7

supplemental table 2 and online supplemental figures 1–8, respectively.

Based on the NOS, 84 (77.1%) of the included studies were of high quality, with 17 (15.6%) studies and 8 (7.3%) studies being of moderate and low quality, respectively.

Regarding the assessment of publication bias, there was observed publication bias, especially in studies about the prevalence of diabetic nephropathy, peripheral neuropathy and attainment of optimal BP control. The proportion of studies investigating the prevalence of diabetic nephropathy, peripheral neuropathy, peripheral arterial disease, retinopathy and foot ulcers located within the funnel plot was 30% (n=12), 46.1% (n=13), 55.6% (n=10), 57% (29) and 90% (n=26), respectively. About 46%, 65% and 73% of studies that reported the proportion of attainment of optimal BP, HbA1c and LDLC treatment goal were located within the funnel plot, respectively.

Extent of attainment of optimal HbA1c, BP and LDLC goals

Data on the reported proportions achieving the three diabetes treatment goals are summarised in tables 2–4 and as forest plots in figures 2–4.

Data on attainment of optimal HbA1c, BP and LDLC goals were reported in 34 studies,^{3 20 21 23 35–37 44–47 59–61 63 64 67 84 87 90 92 93 97–99 104 105 111 116 117 120 124} 26 studies,^{3 18 20 21 24 36 40 41 45 47 61 64 67 70 77 87 90 91 96 97 105 107 111 113 120 121} and 11 studies,^{21 37 39 47 61 87 90 94 97 111 116} respectively. The pooled proportion of attainment of an optimal HbA1c, BP and LDLC goal in the respective studies was 27% (95% CI 24 to 30, I²=94.7%), 38% (95% CI 30 to 46, I²=98.7%) and 42% (95% CI 32 to 52, I²=97.4%), respectively.

The lowest proportion of attainment of optimal HbA1c was reported in a study performed in Egypt (4.4%)⁹⁷ and the highest in a study performed in Nigeria (52.5%).⁶⁴ Among studies reporting the extent of attainment of an optimal BP goal, the proportion ranged from 1.5% in a study performed in Uganda⁴⁷ to 85.9% in a study performed in Ethiopia.²⁴ Among the studies reporting information on the optimal LDLC goal, attainment of optimal targets ranged from 20.4% in a study performed in Botswana¹¹¹ to 84.8% in a study performed in Sudan.⁹⁴

Regarding the attainment of the diabetes treatment goals in each region of Africa surveyed, the lowest and highest proportion of attainment of an optimal HbA1c goal was noted in the Central (20%, 95% CI 16 to 23)

and Western regions (37%, 95% CI 29 to 46), respectively. For the attainment of an optimal BP control, the Western region had the least proportion (31%, 95% CI 20 to 43), while the Northern region had the highest (42%, 95% CI 24 to 61). An optimal LDLC target was least achieved in the Southern region (27%, 95% CI 24 to 30) and most achieved in the Northern region (53%, 95% CI 32 to 74).

Prevalence of diabetic retinopathy, peripheral neuropathy, nephropathy, foot ulcers and peripheral arterial disease

Information on the pooled and specific prevalence of diabetes complications as reported by the different studies is summarised in tables 5–9 and as forest plots in figures 5–9.

The prevalence of diabetic retinopathy, nephropathy, peripheral neuropathy, foot ulcers and peripheral arterial disease was reported in 51 studies,^{3 19 24 26 28 30 38 41 48 51 53 54 56–58 65–67 70 72 74 76 77 81 82 86 88 89 91 95–97 102–107 109 112–116 118 120–123 125} 40 studies,^{3 19 21 27 28 30–32 38 46 48 53 57 60 62 64 66 67 69 70 76 81 82 86 88 89 91 96 97 100 105 108–110 113 114 117–119 125} 36 studies,^{3 19 25 27 28 30 33 34 37 38 43 48 51–53 55 57 58 65 67 68 73 76 79 81 85–88 96 97 105 109 118 125} 29 studies,^{3 16–19 21 22 25 27 29 38 42 43 48 49 51 53 54 57 58 67 80 85 87 95 97 113 114 125} and 18 studies,^{3 20 25 30 43 47 50 52 61 67 70 75 78 85 86 91 97 105} respectively.

Prevalence of diabetic peripheral neuropathy and retinopathy

Diabetic peripheral neuropathy and retinopathy were the most prevalent diabetes complications in the included studies with a pooled prevalence of 38% (95% CI 31 to 45, I²=98.2%) and 32% (95% CI 28 to 36, I²=98%), respectively. A wide variation was noted in the prevalence of diabetic peripheral neuropathy across the studies, with prevalence ranging from 4% in a study conducted in Eritrea⁵¹ to 83.3% in a study conducted in Nigeria.⁶⁸ A study by Makwero and colleagues¹⁰⁹ conducted in Lesotho reported the lowest prevalence of diabetic retinopathy of 4.7%, while the study by Megalla and colleagues⁹⁷ conducted in Egypt reported the highest (90%).

According to the regions of Africa surveyed, the lowest and highest prevalence of diabetic peripheral neuropathy was noted in the Central (22%, 95% CI 18 to 27) and Western regions (61%, 95% CI 45 to 75), respectively. Studies conducted in the Eastern region reported the lowest prevalence of diabetic retinopathy (23%, 95% CI

**Table 6** Prevalence of diabetic peripheral neuropathy

Prevalence of diabetic peripheral neuropathy (n=36 studies): **pooled prevalence=38% (95% CI 31 to 45, I²=98.2% 95% CI 98.7 to 99.0) and I² after meta-regression=88%.**

Prevalence of diabetic peripheral neuropathy per region: **Central: 22% (95% CI 18 to 27), Eastern: 26% (95% CI 16 to 38), Northern: 45% (95% CI 30 to 61), Southern: 46% (95% CI 42 to 49) and Western: 61% (95% CI 45 to 75).**

Author and year	No. of study participants	Country(ies)	Region of Africa	Mean age of participants	% of females	Prevalence of neuropathy, %
Abejew <i>et al</i> 2015 ¹⁹	216	Ethiopia	Eastern	45.0	42.6	14.4
Albalawi <i>et al</i> 2020 ⁸³	159	Sudan	Northern	58.1	65.4	40.3
Assaad-Khalil <i>et al</i> 2014 ⁸⁵	958	Egypt	Northern	57.3	50.0	29.3
Awadalla <i>et al</i> 2017 ⁸⁷	424	Sudan	Northern	–	49.3	68.2
Bello <i>et al</i> 2019 ⁶⁵	175	Nigeria	Western	59.8	57.7	41.7
Bentata <i>et al</i> 2015 ⁸⁸	637	Morocco	Northern	58.5	62.3	39.6
Chiwanga <i>et al</i> 2015 ²⁵	404	Tanzania	Eastern	53.6	55.4	44.0
Cohen <i>et al</i> 2010 ¹⁰⁵	620	Malawi	Southern	52.2	60.1	46.4
Deribe <i>et al</i> 2014 ²⁷	216	Ethiopia	Eastern	50.7	40.3	10.6
Dzudie <i>et al</i> 2012 ¹¹⁸	420	Cameroon	Central	56.7	51.0	22.4
Ede <i>et al</i> 2018 ⁶⁸	90	Nigeria	Western	58.6	34.4	83.3
Ekoru <i>et al</i> 2019	2784	Nigeria, Ghana, Kenya	Western and Eastern	56.0	61.0	46.0
Fasil, <i>et al</i> 2019 ²⁸	367	Ethiopia	Eastern	48.6	59.3	7.9
Gill <i>et al</i> 2008 ³⁰	105	Ethiopia	Eastern	41.0	30.0	41.0
Jarso <i>et al</i> 2011 ³³	384	Ethiopia	Eastern	–	54.1	77.0
Jember <i>et al</i> 2017 ³⁴	368	Ethiopia	Eastern	49.0	41.6	52.2
Kahloun <i>et al</i> 2014 ⁹⁶	2320	Tunisia	Northern	–	60.2	18.7
Khalil <i>et al</i> 2019 ⁸⁶	506	Egypt	Northern	–	–	20.0
Kisozi <i>et al</i> 2017 ³⁷	288	Uganda	Eastern	48.5	38.0	29.4
Kuate-Tegueu <i>et al</i> 2016 ⁷³	321	Cameroon	Western	59.8	64.1	33.3
Lebeta <i>et al</i> 2017 ³⁸	344	Ethiopia	Eastern	40.5	42.7	7.7
Makwero <i>et al</i> 2018 ¹⁰⁹	150	Lesotho	Southern	58.2	80.7	43.3
Megallaa <i>et al</i> 2019 ⁹⁷	180	Egypt	Northern	–	24.4	82.0
Miriam <i>et al</i> 2017 ⁴³	279	Ethiopia	Eastern	48.8	44.8	10.0
Mohmad <i>et al</i> 2011 ⁸¹	71	Sudan	Central	–	42.0	69.0
Neuhann <i>et al</i> 2001 ⁴⁸	474	Tanzania	Eastern	53.8	46.0	44.0
Olamoyegun <i>et al</i> 2015 ⁷⁶	90	Nigeria	Western	62.5	50.0	69.6
Seyum <i>et al</i> 2010 ⁵¹	429	Eritrea	Eastern	57.4	–	4.0
Smide <i>et al</i> 2009 ⁵²	145	Tanzania	Eastern	46.0	48.0	30.0
Sobngwi <i>et al</i> 2011 ³	2352	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	53.0	61.1	48.4
Tesfaye <i>et al</i> 2015 ⁵³	247	Ethiopia	Eastern	–	40.5	10.1
Tilahun <i>et al</i> 2017 ⁵⁴	236	Ethiopia	Eastern	47.8	46.6	25.4
Ugoya <i>et al</i> 2006 ⁷⁹	180	Nigeria	Western	53.0	51.6	75.0
Uloko <i>et al</i> 2012 ⁶⁷	531	Nigeria	Western	57.1	60.5	59.2
Vogt <i>et al</i> 2017 ⁵⁵	100	Zanzibar	Eastern	54.0	49.0	45.0
Worku <i>et al</i> 2010 ⁵⁷	305	Ethiopia	Eastern	44.4	37.1	29.5

Table 7 Prevalence of diabetic retinopathy

 Prevalence of diabetic retinopathy (n=51 studies): **pooled prevalence=32% (95% CI 28-36, I²=98% 95% CI 97.8 to 98.3) and I² after meta-regression=88.5%**.

 Prevalence of diabetic retinopathy per region: **Eastern: 23% (95% CI 19 to 28), Western: 27% (95% CI 19 to 36), Southern: 30% (95% CI 23 to 37), Central: 34% (95% CI 22 to 47) and Northern: 51% (95% CI 37 to 65)**.

Author and year	No. of study participants	Country (ies)	Region of Africa	Mean age of participants	% of females	Prevalence of retinopathy, %
Abejew <i>et al</i> 2015 ¹⁹	216	Ethiopia	Eastern	45.0	42.6	28.9
Ahmed <i>et al</i> 2017 ⁸²	316	Sudan	Northern	58.0	41.5	39.8
Albalawi <i>et al</i> 2020 ⁸³	159	Sudan	Northern	58.1	65.4	34.6
Assaad-Khalil <i>et al</i> 2019 ⁸⁵	506	Egypt	Northern	–	–	34.6
Awadalla <i>et al</i> 2017 ⁸⁷	424	Sudan	Northern	–	49.3	72.6
Bello <i>et al</i> 2019 ⁶⁵	175	Nigeria	Western	59.8	57.7	33.1
Bello <i>et al</i> 2017 ⁶⁶	358	Nigeria	Western	57.8	61.7	20.1
Bentata <i>et al</i> 2015 ⁸⁸	637	Morocco	Northern	58.5	62.3	35.6
Blake <i>et al</i> 2015 ¹⁰²	1307	Botswana	Southern	55.0	67.9	17.7
Bouaziz <i>et al</i> 2012 ⁸⁹	73	Tunisia	Northern	59.3	–	27.0
Burgress <i>et al</i> 2014 ¹⁰³	322	Malawi	Southern	55.2	64.6	50.1
Chahbi <i>et al</i> 2018 ⁹¹	300	Morocco	Northern	–	93.0	34.3
Chisha <i>et al</i> 2017 ²⁴	270	Ethiopia	Eastern	–	48.9	13.0
Cleland <i>et al</i> 2015 ²⁶	5729	Tanzania	Eastern	60.8	60.3	27.9
Cohen <i>et al</i> 2010 ¹⁰⁵	620	Malawi	Southern	52.2	60.1	34.7
Dzudie <i>et al</i> 2012 ¹¹⁸	420	Cameroon	Central	56.7	51.0	15.7
Ekoru <i>et al</i> 2019	2784	Nigeria, Ghana, Kenya	Western and Eastern	56.0	61.0	15.0
Elwali <i>et al</i> 2017 ⁹⁵	316	Sudan	Northern	58.7	40.8	82.6
Fasil <i>et al</i> 2019 ²⁸	367	Ethiopia	Eastern	48.6	59.3	17.7
Gill <i>et al</i> 2008 ³⁰	105	Ethiopia	Eastern	41.0	30.0	21.0
Glover <i>et al</i> 2011 ¹⁰⁶	281	Malawi	Southern	56.4	72.8	32.5
Hall <i>et al</i> 2017 ^{5 120}	261	Cameroon	Central	56.0	56.3	27.2
Hayfron-Benjamin <i>et al</i> 2019 ⁷⁰	206	Ghana	Western	52.9	68.9	11.0
Jingi <i>et al</i> 2014 ¹²²	407	Cameroon	Central	54.2	41.8	38.8
Jingi <i>et al</i> 2015 ¹²¹	407	Cameroon	Central	–	41.8	40.3
Kahloun <i>et al</i> 2014 ⁹⁶	2320	Tunisia	Northern	–	60.2	26.3
Kizor-Akarairwe <i>et al</i> 2018 ⁷²	80	Nigeria	Western	61.2	48.8	32.1
Lartey <i>et al</i> 2018 ⁷⁴	208	Ghana	Western	57.5	70.7	15.5
Lebeta <i>et al</i> 2017 ³⁸	344	Ethiopia	Eastern	40.5	42.7	25.5
Lewis <i>et al</i> 2018 ¹⁰⁷	921	Zambia	Southern	56.0	45.0	44.0
Magan <i>et al</i> 2019 ⁴¹	44	Uganda	Eastern	50.4	63.4	19.5
Makwero <i>et al</i> 2018 ¹⁰⁹	150	Lesotho	Southern	58.2	80.7	4.7
Megallaa <i>et al</i> , 2019 ⁹⁷	180	Egypt	Northern	–	24.4	90.0
Mohmad <i>et al</i> 2011 ⁸¹	71	Sudan	Central	–	42.0	71.2
Neuhann <i>et al</i> 2001 ⁴⁸	474	Tanzania	Eastern	53.8	46.0	14.0
Njikam <i>et al</i> 2016 ¹²³	371	Cameroon	Central	59.2	54.7	49.9
Olamoyegun <i>et al</i> 2015 ⁷⁶	90	Nigeria	Western	62.5	50.0	48.9
Onakpoya <i>et al</i> 2015 ⁷⁷	133	Nigeria	Western	–	48.1	27.8
Pirie <i>et al</i> 2014 ¹¹²	292	South Africa	Southern	59.2	79.0	39.0

Continued

**Table 7** Continued

Prevalence of diabetic retinopathy (n=51 studies): **pooled prevalence=32% (95% CI 28-36, I²=98% 95% CI 97.8 to 98.3) and I² after meta-regression=88.5%).**

Prevalence of diabetic retinopathy per region: **Eastern: 23% (95% CI 19 to 28), Western: 27% (95% CI 19 to 36), Southern: 30% (95% CI 23 to 37), Central: 34% (95% CI 22 to 47) and Northern: 51% (95% CI 37 to 65).**

Author and year	No. of study participants	Country (ies)	Region of Africa	Mean age of participants	% of females	Prevalence of retinopathy, %
Rotchford <i>et al</i> 2002 ¹¹³	253	South Africa	Southern	56.5	73.1	40.3
Seyum <i>et al</i> 2010 ⁵¹	429	Eritrea	Eastern	57.4	–	33.0
Sobngwi <i>et al</i> 2011 ³	2352	Tanzania, Kenya, Cameroon, Ghana, Senegal, and Nigeria	Eastern, Western, and Central	53.0	61.1	18.3
Tesfaye <i>et al</i> 2015 ⁵³	247	Ethiopia	Eastern	–	40.5	11.7
Thinyane <i>et al</i> 2013 ¹¹⁴	80	Lesotho	Southern	49.0	49.0	35.0
Thomas <i>et al</i> 2013 ¹¹⁵	3978	South Africa	Southern	56.8	33.3	20.5
Tilahun <i>et al</i> 2017 ⁵⁴	236	Ethiopia	Eastern	47.8	46.6	20.3
Uloko <i>et al</i> 2012 ⁶⁷	531	Nigeria	Western	57.1	60.5	35.5
Webb <i>et al</i> 2016 ¹¹⁶	599	South Arica	Southern	57.8	68.0	24.9
Woodward <i>et al</i> 2020 ⁵⁶	91	Tanzania	Eastern	59.2	62.6	42.9
Worku <i>et al</i> 2010 ⁵⁷	305	Ethiopia	Eastern	44.4	37.1	33.8

19 to 28) while studies conducted in the Northern region reported the highest prevalence (51%, 95% CI 37 to 65).

Prevalence of diabetic nephropathy, peripheral arterial disease and foot ulcers

The pooled prevalence of diabetic nephropathy, peripheral arterial disease and foot ulcers in the included studies was 31% (95% CI 22 to 41, I²=99.3%), 19% (95% CI 12 to 25, I²=98.1%) and 11% (95% CI 9 to 14, I²=97.4%), respectively.

The prevalence of diabetic nephropathy and peripheral arterial disease ranged from 2.2% in Ethiopia¹⁹ to 90% in Nigeria⁶⁴ and 2.7% in a study performed in Morocco⁹¹ to 52.5% in a study performed in Nigeria,⁷⁸ respectively. Regarding the burden of diabetic foot ulcers, there was also an observed heterogeneity, with prevalence ranging from 0.4% in Ethiopia⁵³ to 86.7% in Egypt.⁹⁷

Studies conducted in the Central, Eastern and Southern regions reported a comparable prevalence of diabetic nephropathy (22%, 25% and 28%, respectively) with the highest prevalence reported in studies conducted in the Western region (47%). Regarding the prevalence of PAD, studies conducted in the Southern (8%, 95% CI 6 to 10) and Western (29%, 95% CI 13 to 48) regions reported the lowest and highest prevalence, respectively. A comparable prevalence of diabetic foot ulcers was noted in studies conducted in the Southern, Western and Eastern regions (7%, 8% and 10%, respectively), with the highest prevalence noted in studies conducted in the Northern region (21%).

On sensitivity analysis considering only high-quality studies, the pooled prevalence of the five diabetic complications and the proportion of attainment of the three

optimal diabetes treatment goals did not differ from those obtained in the preliminary analysis with all eligible studies included. The pooled prevalence of diabetic foot ulcers, peripheral arterial disease, diabetic nephropathy, diabetic retinopathy and diabetic peripheral neuropathy after sensitivity analysis was 9% (95% CI 7 to 12, I²=92.9%), 20% (95% CI 13 to 28, I²=98.4%), 31% (95% CI 21 to 42, I²=99.4%), 33% (95% CI 28 to 37, I²=98.2%) and 40% (95% CI 32 to 48, I²=99%), respectively. The pooled proportion of attainment of optimal HbA1c, BP and LDLC treatment goal was 27% (95% CI 23 to 30, I²=94.5%), 37% (95% CI 29 to 46, I²=99.0%) and 43% (95% CI 31 to 55, I²=97.9%), respectively.

DISCUSSION

To our knowledge, this is the first systematic review and meta-analysis to simultaneously document the proportion of attainment of the three key indicators of optimal diabetes care (HbA1c, BP, and LDLC goals) and the burden of five diabetes complications in an indigenous adult population with type 2 diabetes in Africa. In this study of a total of 63 890 study participants, we report that, generally, a small proportion of adult patients with type 2 diabetes in Africa attain optimal diabetes treatment targets, especially HbA1c and BP goals (less than 40%). In addition, diabetes complications are relatively common with diabetic neuropathy being the most prevalent (38%) followed by diabetic retinopathy (32%), nephropathy (31%), peripheral arterial disease (19%) and foot ulcers (11%).

Table 8 Prevalence of diabetic foot ulcers

Prevalence of diabetic foot ulcers (n=29 studies): **pooled prevalence=11% (95% CI 9 to 14, I²=97.4% 95% CI 96.9 to 97.8), and I² after meta-regression :1.4%).**
 Prevalence of diabetic foot ulcers per region: **Southern: 7% (95% CI 5 to 11), Western: 8% (95% CI 6 to 10), Eastern: 10% (95% CI 8 to 12) and Northern: 21% (95% CI 4 to 48).**

Author and year	No. of study participants	Country(ies)	Region of Africa	Mean age of participants	% of females	Prevalence of foot ulcers, %
Abbas <i>et al</i> 2002 ¹⁶	627	Tanzania	Eastern	53.0	35.0	15.0
Abbas <i>et al</i> 2011 ¹⁷	11 866	Tanzania	Eastern	–	–	12.0
Abdissa <i>et al</i> 2020 ¹⁸	229	Ethiopia	Eastern	–	40.4	12.7
Abejew <i>et al</i> 2015 ¹⁹	216	Ethiopia	Eastern	45.0	42.6	4.4
Albalawi <i>et al</i> 2020 ⁸³	159	Sudan	Northern	58.1	65.4	2.5
Amour <i>et al</i> 2019 ²¹	315	Tanzania	Eastern	57.2	65.7	10.0
Assaad-Khalil <i>et al</i> 2014 ⁸⁵	958	Egypt	Northern	57.3	50.0	6.1
Awadalla <i>et al</i> 2017 ⁸⁷	424	Sudan	Northern	–	49.3	12.7
Chalya <i>et al</i> 2011 105 ²²	136	Tanzania	Eastern	54.3	45.6	3.2
Chiwanga <i>et al</i> 2015 ²⁵	404	Tanzania	Eastern	53.6	55.4	15.0
Deribe <i>et al</i> 2014 ²⁷	216	Ethiopia	Eastern	50.7	40.3	14.8
Ekoru K <i>et al</i> 2019	2784	Nigeria, Ghana, Kenya	Western and Eastern	56.0	61.0	5.0
Elwali <i>et al</i> 2017 ⁹⁵	316	Sudan	Northern	58.7	40.8	17.7
Gebrekirstos <i>et al</i> 2015 ²⁹	228	Ethiopia	Eastern	–	38.0	12.0
Lebeta <i>et al</i> 2017 ³⁸	344	Ethiopia	Eastern	40.5	42.7	21.2
Mamo <i>et al</i> 2015 ⁴²	200	Ethiopia	Eastern	50.0	72.5	15.0
Mariam <i>et al</i> 2017 ⁴³	279	Ethiopia	Eastern	48.8	44.8	13.6
Megallaa <i>et al</i> 2019 ⁹⁷	180	Egypt	Northern	–	24.4	86.7
Neuhann <i>et al</i> 2001 ⁴⁸	474	Tanzania	Eastern	53.8	46.0	10.0
Nyamu <i>et al</i> 2003 ⁴⁹	1788	Kenya	Eastern	56.9	–	4.6
Rotchford <i>et al</i> 2002 ¹¹³	253	South Africa	Southern	56.5	73.1	6.0
Seyum <i>et al</i> 2010 ⁵¹	429	Eritrea	Eastern	57.4	–	14.0
Sobngwi <i>et al</i> 2011 ³	2352	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	53.0	61.1	11.7
Tesfaye <i>et al</i> 2015 ⁵³	247	Ethiopia	Eastern	–	40.5	0.4
Thinyane <i>et al</i> 2013 ¹¹⁴	80	Lesotho	Southern	49.0	49.0	14.0
Tilahun <i>et al</i> 2017 ⁵⁴	236	Ethiopia	Eastern	47.8	46.6	8.5
Uloko <i>et al</i> 2012 ⁶⁷	531	Nigeria	Western	57.1	60.5	3.8
Unachukwu <i>et al</i> 2006 ⁸⁰	315	Nigeria	Western	54.6	36.7	19.1
Worku <i>et al</i> 2010 ⁵⁷	305	Ethiopia	Eastern	44.4	37.1	4.6

Proportions of attainment of the optimal diabetes treatment goals

A wide heterogeneity in the attainment of the optimal diabetes treatment goals was noted across all five regions of Africa. This could probably be explained by the marked differences in the populations studied, health-care systems and knowledge-practice gaps among health-care practitioners.

Similar to our study findings, achievement of optimal HbA1c, BP and LDLC treatment goals has also been widely reported to be a significant clinical challenge in several studies performed in Caucasian and Asian populations with type 2 diabetes in high-income and middle-income countries.^{126–131} In one large registry-based study of >100 000 adults with a self-reported diagnosis of diabetes carried out between 1999 and 2010 in USA,

Table 9 Prevalence of peripheral arterial disease

Prevalence of peripheral arterial disease (PAD) (n=18 studies): **Pooled prevalence=19% (95% CI 12 to 25, I²=98.1% 95% CI 97.6 to 98.4) and I² after meta-regression: 70.9%**.
 Prevalence of PAD per region: **Southern: 8% (95% CI 6 to 10), Northern: 15% (95% CI 4 to 29), Eastern: 18% (95% CI 11 to 27) and Western: 29% (95% CI 13 to 48).**

Author and year	No. of study participants	Country(ies)	Region of Africa	Mean age of participants	% of females	Prevalence of PAD, %
Agboghoroma <i>et al</i> 2020 ⁶¹	200	Nigeria	Western	–	–	38.5
Akalu <i>et al</i> 2020 ²⁰	280	Ethiopia	Eastern	–	38.6	30.7
Assaad-Khalil <i>et al</i> 2014 ⁸⁵	958	Egypt	Northern	57.3	50.0	11.0
Chahbi <i>et al</i> 2018 ⁹¹	300	Morocco	Northern	–	93.0	2.7
Chiwanga <i>et al</i> 2015 ²⁵	404	Tanzania	Eastern	53.6	55.4	15.0
Cohen <i>et al</i> 2010 ¹⁰⁵	620	Malawi	Southern	52.2	60.1	7.6
Gill <i>et al</i> 2008 ³⁰	105	Ethiopia	Eastern	41.0	30.0	6.0
Hayfron-Benjamin <i>et al</i> 2019 ⁷⁰	206	Ghana	Western	52.9	68.9	11.2
Khalil <i>et al</i> 2019 ⁸⁶	506	Egypt	Northern	–	–	32.6
Mariam <i>et al</i> 2017 ⁴³	279	Ethiopia	Eastern	48.8	44.8	9.7
Megallaa <i>et al</i> 2019 ⁹⁷	180	Egypt	Northern	–	24.4	20.0
Mwebaze <i>et al</i> 2014 ⁴⁷	146	Uganda	Eastern	53.9	48.6	39.0
Ogbera <i>et al</i> 2015 ⁷⁵	225	Nigeria	Western	61.4	57.0	40.0
Okello <i>et al</i> 2014 ⁵⁰	229	Uganda	Eastern	60.0	63.7	24.0
Oyelade <i>et al</i> 2012 ⁷⁸	219	Nigeria	Western	–	58.9	52.5
Smide <i>et al</i> 2008 ⁵²	145	Tanzania	Eastern	46.0	48.0	13.0
Sobngwi <i>et al</i> 2011 ³	2352	Tanzania, Kenya, Cameroon, Ghana, Senegal and Nigeria	Eastern, Western and Central	53.0	61.1	4.7
Uloko <i>et al</i> 2012 ⁶⁷	531	Nigeria	Western	57.1	60.5	10.7

33.4%–48.7% of adult patients with diabetes did not achieve the recommended HbA1c, BP and LDLC treatment targets. Less than 15% met all the three treatment targets in addition to smoking cessation.¹²⁶

Similarly, a low proportion of achievement of an optimal HbA1c target was also reported by a large international, multicentre observational study of 2704 multiracial adult populations with diabetes from 10 countries (two from Africa, five from the Middle East and three from South Asia). About 46% of the participants were Caucasian. An optimal HbA1c goal of <7% (53 mmol/mol) was reported in only 25.8% of the participants.¹²⁸

In the Japan Epidemiology Collaboration on Occupational Health study, which enrolled 3070 adult employees of large manufacturing companies, optimal HbA1c, BP and LDLC goals as recommended by the ADA were noted in 44.9%, 76.6% and 27.1% of participants, respectively. Only 11.2% of participants attained all three treatment goals.¹²⁹

The burden of diabetes complications in Africa

Regarding studies on the burden of diabetes complications in Africa, there were few that investigated the prevalence of diabetic foot ulcers and peripheral arterial

disease with diabetic retinopathy, peripheral nephropathy and neuropathy being the most studied. Diabetic peripheral neuropathy and retinopathy remain the most prevalent diabetes complication and diabetic foot ulcers the least prevalent.

With regards to the prevalence of diabetic foot ulcers, an earlier published systematic review and meta-analysis on the characteristics, prevalence and outcomes of diabetic foot ulcers in Africa by Rigato *et al*¹³² reported a pooled prevalence of diabetic foot ulcers of 13%, a finding close to what we observed (11%). In another systematic review and meta-analysis on the prevalence of diabetic peripheral neuropathy in African populations with DM, Shiferaw *et al*¹³³ reported a slightly higher overall prevalence of 46% compared with what we found in our study (38%) while including fewer studies (n=23).

Similar to our study, considerable heterogeneity was also reported in the documented prevalence of the varied diabetes complications in Africa in most previously published systematic reviews. This may be due to variations in clinical definitions of diabetes complications in the studies. Burgess *et al*¹³⁴ and Achigbu *et al*¹³⁵ reported a wide disparity in the prevalence of diabetic retinopathy




Figure 5 Forest plot summarising studies on the prevalence of diabetic retinopathy. ES, effect size.

in the included studies of 7%–62.4%, and 13%–82.6%, respectively. Noubiap *et al*¹³⁶ in a systematic review on the burden of diabetic nephropathy in 2015 reported an overall prevalence of chronic kidney disease in patients with diabetes ranging between 11% and 83.7%. Johnston *et al* in a systematic review that aimed to assess the




Figure 6 Forest plot summarising studies on the prevalence of diabetic foot ulcers. ES, effect size.




Figure 7 Forest plot summarising studies on the prevalence of diabetic nephropathy. ES, effect size.

epidemiological and clinical reports regarding Peripheral arterial disease (PAD) in Sub-saharan Africa (SSA) documenting the prevalence of PAD in patients with diabetes as reported by three studies to range from 39% to 52%.¹³⁷

Compared with Caucasian and Asian adult populations with type 2 diabetes, our study has demonstrated that




Figure 8 Forest plot summarising studies on the prevalence of diabetic neuropathy. ES, effect size.




Figure 9 Forest plot summarising studies on the prevalence of peripheral arterial disease. ES, effect size.

adult African patients are disproportionately affected by complications of DM. The Joint Asia Diabetes Evaluation programme that undertook comprehensive risk assessments of 3687 adult patients with type 2DM recruited from seven Asian countries reported a prevalence of peripheral arterial disease, diabetic neuropathy, macroalbuminuria and microalbuminuria and diabetic retinopathy of 3.1%, 15%, 18.8% and 20.4%, respectively.¹³⁸

The National Health and Nutrition Examination Survey conducted from 1988 to 1994 and 1999–2018 in USA in 1486 non-pregnant adults (aged ≥ 20 years) with newly diagnosed diabetes (diagnosed within the past 2 years) also documented a low burden of most diabetes complications. Diabetic foot ulcers, peripheral arterial disease, diabetic retinopathy, neuropathy and nephropathy (albuminuria) were prevalent in 6.3%, 9.2%, 12.1%, 14.5% and 18.7%, respectively.¹³⁹

The documented low proportions of attainment of optimal diabetes treatment goals (optimal HbA1c, BP and LDLC targets) in Africa is associated with an increased risk of onset and progression of diabetes complications, hence increasing morbidity and mortality in addition to causing a significant economic strain on the meagre health resources. This generally observed low proportion of attainment of key diabetes treatment goals and high prevalence of diabetes complications, notably diabetic neuropathy, retinopathy and nephropathy in Africa, exists broadly due to challenges related to screening, diagnosis and management of DM.

Awareness of diabetes in the general African population and healthcare practitioners remains very poor, resulting in delayed diagnosis of diabetes. The challenge of ready access to affordable essential diabetes medicines like insulin and statins and diagnostic tests or equipment like glucometers for home self-monitoring of glucose,

HbA1c and lipid profile tests remains highly prevalent in most African countries.^{140–144}

Effective management of diabetes and its related cardiovascular risk factors like hypertension and dyslipidaemia in most healthcare settings in Africa also remains a significant clinical challenge.³ Most healthcare facilities especially the lower tier ones lack local or institution-specific comprehensive diabetes treatment guidelines to guide healthcare practitioners on how to optimally manage diabetes, in addition to the evident knowledge–practice gaps among healthcare practitioners.²

Healthcare systems in most African countries remain poorly structured to optimally manage most NCDs like diabetes along with an inadequately funded health sector. Most African countries have not yet fulfilled the 2001 Abuja Declaration of allocating 15% of their national annual budget to the health sector.^{2 145}

This systematic review and meta-analysis had its strengths and limitations. To our knowledge, it is the first to simultaneously investigate the status of attainment of the three key diabetes treatment goals and the burden of five common diabetes complications in an adult indigenous African population with type 2 diabetes. The systematic review and meta-analysis included a large number of studies that assessed the extent of attainment of diabetes treatment goals and the prevalence of diabetes complications based on recommendations or definitions by internationally recognised associations.

It also had its limitations. There was considerable heterogeneity in the included studies. This could be explained by the differences in study sites (tertiary vs low-tier hospitals or private vs public hospitals), patient characteristics (age, duration of diabetes, coexisting medical conditions), regions where the studies were conducted and diagnostic modalities used to identify diabetes complications. The systematic review also excluded studies published in French, which is the official language of some African countries. However, these were very few. There was evidence of publication bias in some of the included studies especially studies investigating the prevalence of diabetic nephropathy and peripheral neuropathy and the proportion of attainment of an optimal BP goal. About 23% of the included studies were moderate and low quality on assessment using the NOS for cross-sectional studies.

CONCLUSION

Achievement of optimal diabetes treatment goals, especially HbA1c and BP, in adult African patients with type 2 diabetes remains low in Africa. Diabetes complications especially diabetic peripheral neuropathy and retinopathy also remain highly prevalent. Implementation of universal diabetes screening and education initiatives coupled with improving knowledge about diabetes management among healthcare practitioners and ready access to affordable essential diabetes diagnostic tests and medicines in Africa are integral in improving

overall optimal diabetes care and reducing the burden of diabetes complications.

Considering the projected future increase in the prevalence of diabetes globally, especially in the African region, there is an urgent need to address glaring gaps in diabetes care and to develop simple and pragmatic interventions to improve treatment outcomes and reduce the burden of diabetes complications.

Author affiliations

- ¹Department of Medicine, Lubaga Hospital, Kampala, Uganda
²Department of Internal Medicine, Kilimanjaro Christian Medical Centre, Moshi, Kilimanjaro, Tanzania
³Department of Medicine, Kilimanjaro Christian Medical University College, Moshi, Kilimanjaro, Tanzania
⁴Department of Internal Medicine, Makerere University College of Health Sciences, Kampala, Uganda
⁵Department of Immunomodulation and Vaccines, MRC/UVRI and LSHTM Uganda Research Unit, Entebbe, Uganda
⁶Non-Communicable Diseases Program, Medical Research Council/Uganda Virus Research Institute & London School of Hygiene and Tropical Medicine Uganda Research Unit, Entebbe, Uganda
⁷Department of Immunology, Makerere University College of Health Sciences, Kampala, Uganda
⁸Department of Medical Microbiology and Immunology, Faculty of Medicine, Gulu University, Gulu, Uganda
⁹Department of Medicine, NIMR-Mbeya Medical Research Programme, Mbeya, Mbeya, Tanzania
¹⁰Department of Medical Statistics, NIMR-Mbeya Medical Research Programme, Mbeya, Mbeya, Tanzania
¹¹Department of Paediatrics and Child Health, NIMR-Mbeya Medical Research Programme, Mbeya, Mbeya, Tanzania
¹²Centre for International Health, University of Otago, Dunedin, New Zealand
¹³Department of Pharmacology, Radboud University Nijmegen, Nijmegen, Gelderland, The Netherlands
¹⁴Department of Medicine, Radboud University Nijmegen, Nijmegen, Gelderland, The Netherlands
¹⁵Department of Internal Medicine, Radboud University Nijmegen, Nijmegen, Gelderland, The Netherlands
¹⁶University of Oxford Centre for Tropical Medicine and Global Health, Oxford, UK
¹⁷Population Health Research Institute, St George's University of London, London, UK

Twitter Phillip Ssekamatte @PWSsekamatte

Contributors DK and NC: conceived the research idea, performed the preliminary screening of titles and abstracts to identify potentially eligible articles and wrote the initial draft of the manuscript; DK, NC, IA-B, SNL, ISe, APK and SN: retrieved full texts and identified the eligible articles; KK, SNL, AP-K, SN, PS, FB, LEM, WO, TDM, NEN, ISA: extracted data from the identified eligible articles; DK and ISe performed the data analysis and interpretation; NC, KK and SNL: performed the assessment of the quality of studies; KS, PH, LtB, JV, RvC and JAC: offered additional data interpretation and supervised this work. All the authors reviewed the different versions of the manuscript and read and approved the final draft of the manuscript. DK is the overall guarantor and accepts full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and

responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Davis Kibirige <http://orcid.org/0000-0001-5127-3031>
 Phillip Ssekamatte <http://orcid.org/0000-0002-3418-8414>
 Felix Bongomin <http://orcid.org/0000-0003-4515-8517>
 Julia Alison Critchley <http://orcid.org/0000-0002-5248-4188>

REFERENCES

- 1 IDF. *International diabetes Federation diabetes atlas*. 10th edition, 2021. <https://diabetesatlas.org>
- 2 Atun R, Davies JI, Gale EAM, *et al*. Diabetes in sub-Saharan Africa: from clinical care to health policy. *Lancet Diabetes Endocrinol* 2017;5:622–67.
- 3 Sobngwi E, Ndur-Mbaye M, Boateng KA, *et al*. Type 2 diabetes control and complications in specialised diabetes care centres of six sub-Saharan African countries: the Diabcare Africa study. *Diabetes Res Clin Pract* 2012;95:30–6.
- 4 Gill GV, Mbanya J-C, Ramaiya KL, *et al*. A sub-Saharan African perspective of diabetes. *Diabetologia* 2009;52:8–16.
- 5 Hall V, Thomsen RW, Henriksen O, *et al*. Diabetes in sub Saharan Africa 1999–2011: epidemiology and public health implications. A systematic review. *BMC Public Health* 2011;11:564.
- 6 Nuche-Berenguer B, Kupfer LE. Readiness of sub-Saharan Africa healthcare systems for the new pandemic, diabetes: a systematic review. *J Diabetes Res* 2018;2018:9262395.
- 7 American Diabetes Association. 9. Cardiovascular Disease and Risk Management: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S125–50.
- 8 American Diabetes Association. 9. Pharmacologic Approaches to Glycemic Treatment: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S111.
- 9 IDF. International Diabetes Federation. IDF clinical practice recommendations for managing type 2 diabetes in primary care, 2018. Available: <https://www.idf.org/e-library/guidelines/128-idf-clinical-practice-recommendations-for-managing-type-2-diabetes-in-primary-care.html> [Accessed 17 Sep 2021].
- 10 Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 11 American Diabetes Association. 2. Classification and Diagnosis of Diabetes: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S15–33.
- 12 Wells GA, Shea B, O'Connell D, *et al*. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses, 2021. Available: http://www.ohrica.com/clinical_epidemiology/oxfordasp [Accessed 02 Sep 2021].
- 13 DerSimonian R, Laird N. Meta-Analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
- 14 Cumpston M, Li T, Page MJ, *et al*. Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for systematic reviews of interventions. *Cochrane Database Syst Rev* 2019;10:ED000142.
- 15 Egger M, Davey Smith G, Schneider M, *et al*. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- 16 Gulam-Abbas Z, Lutale JK, Morbach S, *et al*. Clinical outcome of diabetes patients hospitalized with foot ulcers, Dar ES Salaam, Tanzania. *Diabet Med* 2002;19:575–9.
- 17 Abbas ZG, Lutale JK, Bakker K, *et al*. The 'Step by Step' Diabetic Foot Project in Tanzania: a model for improving patient outcomes in less-developed countries. *Int Wound J* 2011;8:169–75.
- 18 Abdissa D, Adugna T, Gerema U, *et al*. Prevalence of diabetic foot ulcer and associated factors among adult diabetic patients on follow-up clinic at Jimma medical center, Southwest Ethiopia, 2019: an Institutional-Based cross-sectional study. *J Diabetes Res* 2020;2020:1–6.

- 19 Abejew AA, Belay AZ, Kerie MW. Diabetic complications among adult diabetic patients of a tertiary hospital in northeast Ethiopia. *Adv Public Health* 2015;2015:1–7.
- 20 Akalu Y, Belsti Y. Hypertension and its associated factors among type 2 diabetes mellitus patients at Debre Tabor General Hospital, Northwest Ethiopia. *Diabetes Metab Syndr Obes* 2020;13:1621–31.
- 21 Amour AA, Chamba N, Kayandabila J, et al. Prevalence, patterns, and factors associated with peripheral neuropathies among diabetic patients at tertiary hospital in the Kilimanjaro region: descriptive cross-sectional study from north-eastern Tanzania. *Int J Endocrinol* 2019;2019:1–7.
- 22 Chalya PL, Mabula JB, Dass RM, et al. Surgical management of diabetic foot ulcers: a Tanzanian university teaching hospital experience. *BMC Res Notes* 2011;4:365.
- 23 Chamba NG, Shao ER, Tolbert S, et al. Lipid profile of type 2 diabetic patients at a tertiary hospital in Tanzania: cross sectional study. *J Endocrinol Diab* 2017;4:1–6.
- 24 Chisha Y, Terefe W, Assefa H, et al. Prevalence and factors associated with diabetic retinopathy among diabetic patients at Arbaminch General Hospital, Ethiopia: cross sectional study. *PLoS One* 2017;12:e0171987.
- 25 Chiwanga FS, Njelekela MA. Diabetic foot: prevalence, knowledge, and foot self-care practices among diabetic patients in Dar es Salaam, Tanzania - a cross-sectional study. *J Foot Ankle Res* 2015;8:20.
- 26 Cleland CR, Burton MJ, Hall C, et al. Diabetic retinopathy in Tanzania: prevalence and risk factors at entry into a regional screening programme. *Trop Med Int Health* 2016;21:417–26.
- 27 Deribe B, Woldemichael K, Nemera G. Prevalence and factors influencing diabetic foot ulcer among diabetic patients attending Arbaminch Hospital, South Ethiopia. *J Diabetes Metab Disord* 2014;2:322.
- 28 Fasil A, Biadgo B, Abebe M. Glycemic control and diabetes complications among diabetes mellitus patients attending at University of Gondar Hospital, Northwest Ethiopia. *Diabetes Metab Syndr Obes* 2018;12:75–83.
- 29 Gebrekirstos K, Gebrekiros S, Fantahun A. Prevalence and factors associated with diabetic foot ulcer among adult patients in Ayder referral hospital diabetic clinic Mekelle, North Ethiopia, 2013. *J Diabetes Metab* 2015;6:2.
- 30 Gill G, Gebrekidan A, English P, et al. Diabetic complications and glycaemic control in remote North Africa. *QJM* 2008;101:793–8.
- 31 Kumela Goro K, Desalegn Wolide A, Kerga Dibaba F, et al. Patient awareness, prevalence, and risk factors of chronic kidney disease among diabetes mellitus and hypertensive patients at Jimma University medical center, Ethiopia. *Biomed Res Int* 2019;2019:1–8.
- 32 Janmohamed MN, Kalluvya SE, Mueller A, et al. Prevalence of chronic kidney disease in diabetic adult out-patients in Tanzania. *BMC Nephrol* 2013;14:183–83.
- 33 Jarso G, Ahmed A, Feleke Y. The prevalence, clinical features and management of peripheral neuropathy among diabetic patients in Tikur Anbessa and St. Paul's specialized university hospitals, Addis Ababa, Ethiopia. *Ethiop Med J* 2011;49:299–311.
- 34 Jember G, Melsew YA, Fisseha B, et al. Peripheral sensory neuropathy and associated factors among adult diabetes mellitus patients in Bahr Dar, Ethiopia. *J Diabetes Metab Disord* 2017;16:16.
- 35 Kibirige D, Akabwai GP, Kampiire L, et al. Frequency and predictors of suboptimal glycemic control in an African diabetic population. *Int J Gen Med* 2017;10:33–8.
- 36 Kimando MW, Otieno FCF, Ogola EN, et al. Adequacy of control of cardiovascular risk factors in ambulatory patients with type 2 diabetes attending diabetes out-patients clinic at a County Hospital, Kenya. *BMC Endocr Disord* 2017;17:73.
- 37 Kisozi T, Mutebi E, Kisekka M, et al. Prevalence, severity and factors associated with peripheral neuropathy among newly diagnosed diabetic patients attending Mulago Hospital: a cross-sectional study. *Afr Health Sci* 2017;17:463–73.
- 38 Reba Lebeta K, Lebeta KR, Argaw Z. Prevalence of diabetic complications and its associated factors among diabetes mellitus patients attending diabetes mellitus clinics; institution based cross sectional study. *American Journal of Health Research* 2017;5:38–43.
- 39 Lumu W, Kampiire L, Akabwai GP, et al. Dyslipidaemia in a black African diabetic population: burden, pattern and predictors. *BMC Res Notes* 2017;10:587.
- 40 Lumu W, Kampiire L, Akabwai GP, et al. Statin therapy reduces the likelihood of suboptimal blood pressure control among Ugandan adult diabetic patients. *Ther Clin Risk Manag* 2017;13:215–21.
- 41 Magan T, Pouncey A, Gadhi K, et al. Prevalence and severity of diabetic retinopathy in patients attending the endocrinology diabetes clinic at Mulago hospital in Uganda. *Diabetes Res Clin Pract* 2019;152:65–70.
- 42 Mamo T, Yifter H, Lemessa T. Risk factors assessment of diabetic foot ulcer using the sixty seconds screening tool: a hospital based cross-sectional study at TIKUR ANBESSA specialized Hospital. *Ethiop Med J* 2015;2:45–9.
- 43 Mariam TG, Alemayehu A, Tesfaye E, et al. Prevalence of diabetic foot ulcer and associated factors among adult diabetic patients who attend the diabetic follow-up clinic at the University of Gondar referral Hospital, North West Ethiopia, 2016: Institutional-Based cross-sectional study. *J Diabetes Res* 2017;2017:1–8.
- 44 Mbwete GW, Kilonzo KG, Shao ER, et al. Suboptimal blood pressure control, associated factors, and choice of antihypertensive drugs among type 2 diabetic patients at KCMC, Tanzania. *J Diabetes Res* 2020;2020:1–9.
- 45 Muddu M, Mutebi E, Mondo C. Prevalence, types and factors associated with echocardiographic abnormalities among newly diagnosed diabetic patients at Mulago Hospital. *Afr Health Sci* 2016;16:183–93.
- 46 Muddu M, Mutebi E, Ssinabulya I, et al. Utility of albumin to creatinine ratio in screening for microalbuminuria among newly diagnosed diabetic patients in Uganda: a cross sectional study. *Afr Health Sci* 2019;19:1607–16.
- 47 Mwebaze RM, Kibirige D. Peripheral arterial disease among adult diabetic patients attending a large outpatient diabetic clinic at a national referral hospital in Uganda: a descriptive cross sectional study. *PLoS One* 2014;9:e105211.
- 48 Neuhann HF, Warter-Neuhann C, Lyaruu I, et al. Diabetes care in Kilimanjaro region: clinical presentation and problems of patients of the diabetes clinic at the regional referral hospital-an inventory before structured intervention. *Diabet Med* 2002;19:509–13.
- 49 Nyamu PN, Otieno CF, Amayo EO, et al. Risk factors and prevalence of diabetic foot ulcers at Kenyatta national Hospital, Nairobi. *East Afr Med J* 2003;80:36–43.
- 50 Okello S, Millard A, Owori R, et al. Prevalence of lower extremity peripheral artery disease among adult diabetes patients in southwestern Uganda. *BMC Cardiovasc Disord* 2014;14:75.
- 51 Seyum B, Mebrahtu G, Usman A, et al. Profile of patients with diabetes in Eritrea: results of first phase registry analyses. *Acta Diabetol* 2010;47:23–7.
- 52 Smide B. Outcome of foot examinations in Tanzanian and Swedish diabetic patients, a comparative study. *J Clin Nurs* 2009;18:391–8.
- 53 Tesfaye DJ, Tessema F, Taha M. Coexistence of chronic complications among diabetic patients at Nigist Eleni Mohammed Memorial Hospital, Hossana, South Ethiopia. *OALib* 2015;02:1–10.
- 54 Tilahun AN, Waktola C, Tewodros GM, et al. Major micro vascular complications and associated risk factors among diabetic outpatients in Southwest Ethiopia. *Endocrinol Metab Syndr* 2017;6:4.
- 55 Vogt EC, Øksnes M, Suleiman F, et al. Assessment of diabetic polyneuropathy in Zanzibar: comparison between traditional methods and an automated point-of-care nerve conduction device. *J Clin Transl Endocrinol* 2017;10:9–14.
- 56 Woodward R, Mgaya E, Mwanansao C, et al. Retinopathy in adults with hypertension and diabetes mellitus in Western Tanzania: a cross-sectional study. *Trop Med Int Health* 2020;25:1214–25.
- 57 Worku D, Hamza L, Woldemichael K. Patterns of diabetic complications at jimma university specialized Hospital, Southwest Ethiopia. *Ethiop J Health Sci* 2010;20:33–9.
- 58 Ekoru K, Doumatey A, Bentley AR, et al. Type 2 diabetes complications and comorbidity in sub-Saharan Africans. *EClinicalMedicine* 2019;16:30–41.
- 59 Camara A, Baldé NM, Sobngwi-Tambekou J, et al. Poor glycemic control in type 2 diabetes in the South of the Sahara: the issue of limited access to an HbA1c test. *Diabetes Res Clin Pract* 2015;108:187–92.
- 60 Adetunji OR, Adeleye JO, Agada NO, et al. Microalbuminuria and clinical correlates in black African patients with type 2 diabetes. *West Afr J Med* 2006;25:279–83.
- 61 Agboghoro OF, Akemokwe FM, Puepet FH. Peripheral arterial disease and its correlates in patients with type 2 diabetes mellitus in a teaching hospital in northern Nigeria: a cross-sectional study. *BMC Cardiovasc Disord* 2020;20:102.
- 62 Alebiosu CO. Clinical diabetic nephropathy in a tropical African population. *West Afr J Med* 2003;22:152–5.
- 63 Attoye TE, Adebobola PA, Inem V. An assessment of glycaemic control and modes of health financing among type 2 diabetic patients attending a teaching hospital in south-western Nigeria. *West Afr J Med* 2020;37:237–47.
- 64 Balogun WO, Abbiyesuku FM. Excess renal insufficiency among type 2 diabetic patients with dip-stick positive proteinuria in a tertiary hospital. *Afr J Med Med Sci* 2011;40:399–403.

- 65 Bello A, Biliaminu S, Wahab K, *et al.* Distal symmetrical polyneuropathy and cardiovascular autonomic neuropathy among diabetic patients in Ilorin: prevalence and predictors. *Niger Postgrad Med J* 2019;26:123–8.
- 66 Bello BT, Amira CO. Pattern and predictors of urine protein excretion among patients with type 2 diabetes attending a single tertiary hospital in Lagos, Nigeria. *Saudi J Kidney Dis Transpl* 2017;28:1381–8.
- 67 Uloko AE, Ofoegbu EN, Chinenye S, *et al.* Profile of Nigerians with diabetes mellitus - Diabcare Nigeria study group (2008): Results of a multicenter study. *Indian J Endocrinol Metab* 2012;16:558–64.
- 68 Ede O, Eyichukwu GO, Madu KA, *et al.* Evaluation of peripheral neuropathy in diabetic adults with and without foot ulcers in an African population. *Journal of Biosciences and Medicines* 2018;06:71–8.
- 69 Eghan BA, Frempong MT, Adjei-Poku M. Prevalence and predictors of microalbuminuria in patients with diabetes mellitus: a cross-sectional observational study in Kumasi, Ghana. *Ethn Dis* 2007;17:726–30.
- 70 Hayfron-Benjamin C, van den Born B-J, Maitland-van der Zee AH, *et al.* Microvascular and macrovascular complications in type 2 diabetes Ghanaian residents in Ghana and Europe: the RODAM study. *J Diabetes Complications* 2019;33:572–8.
- 71 Iwuala S, Olamoyegun M, Sabir A, *et al.* The relationship between self-monitoring of blood glucose and glycaemic control among patients attending an urban diabetes clinic in Nigeria. *Ann Afr Med* 2015;14:182–7.
- 72 Kizor-Akaraiwe NN, Ezegwui IR, Oguego N, *et al.* Prevalence, awareness and determinants of diabetic retinopathy in a screening centre in Nigeria. *J Community Health* 2016;41:767–71.
- 73 Kuate-Tegueu C, Temfack E, Ngankou S, *et al.* Prevalence and determinants of diabetic polyneuropathy in a sub-Saharan African referral hospital. *J Neurol Sci* 2015;355:108–12.
- 74 Lartey SY, Aikins AK. Visual impairment amongst adult diabetics attending a tertiary outpatient clinic. *Ghana Med J* 2018;52:84–7.
- 75 Ogbera AO, Adeleye O, Solagberu B, *et al.* Screening for peripheral neuropathy and peripheral arterial disease in persons with diabetes mellitus in a Nigerian university teaching hospital. *BMC Res Notes* 2015;8:533.
- 76 Olamoyegun M, Ibraheem W, Iwuala S, *et al.* Burden and pattern of micro vascular complications in type 2 diabetes in a tertiary health institution in Nigeria. *Afr Health Sci* 2015;15:1136–41.
- 77 Onakpoya OH, Kolawole BA, Adeoye AO, *et al.* Visual impairment and blindness in type 2 diabetics: Ife-Ijesa diabetic retinopathy study. *Int Ophthalmol* 2016;36:477–85.
- 78 Oyelade BO, OlaOlorun AD, Odeigah LO, *et al.* The prevalence of peripheral arterial disease in diabetic subjects in south-west Nigeria. *Afr J Prim Health Care Fam Med* 2012;4:354.
- 79 Ugoya SO, Echejoh GO, Ugoya TA, *et al.* Clinically diagnosed diabetic neuropathy: frequency, types and severity. *J Natl Med Assoc* 2006;98:1763–6.
- 80 Unachukwu C, Babatunde S, Ihekwa AE. Diabetes, hand and/or foot ulcers: a cross-sectional hospital-based study in Port Harcourt, Nigeria. *Diabetes Res Clin Pract* 2007;75:148–52.
- 81 Mohamad AH, Hassan A. Correlation between retinopathy, nephropathy and peripheral neuropathy among adult Sudanese diabetic patients. *Sud Jnl Med Sci.* 2011;6.
- 82 Ahmed MH, Elwali ES, Awadalla H, *et al.* The relationship between diabetic retinopathy and nephropathy in Sudanese adult with diabetes: population based study. *Diabetes Metab Syndr* 2017;11 Suppl 1:S333–6.
- 83 Albalawi HB, Mamdouh Alali N, H. ALEnezi S, *et al.* The relationship between periodontitis and diabetic retinopathy: a cross-sectional longitudinal study. *Australasian Medical Journal* 2020;13:50–4.
- 84 Ashur ST, Shah SA, Bosseri S, *et al.* Glycaemic control status among type 2 diabetic patients and the role of their diabetes coping behaviours: a clinic-based study in Tripoli, Libya. *Libyan J Med* 2016;11:31086.
- 85 Assaad-Khalil SH, Zaki A, Abdel Rehim A, *et al.* Prevalence of diabetic foot disorders and related risk factors among Egyptian subjects with diabetes. *Prim Care Diabetes* 2015;9:297–303.
- 86 Khalil SA, Megallaa MH, Rohoma KH, *et al.* Prevalence of chronic diabetic complications in newly diagnosed versus known type 2 diabetic subjects in a sample of Alexandria population, Egypt. *Curr Diabetes Rev* 2019;15:74–83.
- 87 Awadalla H, Noor SK, Elmahoun WM, *et al.* Diabetes complications in Sudanese individuals with type 2 diabetes: overlooked problems in sub-Saharan Africa? *Diabetes Metab Syndr* 2017;11 Suppl 2:S1047–51.
- 88 Bentata Y, Chemlal A, Karimi I, *et al.* Diabetic kidney disease and vascular comorbidities in patients with type 2 diabetes mellitus in a developing country. *Saudi J Kidney Dis Transpl* 2015;26:1035–43.
- 89 Bouaziz A, Zidi I, Zidi N, *et al.* Nephropathy following type 2 diabetes mellitus in Tunisian population. *West Indian Med J* 2012;61:881–9.
- 90 Chadli A, El Aziz S, El Ansari N, *et al.* Management of diabetes in Morocco: results of the International diabetes management practices study (IDMPS) – wave 5. *Ther Adv Endocrinol Metab* 2016;7:101–9.
- 91 Chahbi Z, Lahmar B, Hadri SE, *et al.* The prevalence of painful diabetic neuropathy in 300 Moroccan diabetics. *Pan Afr Med J* 2018;31:158–58.
- 92 Chetoui A, Kaoutar K, Elmoussaoui S, *et al.* Prevalence and determinants of poor glycaemic control: a cross-sectional study among Moroccan type 2 diabetes patients. *Int Health* 2020;14:390–7.
- 93 Diaf M, Khaled BM. Metabolic profile, nutritional status and determinants of glycaemic control in Algerian type 2 diabetic patients. *Kuwait Medical Journal* 2017;49:135–41.
- 94 Elnasri HA, Ahmed AM. Patterns of lipid changes among type 2 diabetes patients in Sudan. *East Mediterr Health J* 2008;14:314–24.
- 95 Elwali ES, Almobarak AO, Hassan MA, *et al.* Frequency of diabetic retinopathy and associated risk factors in Khartoum, Sudan: population based study. *Int J Ophthalmol* 2017;10:948–54.
- 96 Kahloun R, Jelliti B, Zaouali S, *et al.* Prevalence and causes of visual impairment in diabetic patients in Tunisia, North Africa. *Eye* 2014;28:986–91.
- 97 Megallaa MH, Ismail AA, Zeitoun MH, *et al.* Association of diabetic foot ulcers with chronic vascular diabetic complications in patients with type 2 diabetes. *Diabetes Metab Syndr* 2019;13:1287–92.
- 98 Noor SK, Elmahoun WM, Bushara SO, *et al.* Glycaemic control in Sudanese individuals with type 2 diabetes: population based study. *Diabetes Metab Syndr* 2017;11 Suppl 1:S147–51.
- 99 Omar SM, Musa IR, Osman OE, *et al.* Assessment of glycemic control in type 2 diabetes in the eastern Sudan. *BMC Res Notes* 2018;11:373.
- 100 Adeniyi OV, Owolabi EO. Cross-Sectional study of diabetes kidney disease in the eastern Cape, South Africa. *Medicine* 2020;99:e23303.
- 101 Amod A, Riback W, Schoeman HS. Diabetes guidelines and clinical practice: is there a gap? the South African cohort of the International diabetes management practices study. *Journal of Endocrinology, Metabolism and Diabetes of South Africa* 2012;17:85–90.
- 102 Blake AM, Munby HN, Katlego PM, *et al.* Characteristics of patients with diabetic retinopathy in Gaborone, Botswana. *Tanzan J Health Res* 2015;17:thrb.v17i1.
- 103 Burgess PI, Allain TJ, García-Fiñana M, *et al.* High prevalence in Malawi of sight-threatening retinopathy and visual impairment caused by diabetes: identification of population-specific targets for intervention. *Diabet Med* 2014;31:1643–50.
- 104 Cairncross JP, Steinberg WJ, Labuschagne MJ. Prevalence of eye pathology in a group of diabetic patients at national district hospital outpatient department in Bloemfontein, South Africa. *Afr J Prim Health Care Fam Med* 2017;9:e1–7.
- 105 Cohen DB, Allain TJ, Glover S, *et al.* A survey of the management, control, and complications of diabetes mellitus in patients attending a diabetes clinic in Blantyre, Malawi, an area of high HIV prevalence. *Am J Trop Med Hyg* 2010;83:575–81.
- 106 Glover SJ, Burgess PI, Cohen DB, *et al.* Prevalence of diabetic retinopathy, cataract and visual impairment in patients with diabetes in sub-Saharan Africa. *Br J Ophthalmol* 2012;96:156–61.
- 107 Lewis AD, Hogg RE, Chandran M, *et al.* Prevalence of diabetic retinopathy and visual impairment in patients with diabetes mellitus in Zambia through the implementation of a mobile diabetic retinopathy screening project in the Copperbelt Province: a cross-sectional study. *Eye* 2018;32:1201–8.
- 108 Machingura PI, Chikwasha V, Okwanga PN, *et al.* Prevalence of and factors associated with nephropathy in diabetic patients attending an outpatient clinic in Harare, Zimbabwe. *Am J Trop Med Hyg* 2017;96:477–82.
- 109 Makwero MT, Mollentze WF, Joubert G, *et al.* Anthropometric profile and complications in patients with diabetes mellitus seen at Maluti Adventist Hospital, Lesotho. *South African Family Practice* 2018;60:97–102.
- 110 Molefe-Baikai OJ, Molefi M, Cainelli F, *et al.* The prevalence of microalbuminuria and associated factors among patients with type 2 diabetes mellitus in Botswana. *Niger J Clin Pract* 2018;21:1430–7.
- 111 Mwita JC, Francis JM, Omech B, *et al.* Glycaemic, blood pressure and low-density lipoprotein-cholesterol control among patients



- with diabetes mellitus in a specialised clinic in Botswana: a cross-sectional study. *BMJ Open* 2019;9:e026807.
- 112 Pirie FJ, Maharaj S, Esterhuizen TM, *et al*. Retinopathy in subjects with type 2 diabetes at a tertiary diabetes clinic in Durban, South Africa: clinical, biochemical and genetic factors. *J Clin Transl Endocrinol* 2014;1:e9–12.
 - 113 Rotchford AP, Rotchford KM. Diabetes in rural South Africa—an assessment of care and complications. *S Afr Med J* 2002;92:536–41.
 - 114 Thinyane KH, Theketsa CE. Characteristics of patients admitted with diabetes in Maseru, Lesotho. *African Journal of Diabetes Medicine* 2013;21:17–19.
 - 115 Thomas RL, Distiller L, Luzio SD, *et al*. Ethnic differences in the prevalence of diabetic retinopathy in persons with diabetes when first presenting at a diabetes clinic in South Africa. *Diabetes Care* 2013;36:336–41.
 - 116 Webb EM, Rheeder P, Van Zyl DG. Diabetes care and complications in primary care in the Tshwane district of South Africa. *Prim Care Diabetes* 2015;9:147–54.
 - 117 Blum J, Chaney M, Mudji J, *et al*. Glycaemic control among patients with type 2 diabetes followed in a rural African primary care setting - A reality check in the Democratic Republic of Congo. *Prim Care Diabetes* 2020;14:139–46.
 - 118 Dzudie A, Choukem S-PN, Adam AK, *et al*. Prevalence and determinants of electrocardiographic abnormalities in sub-Saharan African individuals with type 2 diabetes. *Cardiovasc J Afr* 2012;23:533–7.
 - 119 Efundem NT, Assob JCN, Fetei VF, *et al*. Prevalence and associations of microalbuminuria in proteinuria-negative patients with type 2 diabetes in two regional hospitals in Cameroon: a cross-sectional study. *BMC Res Notes* 2017;10:477.
 - 120 Hall KK, Tambekou J, Penn L, *et al*. Association between depression, glycaemic control and the prevalence of diabetic retinopathy in a diabetic population in Cameroon. *S Afr J Psychiatry* 2017;23:983–83.
 - 121 Jingi AM, Nansseu JRN, Noubiap JJN, *et al*. Diabetes and visual impairment in sub-Saharan Africa: evidence from Cameroon. *J Diabetes Metab Disord* 2015;14:21.
 - 122 Jingi AM, Noubiap JJN, Ellong A, *et al*. Epidemiology and treatment outcomes of diabetic retinopathy in a diabetic population from Cameroon. *BMC Ophthalmol* 2014;14:19.
 - 123 Njikam EJ, Kariuki MM, Kollmann MKH, *et al*. The magnitude and pattern of diabetic retinopathy in Yaoundé, Cameroon - a cross-sectional hospital-based study. *Acta Ophthalmol* 2016;94:e156–7.
 - 124 Iwuola SO, Olamoyegun MA, Sabir AA, *et al*. The relationship between self-monitoring of blood glucose and glycaemic control among patients attending an urban diabetes clinic in Nigeria. *Ann Afr Med* 2015;14:182–7.
 - 125 Albalawi HB, Mamdouh Alali N, H. ALenezi S, *et al*. The relationship between periodontitis and diabetic retinopathy: a cross-sectional longitudinal study. *Australas Med J* 2020;13:50–4.
 - 126 Ali MK, Bullard KM, Saaddine JB, *et al*. Achievement of goals in U.S. diabetes care, 1999–2010. *N Engl J Med* 2013;368:1613–24.
 - 127 Carls G, Huynh J, Tuttle E, *et al*. Achievement of glycosylated hemoglobin goals in the US remains unchanged through 2014. *Diabetes Ther* 2017;8:863–73.
 - 128 Al Mansari A, Obeid Y, Islam N, *et al*. Goal study: clinical and non-clinical predictive factors for achieving glycemic control in people with type 2 diabetes in real clinical practice. *BMJ Open Diabetes Res Care* 2018;6:e000519.
 - 129 Hu H, Hori A, Nishiura C, *et al*. Hba1C, blood pressure, and lipid control in people with diabetes: Japan epidemiology collaboration on occupational health study. *PLoS One* 2016;11:e0159071.
 - 130 Schmieder RE, Tschöpe D, Koch C, *et al*. Individualised treatment targets in patients with type-2 diabetes and hypertension. *Cardiovasc Diabetol* 2018;17:18.
 - 131 Kumar KVSH, Modi KD. A1C, blood pressure and cholesterol goal achievement in patients of type 2 diabetes. *Med J DY Patil Univ* 2016;9:195–9.
 - 132 Rigato M, Pizzol D, Tiago A, *et al*. Characteristics, prevalence, and outcomes of diabetic foot ulcers in Africa. A systemic review and meta-analysis. *Diabetes Res Clin Pract* 2018;142:63–73.
 - 133 Shiferaw WS, Akalu TY, Work Y, *et al*. Prevalence of diabetic peripheral neuropathy in Africa: a systematic review and meta-analysis. *BMC Endocr Disord* 2020;20:49.
 - 134 Burgess PI, MacCormick IJC, Harding SP, *et al*. Epidemiology of diabetic retinopathy and maculopathy in Africa: a systematic review. *Diabet Med* 2013;30:399–412.
 - 135 Achigbu EO, Agweye CT, Achigbu KI, *et al*. Diabetic retinopathy in sub-Saharan Africa: a review of magnitude and risk factors. *Nigerian Journal of Ophthalmology* 2021;29:3–12.
 - 136 Noubiap JJN, Naidoo J, Kengne AP. Diabetic nephropathy in Africa: a systematic review. *World J Diabetes* 2015;6:759–73.
 - 137 Johnston LE, Stewart BT, Yangni-Angate H, *et al*. Peripheral arterial disease in sub-Saharan Africa: a review. *JAMA Surg* 2016;151:564–72.
 - 138 So W-Y, Raboca J, Sobrepena L, *et al*. Comprehensive risk assessments of diabetic patients from seven Asian countries: the joint Asia diabetes evaluation (JADE) program. *J Diabetes* 2011;3:109–18.
 - 139 Fang M, Selvin E. Thirty-Year trends in complications in U.S. adults with newly diagnosed type 2 diabetes. *Diabetes Care* 2021;44:699–706.
 - 140 Beran D, Yudkin JS, de Courten M. Access to care for patients with insulin-requiring diabetes in developing countries: case studies of Mozambique and Zambia. *Diabetes Care* 2005;28:2136–40.
 - 141 Beran D, Yudkin JS. Looking beyond the issue of access to insulin: what is needed for proper diabetes care in resource poor settings. *Diabetes Res Clin Pract* 2010;88:217–21.
 - 142 Kibirige D, Atuhe D, Kampiire L, *et al*. Access to medicines and diagnostic tests integral in the management of diabetes mellitus and cardiovascular diseases in Uganda: insights from the ACCODAD study. *Int J Equity Health* 2017;16:154.
 - 143 Jingi AM, Noubiap JJN, Ewane Onana A, *et al*. Access to diagnostic tests and essential medicines for cardiovascular diseases and diabetes care: cost, availability and affordability in the West region of Cameroon. *PLoS One* 2014;9:e111812.
 - 144 Mendis S, Al Bashir I, Dissanayake L, *et al*. Gaps in capacity in primary care in low-resource settings for implementation of essential noncommunicable disease interventions. *Int J Hypertens* 2012;2012:1–7.
 - 145 WHO. The Abuja declaration: ten years on, 2011. Available: <https://www.who.int/healthsystems/publications/Abuja10pdf> [Accessed 08 Oct 2021].