

EPE338: USING ROUTINELY COLLECTED BLOOD DONATION DATA FOR EXPANDED HIV AND SYPHILIS SURVEILLANCE IN BLANTYRE DISTRICT, MALAWI

Emmanuel Singogo¹, Thomas Hartney², Sarah Bourdin², Maganizo Chagomerana¹, Sydney Puerto-Meredith^{1,3}, Bridon M'baya⁴, Godfrey Kadewere⁵, Evaristar Kudowa¹, Brian Rice², James Hargreaves², Sharon Weir³, Mina C. Hosseinipour^{1,3}

1. University of North Carolina Project-Lilongwe, Malawi; 2. London School of Hygiene and Tropical Medicine, UK; 3. University of North Carolina-Chapel Hill; Malawi Blood Transfusion Services, Blantyre Malawi; 4. Ministry of Health, Directorate of Health Technical Support Services-Lilongwe, Malawi

Background

- WHO recommends all blood donations be screened for transfusion transmissible infections.
- However, these data are not incorporated into national surveillance systems in Malawi.
- We set out to use **routinely collected data from blood donors in Blantyre district, Malawi**, an area of high HIV and syphilis prevalence, to explore current HIV and syphilis prevalence and identify recent sero-conversions among repeat donors.
- We think these data are an **underutilised source of information** with great potential to contribute to TTI surveillance data sources and subsequent understanding of local variations in HIV disease burden over time.
- Therefore, through this analysis we aim to highlight the feasibility of using blood screening data to improve for HIV and syphilis surveillance.

Results

- A total of **23,280 donations from 5,051 donors** were recorded, with 7 donors (0.1%) donating a maximum of 24 times.
- The majority of donors were male (4,294; 85%) and students (3264; 64.6%) at the time of their first donation.
- Of those screened for HIV and syphilis, 104 (2.1%, 104/5,051) and 206 (4.1%, 206/5,051) tested positive respectively.
- Among repeat donors who previously tested negative, **74 HIV sero-conversions** and **165 syphilis sero-conversions** were identified over the study period, indicating an **HIV incidence rate of 5.89 (4.69, 7.40) per 1,000 person-years** and a **syphilis incidence rate of 13.25 (11.37, 15.43) per 1,000 person-years**.
- Compared to men, female donors had a significant lower risk of any TTI seroconversion (Adjusted RR: 0.66; 0.46, 0.96), and also for both HIV and syphilis in univariate analysis but this significance disappeared when adjusting for other donor covariates such as age, occupation and marital status.
- We noted **differences in geographical distribution of prevalence of HIV and syphilis (Figure 1)**.


Conclusions

- Routinely collected data from national blood donation services may be used to enhance existing population-level disease surveillance systems, particularly in high prevalence areas.
- While blood donors are generally considered a low-risk population for HIV and syphilis, we were able to identify and characterize blood donor populations at increased risk of sero-conversion over the study period.
- This information will provide **insight into priority prevention areas in Blantyre district** and help to inform targeted interventions for improved prevention, testing and treatment.

Methods

- We conducted a **retrospective cohort analysis of blood donation data collected in Blantyre** district by the Malawi Blood Transfusion Service (MBTS) between October 1st 2015 and May 31st 2021.
- All blood donations were routinely screened for WHO-prioritized transfusion-transmissible infections (TTI), including HIV and syphilis.
- We characterized donor demographics as well as screening outcomes, including identifying sero-conversions among repeat donors who previously tested negative.
- Logistic regression was used to model the impact of individual level covariates on the probability of sero-conversion.
- We used geographic information system tools to map the prevalence of both HIV and syphilis in Blantyre (Figure 1)

Figure 1: Prevalence of HIV and Syphilis in Blantyre District



Acknowledgements

We thank Malawi Blood Transfusion Service for providing data for this study. Funding for this study was provided by the MeSH Consortium II at London School of Hygiene and Tropical Medicine (PI: Mina Hosseinipour & Sharon Weir)

